



wwPDB X-ray Structure Validation Summary Report ⓘ

Mar 5, 2026 – 07:57 PM UTC

PDB ID : 4DUF / pdb_00004duf
Title : cytochrome P450 BM3h-2G9 MRI sensor bound to serotonin
Authors : Brustad, E.M.; Lelyveld, V.S.; Snow, C.D.; Crook, N.; Martinez, F.M.; Scholl, T.J.; Jasanoff, A.; Arnold, F.H.
Deposited on : 2012-02-21
Resolution : 1.80 Å (reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0
Mogul : 2022.3.0, CSD as543be (2022)
Xtrriage (Phenix) : 2.0
EDS : 3.0
Buster-report : wwPDB partial adaption of 1.1.7 (2018)
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4 : 9.0.010 (Gargrove)
Density-Fitness : 1.0.12
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.80 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	7662 (1.80-1.80)
Clashscore	190562	8479 (1.80-1.80)
Ramachandran outliers	187476	8391 (1.80-1.80)
Sidechain outliers	187428	8390 (1.80-1.80)
RSRZ outliers	180081	7663 (1.80-1.80)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	471	 85% 11% . .
1	B	471	 83% 11% . .
1	C	471	 79% 15% . .
1	D	471	 85% 12% .

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard

residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
3	SRO	A	501	-	-	X	-
3	SRO	B	501	-	X	X	-
3	SRO	D	501	-	X	-	-

2 Entry composition [i](#)

There are 4 unique types of molecules in this entry. The entry contains 16229 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called cytochrome P450 BM3 variant 2G9.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	456	3733	2380	633	700	20	0	9	0
1	B	456	3737	2383	636	698	20	0	10	0
1	C	456	3732	2379	632	701	20	0	9	0
1	D	456	3704	2363	627	694	20	0	6	0

There are 48 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
A	50	CYS	ARG	engineered mutation	UNP P14779
A	87	LEU	PHE	engineered mutation	UNP P14779
A	268	SER	THR	engineered mutation	UNP P14779
A	438	LEU	THR	engineered mutation	UNP P14779
A	464	LEU	-	expression tag	UNP P14779
A	465	GLU	-	expression tag	UNP P14779
A	466	HIS	-	expression tag	UNP P14779
A	467	HIS	-	expression tag	UNP P14779
A	468	HIS	-	expression tag	UNP P14779
A	469	HIS	-	expression tag	UNP P14779
A	470	HIS	-	expression tag	UNP P14779
A	471	HIS	-	expression tag	UNP P14779
B	50	CYS	ARG	engineered mutation	UNP P14779
B	87	LEU	PHE	engineered mutation	UNP P14779
B	268	SER	THR	engineered mutation	UNP P14779
B	438	LEU	THR	engineered mutation	UNP P14779
B	464	LEU	-	expression tag	UNP P14779
B	465	GLU	-	expression tag	UNP P14779
B	466	HIS	-	expression tag	UNP P14779
B	467	HIS	-	expression tag	UNP P14779
B	468	HIS	-	expression tag	UNP P14779

Continued on next page...

Continued from previous page...


Chain	Residue	Modelled	Actual	Comment	Reference
B	469	HIS	-	expression tag	UNP P14779
B	470	HIS	-	expression tag	UNP P14779
B	471	HIS	-	expression tag	UNP P14779
C	50	CYS	ARG	engineered mutation	UNP P14779
C	87	LEU	PHE	engineered mutation	UNP P14779
C	268	SER	THR	engineered mutation	UNP P14779
C	438	LEU	THR	engineered mutation	UNP P14779
C	464	LEU	-	expression tag	UNP P14779
C	465	GLU	-	expression tag	UNP P14779
C	466	HIS	-	expression tag	UNP P14779
C	467	HIS	-	expression tag	UNP P14779
C	468	HIS	-	expression tag	UNP P14779
C	469	HIS	-	expression tag	UNP P14779
C	470	HIS	-	expression tag	UNP P14779
C	471	HIS	-	expression tag	UNP P14779
D	50	CYS	ARG	engineered mutation	UNP P14779
D	87	LEU	PHE	engineered mutation	UNP P14779
D	268	SER	THR	engineered mutation	UNP P14779
D	438	LEU	THR	engineered mutation	UNP P14779
D	464	LEU	-	expression tag	UNP P14779
D	465	GLU	-	expression tag	UNP P14779
D	466	HIS	-	expression tag	UNP P14779
D	467	HIS	-	expression tag	UNP P14779
D	468	HIS	-	expression tag	UNP P14779
D	469	HIS	-	expression tag	UNP P14779
D	470	HIS	-	expression tag	UNP P14779
D	471	HIS	-	expression tag	UNP P14779

- Molecule 2 is PROTOPORPHYRIN IX CONTAINING FE (CCD ID: HEM) (formula: $C_{34}H_{32}FeN_4O_4$).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	
			Total	C	Fe	N			O
2	A	1	43	34	1	4	4	0	0
2	B	1	43	34	1	4	4	0	0
2	C	1	43	34	1	4	4	0	0
2	D	1	43	34	1	4	4	0	0

- Molecule 3 is SEROTONIN (CCD ID: SRO) (formula: C₁₀H₁₂N₂O).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
3	A	1	Total	C	N	O	0	0
			13	10	2	1		
3	B	1	Total	C	N	O	0	0
			13	10	2	1		
3	C	1	Total	C	N	O	0	0
			13	10	2	1		
3	D	1	Total	C	N	O	0	0
			13	10	2	1		


- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	310	Total	O	0	0
			310	310		
4	B	312	Total	O	0	0
			312	312		
4	C	242	Total	O	0	0
			242	242		
4	D	235	Total	O	0	0
			235	235		


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: cytochrome P450 BM3 variant 2G9



- Molecule 1: cytochrome P450 BM3 variant 2G9





- Molecule 1: cytochrome P450 BM3 variant 2G9





- Molecule 1: cytochrome P450 BM3 variant 2G9



4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	79.69Å 150.49Å 87.71Å 90.00° 90.32° 90.00°	Depositor
Resolution (Å)	29.44 – 1.80 29.44 – 1.80	Depositor EDS
% Data completeness (in resolution range)	98.0 (29.44-1.80) 98.0 (29.44-1.80)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	0.06	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.39 (at 1.80Å)	Xtrriage
Refinement program	REFMAC 5.5.0109	Depositor
R, R_{free}	0.163 , 0.203 0.163 , 0.202	Depositor DCC
R_{free} test set	9607 reflections (5.04%)	wwPDB-VP
Wilson B-factor (Å ²)	20.2	Xtrriage
Anisotropy	0.072	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.37 , 39.9	EDS
L-test for twinning ²	$\langle L \rangle = 0.48$, $\langle L^2 \rangle = 0.31$	Xtrriage
Estimated twinning fraction	0.027 for h,-k,-l	Xtrriage
F_o, F_c correlation	0.96	EDS
Total number of atoms	16229	wwPDB-VP
Average B, all atoms (Å ²)	22.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The analyses of the Patterson function reveals a significant off-origin peak that is 37.96 % of the origin peak, indicating pseudo-translational symmetry. The chance of finding a peak of this or larger height randomly in a structure without pseudo-translational symmetry is equal to 3.9608e-04. The detected translational NCS is most likely also responsible for the elevated intensity ratio.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: SRO, HEM

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	1.40	13/3823 (0.3%)	1.21	14/5167 (0.3%)
1	B	1.43	15/3833 (0.4%)	1.18	5/5180 (0.1%)
1	C	1.42	13/3822 (0.3%)	1.23	17/5165 (0.3%)
1	D	1.39	9/3791 (0.2%)	1.20	13/5125 (0.3%)
All	All	1.41	50/15269 (0.3%)	1.21	49/20637 (0.2%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	C	0	1

The worst 5 of 50 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	D	302	VAL	CA-CB	9.03	1.62	1.54
1	C	441	PRO	C-O	8.80	1.33	1.23
1	B	357	ILE	CA-CB	7.97	1.59	1.53
1	D	385	ILE	CA-CB	6.70	1.60	1.53
1	C	357	ILE	CA-C	6.68	1.58	1.52

The worst 5 of 49 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	436	THR	N-CA-C	-11.89	97.33	108.75
1	A	436	THR	N-CA-C	-9.79	93.94	108.34
1	C	245	THR	N-CA-C	8.38	123.51	113.28
1	A	381	ASN	CA-C-N	7.76	129.54	119.84

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed($^{\circ}$)	Ideal($^{\circ}$)
1	A	381	ASN	C-N-CA	7.76	129.54	119.84

There are no chirality outliers.

All (1) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	C	136	ASP	Peptide

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3733	0	3696	34	0
1	B	3737	0	3709	45	0
1	C	3732	0	3692	54	0
1	D	3704	0	3665	28	0
2	A	43	0	30	1	0
2	B	43	0	30	0	0
2	C	43	0	30	4	0
2	D	43	0	30	3	0
3	A	13	0	11	7	0
3	B	13	0	10	6	0
3	C	13	0	11	1	0
3	D	13	0	11	1	0
4	A	310	0	0	3	0
4	B	312	0	0	6	2
4	C	242	0	0	7	2
4	D	235	0	0	2	0
All	All	16229	0	14925	168	2

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 6.

The worst 5 of 168 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:C:267[A]:GLU:HG2	1:C:438:LEU:HD12	1.41	1.02
1:C:110:GLN:HE22	1:C:113:LYS:NZ	1.60	0.98
1:C:146:THR:HG23	1:C:266:HIS:CE1	2.05	0.92
1:C:146:THR:HG23	1:C:266:HIS:HE1	1.34	0.90
1:C:110:GLN:HE22	1:C:113:LYS:HZ2	0.95	0.89

All (2) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
4:B:618:HOH:O	4:C:807:HOH:O[2_547]	1.15	1.05
4:B:661:HOH:O	4:C:807:HOH:O[2_547]	2.01	0.19

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	463/471 (98%)	451 (97%)	12 (3%)	0	100	100
1	B	464/471 (98%)	452 (97%)	12 (3%)	0	100	100
1	C	463/471 (98%)	452 (98%)	11 (2%)	0	100	100
1	D	460/471 (98%)	445 (97%)	15 (3%)	0	100	100
All	All	1850/1884 (98%)	1800 (97%)	50 (3%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	407/413 (98%)	405 (100%)	2 (0%)	81	80
1	B	408/413 (99%)	399 (98%)	9 (2%)	45	34
1	C	407/413 (98%)	395 (97%)	12 (3%)	37	25
1	D	403/413 (98%)	398 (99%)	5 (1%)	63	57
All	All	1625/1652 (98%)	1597 (98%)	28 (2%)	51	45

5 of 28 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	C	109	GLN
1	D	455	LEU
1	C	218	LYS
1	D	224	LYS
1	C	148	LEU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 28 such sidechains are listed below:

Mol	Chain	Res	Type
1	B	359	GLN
1	D	229	GLN
1	C	70	ASN
1	C	403	GLN
1	C	7	GLN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry

8 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
2	HEM	A	500	3,1	50,50,50	1.77	10 (20%)	67,82,82	1.85	17 (25%)
2	HEM	B	500	3,1	50,50,50	1.80	9 (18%)	67,82,82	2.42	17 (25%)
3	SRO	D	501	2	14,14,14	2.00	3 (21%)	19,19,19	3.79	12 (63%)
2	HEM	C	500	3,1	50,50,50	1.78	11 (22%)	67,82,82	2.21	19 (28%)
3	SRO	A	501	2	14,14,14	2.03	5 (35%)	19,19,19	3.68	9 (47%)
3	SRO	B	501	2	14,14,14	2.39	4 (28%)	19,19,19	4.43	11 (57%)
3	SRO	C	501	2	14,14,14	1.71	1 (7%)	19,19,19	2.78	6 (31%)
2	HEM	D	500	3,1	50,50,50	1.84	15 (30%)	67,82,82	1.81	17 (25%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '2' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	HEM	A	500	3,1	-	2/14/54/54	-
2	HEM	B	500	3,1	-	2/14/54/54	-
3	SRO	D	501	2	-	1/3/3/3	0/2/2/2
2	HEM	C	500	3,1	-	2/14/54/54	-
3	SRO	A	501	2	-	1/3/3/3	0/2/2/2
3	SRO	B	501	2	-	1/3/3/3	0/2/2/2
3	SRO	C	501	2	-	0/3/3/3	0/2/2/2
2	HEM	D	500	3,1	-	2/14/54/54	-

The worst 5 of 58 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	B	501	SRO	CB-CG	-6.54	1.39	1.50

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	A	500	HEM	C3D-C2D	5.87	1.49	1.36
3	D	501	SRO	CD2-CE2	5.76	1.48	1.41
2	B	500	HEM	FE-ND	5.66	2.12	1.94
3	C	501	SRO	CD2-CE2	5.35	1.48	1.41

The worst 5 of 108 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	B	500	HEM	C4D-ND-C1D	9.35	116.28	105.21
3	D	501	SRO	CZ2-CE2-CD2	-8.20	114.25	122.19
3	B	501	SRO	CZ2-CE2-CD2	-7.80	114.64	122.19
3	C	501	SRO	CZ2-CE2-CD2	-7.50	114.93	122.19
3	D	501	SRO	CE3-CD2-CE2	7.49	126.69	119.39

There are no chirality outliers.

5 of 11 torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
3	A	501	SRO	NZ-CA-CB-CG
3	B	501	SRO	NZ-CA-CB-CG
3	D	501	SRO	NZ-CA-CB-CG
2	C	500	HEM	CAD-CBD-CGD-O1D
2	D	500	HEM	CAD-CBD-CGD-O1D


There are no ring outliers.


7 monomers are involved in 22 short contacts:


Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	A	500	HEM	1	0
3	D	501	SRO	1	0
2	C	500	HEM	4	0
3	A	501	SRO	7	0
3	B	501	SRO	6	0
3	C	501	SRO	1	0
2	D	500	HEM	3	0


The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier.

Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.









5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	456/471 (96%)	-0.53	6 (1%) 75 75	6, 19, 42, 58	10 (2%)
1	B	456/471 (96%)	-0.62	3 (0%) 84 85	6, 18, 37, 51	10 (2%)
1	C	456/471 (96%)	-0.29	9 (1%) 65 65	6, 21, 47, 66	9 (1%)
1	D	456/471 (96%)	-0.44	8 (1%) 67 68	9, 20, 43, 58	6 (1%)
All	All	1824/1884 (96%)	-0.47	26 (1%) 73 73	6, 20, 42, 66	35 (1%)

The worst 5 of 26 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	C	457	GLY	4.1
1	A	382	PRO	3.6
1	D	197	ALA	3.4
1	D	245	THR	2.9
1	D	191	ALA	2.8

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)


There are no oligosaccharides in this entry.


6.4 Ligands [i](#)


In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q < 0.9’ lists the number of atoms with occupancy less than 0.9.


Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(\AA^2)	Q<0.9
3	SRO	A	501	13/13	0.85	0.14	14,37,41,41	0
3	SRO	B	501	13/13	0.86	0.14	13,35,40,41	0
3	SRO	D	501	13/13	0.86	0.13	15,35,39,39	0
3	SRO	C	501	13/13	0.89	0.14	17,41,43,43	0
2	HEM	A	500	43/43	0.99	0.04	7,10,14,26	0
2	HEM	B	500	43/43	0.99	0.04	6,9,13,24	0
2	HEM	C	500	43/43	0.99	0.04	7,11,14,26	0
2	HEM	D	500	43/43	0.99	0.04	9,13,16,27	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.









6.5 Other polymers [i](#)

There are no such residues in this entry.