



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 8, 2026 – 05:59 PM UTC

PDB ID : 5DUM / pdb_00005dum
Title : Crystal structure of influenza A virus H5 hemagglutinin globular head in complex with the Fab of antibody 65C6
Authors : Sun, J.; Zuo, T.; Wang, G.; Zhou, P.; Zhang, L.; Wang, X.
Deposited on : 2015-09-19
Resolution : 3.00 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0
Mogul : 2022.3.0, CSD as543be (2022)
Xtriage (Phenix) : 2.0
EDS : 3.0
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4 : 9.0.010 (Gargrove)
Density-Fitness : 1.0.12
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 3.00 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	2672 (3.00-3.00)
Clashscore	190562	2977 (3.00-3.00)
Ramachandran outliers	187476	2877 (3.00-3.00)
Sidechain outliers	187428	2880 (3.00-3.00)
RSRZ outliers	180081	2671 (3.00-3.00)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	H	237	 70% 19% • 8%
2	L	214	 70% 26% • •
3	A	233	 % 61% 21% • 15%

2 Entry composition

There are 4 unique types of molecules in this entry. The entry contains 4922 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called 65C6 Heavy Chain.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	H	219	Total	C	N	O	S	0	1	0
			1694	1085	281	321	7			

- Molecule 2 is a protein called 65C6 Light Chain.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	L	211	Total	C	N	O	S	0	0	0
			1621	1016	275	325	5			

- Molecule 3 is a protein called Hemagglutinin.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
3	A	197	Total	C	N	O	S	0	0	0
			1579	1006	272	295	6			

There are 9 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	42	ALA	-	expression tag	UNP Q1WDM0
A	43	ASP	-	expression tag	UNP Q1WDM0
A	44	PRO	-	expression tag	UNP Q1WDM0
A	269	HIS	-	expression tag	UNP Q1WDM0
A	270	HIS	-	expression tag	UNP Q1WDM0
A	271	HIS	-	expression tag	UNP Q1WDM0
A	272	HIS	-	expression tag	UNP Q1WDM0
A	273	HIS	-	expression tag	UNP Q1WDM0
A	274	HIS	-	expression tag	UNP Q1WDM0

- Molecule 4 is 2-acetamido-2-deoxy-beta-D-glucopyranose (CCD ID: NAG) (formula: C₈H₁₅NO₆).




Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
4	A	1	Total	C	N	O	0	0
			14	8	1	5		
4	A	1	Total	C	N	O	0	0
			14	8	1	5		

3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


• Molecule 1: 65C6 Heavy Chain

Chain H: 




• Molecule 2: 65C6 Light Chain

Chain L: 



• Molecule 3: Hemagglutinin

Chain A: 



4 Data and refinement statistics

Property	Value	Source
Space group	P 2 21 21	Depositor
Cell constants a, b, c, α , β , γ	63.54Å 92.98Å 134.00Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	36.13 – 3.00 36.13 – 3.00	Depositor EDS
% Data completeness (in resolution range)	99.1 (36.13-3.00) 99.0 (36.13-3.00)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$	-	Xtrriage
Refinement program	PHENIX 1.9_1692	Depositor
R, R_{free}	0.215 , 0.269 0.220 , 0.272	Depositor DCC
R_{free} test set	799 reflections (4.85%)	wwPDB-VP
Wilson B-factor (Å ²)	(Not available)	Xtrriage
Anisotropy	(Not available)	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.30 , 17.7	EDS
L-test for twinning ¹	$\langle L \rangle =$ (Not available), $\langle L^2 \rangle =$ (Not available)	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
F_o, F_c correlation	0.92	EDS
Total number of atoms	4922	wwPDB-VP
Average B, all atoms (Å ²)	65.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *(Not available)*

¹Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section:
NAG

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	H	0.71	1/1749 (0.1%)	0.97	2/2385 (0.1%)
2	L	0.64	0/1656	1.01	6/2253 (0.3%)
3	A	0.77	0/1621	0.99	2/2204 (0.1%)
All	All	0.71	1/5026 (0.0%)	0.99	10/6842 (0.1%)

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	H	126	SER	CA-C	6.92	1.57	1.52

All (10) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	A	80	VAL	N-CA-C	9.67	121.92	107.51
2	L	39	MET	CA-C-N	8.50	130.46	119.84
2	L	39	MET	C-N-CA	8.50	130.46	119.84
2	L	29	VAL	N-CA-C	-8.46	104.93	113.47
3	A	51	ILE	N-CA-C	6.51	117.49	108.12
2	L	118	PHE	CA-C-N	6.13	126.70	120.38
2	L	118	PHE	C-N-CA	6.13	126.70	120.38
1	H	52	ASP	CA-C-N	5.76	127.04	119.84
1	H	52	ASP	C-N-CA	5.76	127.04	119.84
2	L	186	TYR	N-CA-C	-5.55	107.05	113.88

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	H	1694	0	1632	37	0
2	L	1621	0	1582	35	0
3	A	1579	0	1533	37	0
4	A	28	0	26	0	0
All	All	4922	0	4773	102	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 11.

All (102) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:H:126:SER:HB3	1:H:127:SER:HA	1.47	0.95
3:A:82:LYS:HE3	3:A:85:PRO:HA	1.58	0.84
3:A:142:SER:OG	3:A:143:PHE:N	2.18	0.77
2:L:37:GLN:HB2	2:L:47:LEU:HD11	1.66	0.75
1:H:103[A]:HIS:HD2	3:A:163:SER:HB3	1.52	0.75
1:H:140:PRO:HG3	1:H:152:LEU:HB3	1.71	0.73
1:H:73:ASP:HB3	1:H:76:ILE:HD11	1.71	0.72
2:L:61:ARG:NH1	2:L:82:ASP:OD2	2.16	0.72
1:H:60:TYR:HB2	1:H:65:GLN:HG3	1.71	0.71
1:H:139:ALA:HB2	1:H:226:GLU:HG3	1.74	0.70
1:H:100:ARG:HB3	1:H:103[A]:HIS:ND1	2.07	0.70
1:H:52:ASP:OD2	1:H:101:ARG:NH1	2.24	0.69
1:H:126:SER:CB	1:H:127:SER:HA	2.18	0.68
2:L:6:GLN:NE2	2:L:86:TYR:O	2.24	0.67
1:H:103[A]:HIS:CD2	3:A:163:SER:HB3	2.29	0.67
2:L:185:ASP:HA	2:L:188:LYS:HD2	1.77	0.66
2:L:149:LYS:HG2	2:L:154:LEU:HD13	1.76	0.66
3:A:175:LEU:HD13	3:A:230:TRP:HB3	1.81	0.63
3:A:152:LYS:HE2	3:A:189:LYS:O	1.99	0.63
1:H:101:ARG:NH1	3:A:161:LYS:HD2	2.15	0.61
1:H:103[A]:HIS:HD2	3:A:163:SER:CB	2.13	0.60
1:H:101:ARG:CZ	3:A:161:LYS:HD2	2.31	0.59
1:H:214:HIS:CE1	1:H:216:PRO:HG2	2.37	0.59

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:H:48:MET:HE1	1:H:81:LEU:HD21	1.85	0.58
3:A:53:ARG:HG3	3:A:54:ASP:H	1.68	0.58
2:L:134:CYS:HB2	2:L:148:TRP:CZ2	2.39	0.56
2:L:184:ALA:O	2:L:185:ASP:HB2	2.04	0.56
1:H:87:LYS:O	1:H:90:ASP:N	2.34	0.56
2:L:182:SER:OG	2:L:184:ALA:O	2.24	0.56
3:A:50:LEU:HD22	3:A:108:ILE:CD1	2.36	0.55
3:A:57:VAL:O	3:A:60:TRP:HB3	2.07	0.55
3:A:125:HIS:CE1	3:A:158:PRO:HG2	2.42	0.54
3:A:129:SER:O	3:A:131:VAL:N	2.40	0.54
1:H:18:LEU:HB2	1:H:86:LEU:HD11	1.90	0.54
1:H:57:TYR:OH	3:A:124:ASP:OD2	2.24	0.54
3:A:50:LEU:HD22	3:A:108:ILE:HD12	1.89	0.54
3:A:79:ILE:HG22	3:A:80:VAL:HG23	1.90	0.53
3:A:54:ASP:HB2	3:A:86:ALA:HB3	1.89	0.53
2:L:184:ALA:HA	2:L:187:GLU:OE2	2.08	0.53
3:A:56:SER:HB3	3:A:85:PRO:HG3	1.90	0.52
2:L:67:SER:O	2:L:69:THR:N	2.39	0.52
3:A:182:ASN:ND2	3:A:186:GLU:OE1	2.42	0.52
2:L:202:SER:OG	2:L:203:SER:N	2.42	0.51
2:L:141:PRO:HB2	2:L:143:GLU:OE1	2.11	0.51
1:H:23:LYS:HB3	1:H:78:THR:HG23	1.92	0.50
2:L:73:LEU:HD12	2:L:74:THR:H	1.77	0.50
3:A:53:ARG:CG	3:A:54:ASP:H	2.23	0.50
3:A:160:ILE:O	3:A:242:GLU:HA	2.11	0.50
1:H:13:LYS:HA	1:H:127:SER:OG	2.11	0.50
3:A:53:ARG:HG3	3:A:54:ASP:N	2.27	0.50
1:H:12:LYS:O	1:H:125:VAL:HA	2.12	0.49
1:H:60:TYR:HE2	1:H:70:ILE:HG13	1.76	0.49
2:L:187:GLU:H	2:L:187:GLU:CD	2.18	0.49
1:H:126:SER:H	1:H:127:SER:HG	1.61	0.49
1:H:174:THR:O	1:H:177:VAL:HG12	2.13	0.48
2:L:143:GLU:O	2:L:198:HIS:HD2	1.97	0.48
2:L:21:LEU:HD12	2:L:73:LEU:HD23	1.96	0.48
2:L:140:TYR:CD2	2:L:141:PRO:HA	2.49	0.48
3:A:61:LEU:O	3:A:146:ASN:ND2	2.39	0.48
3:A:154:ASN:O	3:A:156:THR:N	2.47	0.48
3:A:178:ILE:HB	3:A:198:ILE:HD12	1.96	0.47
3:A:151:ILE:HD12	3:A:190:LEU:HD22	1.97	0.46
1:H:91:THR:HG23	1:H:124:THR:HA	1.97	0.46
1:H:87:LYS:HG2	1:H:89:SER:OG	2.16	0.46

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:H:87:LYS:HG3	1:H:88:ALA:N	2.30	0.46
3:A:182:ASN:HA	3:A:214:ALA:O	2.16	0.46
2:L:128:GLY:C	2:L:183:LYS:HB2	2.42	0.46
2:L:127:SER:OG	2:L:128:GLY:N	2.47	0.45
3:A:183:ASP:OD1	3:A:185:ALA:HB3	2.16	0.45
1:H:13:LYS:HG2	1:H:127:SER:HB3	1.98	0.45
3:A:183:ASP:CG	3:A:185:ALA:HB3	2.42	0.45
3:A:182:ASN:HB2	3:A:215:THR:HA	1.98	0.45
2:L:30:SER:O	2:L:30:SER:OG	2.22	0.45
2:L:39:MET:O	2:L:42:GLN:HB2	2.17	0.45
3:A:130:GLY:HA3	3:A:149:TRP:HB3	1.98	0.44
3:A:64:ASN:HB3	3:A:67:CYS:SG	2.57	0.44
1:H:73:ASP:OD1	1:H:75:SER:HB3	2.17	0.44
2:L:94:TRP:HA	2:L:95:PRO:C	2.43	0.44
2:L:151:ASP:C	2:L:153:ALA:H	2.26	0.43
3:A:64:ASN:ND2	3:A:90:CYS:SG	2.91	0.43
3:A:198:ILE:HD11	3:A:247:PHE:HA	2.00	0.43
1:H:87:LYS:O	1:H:89:SER:N	2.51	0.43
2:L:78:LEU:HD13	2:L:78:LEU:HA	1.88	0.43
1:H:25:PHE:HZ	1:H:79:VAL:HG23	1.84	0.43
2:L:128:GLY:HA2	2:L:183:LYS:HD3	2.00	0.43
2:L:167:ASP:OD2	2:L:168:SER:N	2.51	0.43
1:H:126:SER:N	1:H:127:SER:OG	2.40	0.42
3:A:175:LEU:CD1	3:A:230:TRP:HB3	2.48	0.42
2:L:124:GLN:O	2:L:127:SER:HB3	2.19	0.42
1:H:76:ILE:H	1:H:76:ILE:HG13	1.59	0.42
1:H:18:LEU:CB	1:H:86:LEU:HD11	2.50	0.42
2:L:198:HIS:CG	2:L:199:GLN:N	2.88	0.42
3:A:256:ILE:HD13	3:A:256:ILE:HA	1.83	0.42
1:H:223:LYS:HE2	2:L:123:GLU:OE1	2.20	0.41
2:L:7:SER:HA	2:L:8:PRO:HA	1.86	0.41
2:L:25:ALA:HB3	2:L:69:THR:HG23	2.02	0.41
2:L:187:GLU:O	2:L:211:ARG:NH2	2.52	0.41
2:L:124:GLN:HG2	2:L:129:THR:O	2.20	0.41
2:L:118:PHE:HA	2:L:119:PRO:HD3	1.69	0.41
1:H:184:LEU:HG	1:H:190:TYR:CE1	2.57	0.40
1:H:38:ARG:NH2	1:H:46:GLU:OE2	2.41	0.40
2:L:185:ASP:N	2:L:187:GLU:OE2	2.55	0.40

There are no symmetry-related clashes.

5.3 Torsion angles

5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	H	216/237 (91%)	188 (87%)	21 (10%)	7 (3%)	3	18
2	L	209/214 (98%)	189 (90%)	14 (7%)	6 (3%)	3	20
3	A	193/233 (83%)	166 (86%)	21 (11%)	6 (3%)	3	19
All	All	618/684 (90%)	543 (88%)	56 (9%)	19 (3%)	3	19

All (19) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	H	226	GLU
2	L	127	SER
3	A	86	ALA
1	H	88	ALA
1	H	201	SER
2	L	68	GLY
3	A	129	SER
3	A	130	GLY
1	H	31	TYR
1	H	227	SER
2	L	128	GLY
2	L	170	ASP
3	A	90	CYS
3	A	134	ALA
3	A	155	ASN
1	H	75	SER
1	H	205	THR
2	L	17	GLU
2	L	40	PRO

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	H	189/204 (93%)	179 (95%)	10 (5%)	20	54
2	L	183/186 (98%)	176 (96%)	7 (4%)	29	63
3	A	176/208 (85%)	170 (97%)	6 (3%)	32	66
All	All	548/598 (92%)	525 (96%)	23 (4%)	26	61

All (23) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	H	5	VAL
1	H	21	SER
1	H	28	SER
1	H	56	SER
1	H	65	GLN
1	H	76	ILE
1	H	84	SER
1	H	87	LYS
1	H	115	GLU
1	H	130	THR
2	L	9	LEU
2	L	11	LEU
2	L	72	THR
2	L	100	GLN
2	L	126	LYS
2	L	187	GLU
2	L	191	VAL
3	A	104	LEU
3	A	135	CYS
3	A	172	LEU
3	A	178	ILE
3	A	220	ASN
3	A	256	ILE

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (10) such sidechains are listed below:

Mol	Chain	Res	Type
1	H	39	GLN
1	H	82	GLN
2	L	38	GLN
2	L	42	GLN
2	L	124	GLN
2	L	147	GLN
3	A	96	ASN
3	A	109	ASN
3	A	155	ASN
3	A	168	ASN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

2 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
4	NAG	A	301	3	14,14,15	0.47	0	17,19,21	0.87	1 (5%)
4	NAG	A	302	3	14,14,15	1.05	2 (14%)	17,19,21	0.84	1 (5%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral

centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	NAG	A	301	3	-	2/6/23/26	0/1/1/1
4	NAG	A	302	3	-	0/6/23/26	0/1/1/1

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
4	A	302	NAG	O5-C1	2.65	1.48	1.43
4	A	302	NAG	C1-C2	2.59	1.55	1.52

All (2) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
4	A	302	NAG	C1-O5-C5	2.82	115.97	112.19
4	A	301	NAG	C1-C2-N2	2.21	113.92	110.43

There are no chirality outliers.

All (2) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
4	A	301	NAG	C4-C5-C6-O6
4	A	301	NAG	O5-C5-C6-O6

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	H	219/237 (92%)	-0.18	1 (0%) 87 72	31, 61, 83, 88	1 (0%)
2	L	211/214 (98%)	-0.18	0 100 100	52, 68, 79, 88	0
3	A	197/233 (84%)	-0.16	2 (1%) 79 59	50, 66, 92, 96	0
All	All	627/684 (91%)	-0.18	3 (0%) 87 72	31, 65, 86, 96	1 (0%)

All (3) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
3	A	49	PRO	2.8
3	A	50	LEU	2.4
1	H	26	ALA	2.1

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
4	NAG	A	301	14/15	0.45	0.13	64,64,64,64	0

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
4	NAG	A	302	14/15	0.78	0.12	64,64,64,64	0

6.5 Other polymers [i](#)

There are no such residues in this entry.