



wwPDB X-ray Structure Validation Summary Report ⓘ

Mar 15, 2026 – 01:34 PM UTC

PDB ID : 6DV9 / pdb_00006dv9
Title : Crystal structure of Mycobacterium tuberculosis transcription initiation complex (ECF sigma factor L) containing 5nt RNA with 4nt spacer
Authors : Lin, W.; Das, K.; Feng, Y.; Ebright, R.H.
Deposited on : 2018-06-23
Resolution : 3.80 Å (reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 3.80 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.




Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	1065 (3.96-3.64)
Clashscore	190562	1012 (3.94-3.66)
Ramachandran outliers	187476	1048 (3.96-3.64)
Sidechain outliers	187428	1043 (3.96-3.64)
RSRZ outliers	180081	1064 (3.96-3.64)
RNA backbone	3983	1007 (4.50-3.10)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	359	49% 13% . 37%
1	B	359	46% 18% . 35%
2	C	1178	79% 16% . .
3	D	1316	78% 17% .

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
4	E	110	
5	F	177	
6	I	5	
7	G	17	
8	H	22	

2 Entry composition

There are 10 unique types of molecules in this entry. The entry contains 24958 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called DNA-directed RNA polymerase subunit alpha.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	225	Total	C	N	O	S	0	0	0
			1716	1080	296	338	2			
1	B	233	Total	C	N	O	S	0	0	0
			1733	1094	297	340	2			

There are 24 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	-11	MET	-	initiating methionine	UNP P9WGZ1
A	-10	GLY	-	expression tag	UNP P9WGZ1
A	-9	HIS	-	expression tag	UNP P9WGZ1
A	-8	HIS	-	expression tag	UNP P9WGZ1
A	-7	HIS	-	expression tag	UNP P9WGZ1
A	-6	HIS	-	expression tag	UNP P9WGZ1
A	-5	HIS	-	expression tag	UNP P9WGZ1
A	-4	HIS	-	expression tag	UNP P9WGZ1
A	-3	HIS	-	expression tag	UNP P9WGZ1
A	-2	HIS	-	expression tag	UNP P9WGZ1
A	-1	HIS	-	expression tag	UNP P9WGZ1
A	0	HIS	-	expression tag	UNP P9WGZ1
B	-11	MET	-	initiating methionine	UNP P9WGZ1
B	-10	GLY	-	expression tag	UNP P9WGZ1
B	-9	HIS	-	expression tag	UNP P9WGZ1
B	-8	HIS	-	expression tag	UNP P9WGZ1
B	-7	HIS	-	expression tag	UNP P9WGZ1
B	-6	HIS	-	expression tag	UNP P9WGZ1
B	-5	HIS	-	expression tag	UNP P9WGZ1
B	-4	HIS	-	expression tag	UNP P9WGZ1
B	-3	HIS	-	expression tag	UNP P9WGZ1
B	-2	HIS	-	expression tag	UNP P9WGZ1
B	-1	HIS	-	expression tag	UNP P9WGZ1
B	0	HIS	-	expression tag	UNP P9WGZ1

- Molecule 2 is a protein called DNA-directed RNA polymerase subunit beta.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	C	1126	Total	C	N	O	S	0	0	0
			8728	5461	1531	1697	39			

- Molecule 3 is a protein called DNA-directed RNA polymerase subunit beta'.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
3	D	1265	Total	C	N	O	S	0	0	0
			9895	6195	1794	1866	40			

- Molecule 4 is a protein called DNA-directed RNA polymerase subunit omega.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
4	E	81	Total	C	N	O		0	0	0
			630	403	106	121				

- Molecule 5 is a protein called ECF RNA polymerase sigma factor SigL.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
5	F	174	Total	C	N	O	S	0	0	0
			1349	839	256	252	2			

- Molecule 6 is a RNA chain called RNA (5'-R(*CP*UP*CP*GP*A)-3').

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
6	I	5	Total	C	N	O	P	0	0	0
			102	47	18	33	4			

- Molecule 7 is a DNA chain called DNA (5'-D(*GP*CP*AP*TP*CP*CP*GP*TP*GP*AP*GP*TP*CP*GP*AP*GP*G)-3').

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
7	G	17	Total	C	N	O	P	0	0	0
			350	166	68	100	16			

- Molecule 8 is a DNA chain called DNA (5'-D(P*CP*GP*TP*GP*TP*CP*AP*GP*AP*GP*TP*GP*TP*CP*AP*CP*GP*GP*AP*TP*GP*C)-3').

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
8	H	22	Total	C	N	O	P	0	0	0
			452	215	85	131	21			

- Molecule 9 is MAGNESIUM ION (CCD ID: MG) (formula: Mg).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
9	D	1	Total	Mg	0	0
			1	1		


- Molecule 10 is ZINC ION (CCD ID: ZN) (formula: Zn).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
10	D	2	Total	Zn	0	0
			2	2		


3 Residue-property plots

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


• Molecule 1: DNA-directed RNA polymerase subunit alpha




• Molecule 1: DNA-directed RNA polymerase subunit alpha




• Molecule 2: DNA-directed RNA polymerase subunit beta







• Molecule 3: DNA-directed RNA polymerase subunit beta'

Chain D: 78% 17% .







- Molecule 4: DNA-directed RNA polymerase subunit omega




- Molecule 5: ECF RNA polymerase sigma factor SigL



- Molecule 6: RNA (5'-R(*CP*UP*CP*GP*A)-3')



- Molecule 7: DNA (5'-D(*GP*CP*AP*TP*CP*CP*GP*TP*GP*AP*GP*TP*CP*GP*AP*GP*G)-3')



- Molecule 8: DNA (5'-D(P*CP*GP*TP*GP*TP*CP*AP*GP*AP*GP*TP*GP*TP*CP*AP*CP*GP*GP*AP*TP*GP*C)-3')



4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	143.27Å 161.44Å 237.69Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	91.47 – 3.80 91.47 – 3.80	Depositor EDS
% Data completeness (in resolution range)	93.6 (91.47-3.80) 95.5 (91.47-3.80)	Depositor EDS
R_{merge}	0.33	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.44 (at 3.78Å)	Xtriage
Refinement program	PHENIX 1.12_2829	Depositor
R, R_{free}	0.184 , 0.232 0.187 , 0.232	Depositor DCC
R_{free} test set	2618 reflections (4.76%)	wwPDB-VP
Wilson B-factor (Å ²)	97.4	Xtriage
Anisotropy	0.626	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.32 , 92.4	EDS
L-test for twinning ²	$\langle L \rangle = 0.45$, $\langle L^2 \rangle = 0.28$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.92	EDS
Total number of atoms	24958	wwPDB-VP
Average B, all atoms (Å ²)	110.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 2.23% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: ZN, MG

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.11	0/1742	0.30	0/2370
1	B	0.12	0/1759	0.34	0/2399
2	C	0.10	0/8887	0.31	0/12048
3	D	0.10	0/10061	0.30	0/13600
4	E	0.11	0/643	0.30	0/877
5	F	0.13	0/1371	0.32	0/1865
6	I	0.07	0/113	0.18	0/174
7	G	0.18	0/393	0.40	0/606
8	H	0.21	0/507	0.46	0/782
All	All	0.11	0/25476	0.31	0/34721

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1716	0	1756	36	0
1	B	1733	0	1752	47	0
2	C	8728	0	8655	123	0
3	D	9895	0	9953	156	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
4	E	630	0	622	11	0
5	F	1349	0	1344	24	0
6	I	102	0	55	2	0
7	G	350	0	192	10	0
8	H	452	0	249	10	0
9	D	1	0	0	0	0
10	D	2	0	0	0	0
All	All	24958	0	24578	362	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 7.

The worst 5 of 362 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:40:ARG:HE	1:B:33:THR:HG22	1.31	0.95
3:D:334:ARG:HD3	5:F:90:VAL:HG21	1.54	0.90
2:C:1024:THR:H	3:D:730:THR:HG21	1.48	0.79
2:C:593:MET:HA	2:C:628:THR:HG21	1.66	0.77
1:A:4:SER:HB3	1:B:144:ARG:HH12	1.51	0.74

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	223/359 (62%)	217 (97%)	6 (3%)	0	100	100
1	B	231/359 (64%)	212 (92%)	17 (7%)	2 (1%)	14	45
2	C	1124/1178 (95%)	1069 (95%)	50 (4%)	5 (0%)	30	62
3	D	1261/1316 (96%)	1202 (95%)	57 (4%)	2 (0%)	43	73

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
4	E	79/110 (72%)	75 (95%)	4 (5%)	0	100	100
5	F	172/177 (97%)	168 (98%)	3 (2%)	1 (1%)	21	54
All	All	3090/3499 (88%)	2943 (95%)	137 (4%)	10 (0%)	36	67

5 of 10 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	B	6	ARG
2	C	370	ILE
1	B	159	ILE
2	C	732	GLU
3	D	593	PRO

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	194/308 (63%)	191 (98%)	3 (2%)	57	69
1	B	190/308 (62%)	184 (97%)	6 (3%)	34	56
2	C	951/998 (95%)	935 (98%)	16 (2%)	53	67
3	D	1050/1095 (96%)	1039 (99%)	11 (1%)	68	74
4	E	66/90 (73%)	66 (100%)	0	100	100
5	F	133/136 (98%)	127 (96%)	6 (4%)	24	48
All	All	2584/2935 (88%)	2542 (98%)	42 (2%)	55	68

5 of 42 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
3	D	431	VAL
3	D	1194	VAL
3	D	535	ASP
3	D	910	LEU
5	F	56	GLU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 18 such sidechains are listed below:

Mol	Chain	Res	Type
3	D	882	GLN
5	F	122	GLN
3	D	1109	GLN
2	C	700	GLN
3	D	767	HIS

5.3.3 RNA ⓘ

Mol	Chain	Analysed	Backbone Outliers	Pucker Outliers
6	I	4/5 (80%)	1 (25%)	0

All (1) RNA backbone outliers are listed below:

Mol	Chain	Res	Type
6	I	5	C

There are no RNA pucker outliers to report.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no oligosaccharides in this entry.

5.6 Ligand geometry ⓘ

Of 3 ligands modelled in this entry, 3 are monoatomic - leaving 0 for Mogul analysis.

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	225/359 (62%)	-0.44	0 100 100	54, 84, 147, 174	0
1	B	233/359 (64%)	-0.28	1 (0%) 88 74	62, 104, 156, 206	0
2	C	1126/1178 (95%)	-0.36	3 (0%) 90 78	51, 95, 169, 212	0
3	D	1265/1316 (96%)	-0.40	0 100 100	48, 101, 187, 239	0
4	E	81/110 (73%)	-0.47	0 100 100	95, 121, 191, 210	0
5	F	174/177 (98%)	-0.35	0 100 100	62, 115, 182, 225	0
6	I	5/5 (100%)	-0.13	0 100 100	120, 131, 144, 156	0
7	G	17/17 (100%)	0.05	0 100 100	121, 141, 179, 180	0
8	H	22/22 (100%)	0.07	0 100 100	135, 177, 224, 228	0
All	All	3148/3543 (88%)	-0.37	4 (0%) 92 87	48, 100, 178, 239	0

All (4) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
2	C	979	GLY	2.6
2	C	492	PRO	2.6
1	B	89	GLU	2.2
2	C	325	GLY	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
9	MG	D	1401	1/1	0.99	0.04	90,90,90,90	0
10	ZN	D	1402	1/1	1.00	0.06	184,184,184,184	0
10	ZN	D	1403	1/1	1.00	0.04	137,137,137,137	0

6.5 Other polymers [i](#)

There are no such residues in this entry.