



Full wwPDB X-ray Structure Validation Report ⓘ

Apr 25, 2026 – 08:59 PM EDT

PDB ID : 7DVL / pdb_00007dvl
Title : Crystal Structure of the Catalytic Domain of Botulinum Neurotoxin Subtype A3
Authors : Wu, Y.; Leka, O.; Kammerer, R.; Li, X.
Deposited on : 2021-01-13
Resolution : 2.01 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0
Xtriage (Phenix) : 2.0
EDS : 3.0
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4 : 9.0.010 (Gargrove)
Density-Fitness : 1.0.12
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.01 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	10052 (2.00-2.00)
Clashscore	190562	11152 (2.00-2.00)
Ramachandran outliers	187476	11031 (2.00-2.00)
Sidechain outliers	187428	11029 (2.00-2.00)
RSRZ outliers	180081	10067 (2.00-2.00)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	419	9% 94% • •
1	B	419	9% 94% • •

2 Entry composition

There are 3 unique types of molecules in this entry. The entry contains 13486 atoms, of which 6408 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Bont/A3.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
1	A	405	Total	C	H	N	O	S	0	2	0
			6481	2102	3203	551	618	7			
1	B	407	Total	C	H	N	O	S	0	0	0
			6481	2102	3205	549	618	7			

There are 8 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	1	GLY	-	expression tag	UNP D3IV24
A	2	SER	-	expression tag	UNP D3IV24
A	226	GLN	GLU	engineered mutation	UNP D3IV24
A	368	PHE	TYR	engineered mutation	UNP D3IV24
B	1	GLY	-	expression tag	UNP D3IV24
B	2	SER	-	expression tag	UNP D3IV24
B	226	GLN	GLU	engineered mutation	UNP D3IV24
B	368	PHE	TYR	engineered mutation	UNP D3IV24



- Molecule 2 is ZINC ION (CCD ID: ZN) (formula: Zn).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	A	1	Total	Zn	0	0
			1	1		
2	B	1	Total	Zn	0	0
			1	1		

- Molecule 3 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	264	Total	O	0	0
			264	264		
3	B	258	Total	O	0	0
			258	258		

- Molecule 1: Bont/A3



4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	66.25Å 97.42Å 141.25Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	48.71 – 2.01 48.71 – 2.01	Depositor EDS
% Data completeness (in resolution range)	96.6 (48.71-2.01) 96.9 (48.71-2.01)	Depositor EDS
R_{merge}	0.14	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.06 (at 2.00Å)	Xtriage
Refinement program	PHENIX (1.15.2_3472: ???)	Depositor
R, R_{free}	0.205 , 0.232 0.206 , 0.234	Depositor DCC
R_{free} test set	3003 reflections (5.00%)	wwPDB-VP
Wilson B-factor (Å ²)	22.9	Xtriage
Anisotropy	0.527	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.34 , 36.1	EDS
L-test for twinning ²	$\langle L \rangle = 0.50$, $\langle L^2 \rangle = 0.33$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.93	EDS
Total number of atoms	13486	wwPDB-VP
Average B, all atoms (Å ²)	37.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.14% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: ZN

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.14	0/3359	0.38	0/4550
1	B	0.14	0/3354	0.36	0/4544
All	All	0.14	0/6713	0.37	0/9094

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3278	3203	3204	8	0
1	B	3276	3205	3204	8	0
2	A	1	0	0	0	0
2	B	1	0	0	0	0
3	A	264	0	0	2	0
3	B	258	0	0	4	0
All	All	7078	6408	6408	16	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 1.

All (16) close contacts within the same asymmetric unit are listed below, sorted by their clash

magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:13:ARG:NH2	3:B:601:HOH:O	2.22	0.72
1:B:213:GLY:N	3:B:604:HOH:O	2.26	0.68
1:B:409:ARG:NH1	3:B:602:HOH:O	2.37	0.57
1:A:258:LEU:HD22	1:A:371:PHE:HZ	1.73	0.53
1:B:315:MET:SD	1:B:318:ILE:HD12	2.50	0.52
1:A:87:ASP:OD2	1:A:91:LYS:HE2	2.14	0.48
1:A:28:ASN:ND2	3:A:612:HOH:O	2.48	0.46
1:B:183:GLY:HA2	1:B:233:ARG:O	2.15	0.46
1:A:60:ASP:OD2	1:A:62:ASN:ND2	2.49	0.45
1:A:28:ASN:OD1	1:A:134:THR:HG22	2.16	0.44
1:A:115:LYS:NZ	3:A:616:HOH:O	2.50	0.44
1:B:225:HIS:CE1	1:B:264:GLU:OE2	2.71	0.43
1:A:392:PHE:C	1:A:394:LEU:HD12	2.45	0.41
1:B:173:ASP:OD1	1:B:173:ASP:N	2.53	0.41
1:A:272:ASP:OD2	1:A:368:PHE:HD2	2.04	0.40
1:B:316:LYS:NZ	3:B:611:HOH:O	2.44	0.40

There are no symmetry-related clashes.

5.3 Torsion angles ⓘ

5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	403/419 (96%)	395 (98%)	8 (2%)	0	100	100
1	B	403/419 (96%)	392 (97%)	11 (3%)	0	100	100
All	All	806/838 (96%)	787 (98%)	19 (2%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	359/367 (98%)	358 (100%)	1 (0%)	86	91
1	B	359/367 (98%)	358 (100%)	1 (0%)	86	91
All	All	718/734 (98%)	716 (100%)	2 (0%)	86	91

All (2) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	394	LEU
1	B	201	SER

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (5) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	62	ASN
1	A	88	ASN
1	B	9	GLN
1	B	172	HIS
1	B	271	ASN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry

Of 2 ligands modelled in this entry, 2 are monoatomic - leaving 0 for Mogul analysis.

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers

There are no such residues in this entry.

5.8 Polymer linkage issues

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	405/419 (96%)	0.50	39 (9%) 13 12	17, 34, 74, 108	2 (0%)
1	B	407/419 (97%)	0.54	37 (9%) 15 13	20, 35, 68, 101	0
All	All	812/838 (96%)	0.52	76 (9%) 14 13	17, 34, 72, 108	2 (0%)

All (76) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	202	LEU	7.1
1	A	369	LEU	6.7
1	A	418	THR	6.0
1	A	371	PHE	5.7
1	B	146	TYR	5.6
1	A	200	GLU	5.5
1	A	251	ALA	5.2
1	A	370	ASN	5.2
1	A	252	TYR	4.9
1	B	30	GLY	4.8
1	B	213	GLY	4.7
1	A	67	ALA	4.7
1	A	255	MET	4.6
1	A	213	GLY	4.4
1	A	368	PHE	4.3
1	A	30	GLY	4.2
1	B	60	ASP	3.8
1	B	395	GLU	3.8
1	A	31	GLN	3.8
1	A	29	ALA	3.7
1	A	256	SER	3.7
1	A	133	ASP	3.7
1	B	329	ALA	3.7
1	A	257	GLY	3.6

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	B	142	PRO	3.5
1	B	418	THR	3.5
1	B	256	SER	3.4
1	B	124	THR	3.4
1	B	248	LYS	3.2
1	B	145	SER	3.2
1	A	28	ASN	3.1
1	B	140	ILE	3.1
1	B	144	GLY	3.0
1	B	65	PRO	3.0
1	A	405	GLU	3.0
1	A	66	GLU	3.0
1	B	141	GLU	3.0
1	A	250	ASN	2.9
1	B	62	ASN	2.9
1	A	254	GLU	2.8
1	A	7	ASN	2.8
1	B	143	GLY	2.8
1	A	214	THR	2.8
1	A	68	LYS	2.8
1	B	201	SER	2.7
1	A	57	GLU	2.7
1	B	214	THR	2.7
1	B	255	MET	2.7
1	B	249	THR	2.6
1	B	331	GLY	2.6
1	B	251	ALA	2.6
1	B	397	ALA	2.6
1	A	56	PRO	2.6
1	A	258	LEU	2.6
1	A	144	GLY	2.5
1	B	3	MET	2.5
1	B	126	ASP	2.4
1	A	2	SER	2.4
1	A	27	PRO	2.4
1	A	65	PRO	2.4
1	B	7	ASN	2.3
1	B	250	ASN	2.3
1	B	258	LEU	2.3
1	A	253	TYR	2.3
1	A	249	THR	2.2
1	A	389	ASN	2.2

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	394	LEU	2.2
1	A	26	ILE	2.2
1	A	63	PRO	2.2
1	B	133	ASP	2.2
1	A	365[A]	ARG	2.2
1	B	127	THR	2.1
1	B	172	HIS	2.1
1	B	409	ARG	2.1
1	B	366	LYS	2.1
1	B	59	GLY	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
2	ZN	A	501	1/1	0.99	0.03	25,25,25,25	0
2	ZN	B	501	1/1	0.99	0.08	34,34,34,34	0

6.5 Other polymers [i](#)

There are no such residues in this entry.