



Full wwPDB NMR Structure Validation Report ⓘ

Mar 6, 2026 – 03:37 PM UTC

PDB ID : 7DVM / pdb_00007dvm
BMRB ID : 36407
Title : DgkA structure in E.coli lipid bilayer
Authors : Li, J.; Yang, J.
Deposited on : 2021-01-13

This is a Full wwPDB NMR Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/NMRValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0
Mogul : 2022.3.0, CSD as543be (2022)
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
wwPDB-RCI : v_1n_11_5_13_A (Berjanski et al., 2005)
PANAV : Wang et al. (2010)
wwPDB-ShiftChecker : v1.2
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

SOLID-STATE NMR

The overall completeness of chemical shifts assignment is 11%.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	NMR archive (#Entries)
Clashscore	229148	14424
Ramachandran outliers	224038	12848
Sidechain outliers	223484	12823

The table below summarises the geometric issues observed across the polymeric chains and their fit to the experimental data. The red, orange, yellow and green segments indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A cyan segment indicates the fraction of residues that are not part of the well-defined cores, and a grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$

Mol	Chain	Length	Quality of chain
1	A	131	75% 17% 8%
1	B	131	75% 17% 8%
1	C	131	75% 17% 8%

2 Ensemble composition and analysis

This entry contains 10 models. Model 1 is the overall representative, medoid model (most similar to other models).

The following residues are included in the computation of the global validation metrics.

Well-defined (core) protein residues			
Well-defined core	Residue range (total)	Backbone RMSD (Å)	Medoid model
1	A:17-A:23, A:25-A:28, A:30-A:45, A:47-A:50, A:52-A:78, A:80-A:115, A:117-A:121, B:17-B:23, B:25-B:28, B:30-B:45, B:47-B:50, B:52-B:78, B:80-B:115, B:117-B:121, C:17-C:23, C:25-C:28, C:30- C:45, C:47-C:50, C:52-C:78, C:80-C:115, C:117-C:121 (297)	1.73	1

Ill-defined regions of proteins are excluded from the global statistics.

Ligands and non-protein polymers are included in the analysis.

The models can be grouped into 2 clusters and 1 single-model cluster was found.

Cluster number	Models
1	1, 3, 4, 5, 6, 7
2	2, 9, 10
Single-model clusters	8

3 Entry composition

There is only 1 type of molecule in this entry. The entry contains 6183 atoms, of which 3153 are hydrogens and 0 are deuteriums.

- Molecule 1 is a protein called Diacylglycerol kinase.

Mol	Chain	Residues	Atoms						Trace
1	A	121	Total	C	H	N	O	S	0
			2061	659	1051	164	168	19	
1	B	121	Total	C	H	N	O	S	0
			2061	659	1051	164	168	19	
1	C	121	Total	C	H	N	O	S	0
			2061	659	1051	164	168	19	

There are 60 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	-9	MET	-	initiating methionine	UNP A0A0H3MWQ1
A	-8	GLY	-	expression tag	UNP A0A0H3MWQ1
A	-7	HIS	-	expression tag	UNP A0A0H3MWQ1
A	-6	HIS	-	expression tag	UNP A0A0H3MWQ1
A	-5	HIS	-	expression tag	UNP A0A0H3MWQ1
A	-4	HIS	-	expression tag	UNP A0A0H3MWQ1
A	-3	HIS	-	expression tag	UNP A0A0H3MWQ1
A	-2	HIS	-	expression tag	UNP A0A0H3MWQ1
A	-1	GLU	-	expression tag	UNP A0A0H3MWQ1
A	0	LEU	-	expression tag	UNP A0A0H3MWQ1
A	14	R1A	ALA	engineered mutation	UNP A0A0H3MWQ1
A	16	R1A	TYR	engineered mutation	UNP A0A0H3MWQ1
A	24	R1A	ALA	engineered mutation	UNP A0A0H3MWQ1
A	29	R1A	ALA	engineered mutation	UNP A0A0H3MWQ1
A	51	R1A	ASP	engineered mutation	UNP A0A0H3MWQ1
A	70	LEU	ILE	engineered mutation	UNP A0A0H3MWQ1
A	79	R1A	VAL	engineered mutation	UNP A0A0H3MWQ1
A	107	ASP	VAL	engineered mutation	UNP A0A0H3MWQ1
A	113	ALA	CYS	engineered mutation	UNP A0A0H3MWQ1
A	116	R1A	LEU	engineered mutation	UNP A0A0H3MWQ1
B	-9	MET	-	initiating methionine	UNP A0A0H3MWQ1
B	-8	GLY	-	expression tag	UNP A0A0H3MWQ1
B	-7	HIS	-	expression tag	UNP A0A0H3MWQ1
B	-6	HIS	-	expression tag	UNP A0A0H3MWQ1
B	-5	HIS	-	expression tag	UNP A0A0H3MWQ1
B	-4	HIS	-	expression tag	UNP A0A0H3MWQ1
B	-3	HIS	-	expression tag	UNP A0A0H3MWQ1

Continued on next page...

Continued from previous page...


Chain	Residue	Modelled	Actual	Comment	Reference
B	-2	HIS	-	expression tag	UNP A0A0H3MWQ1
B	-1	GLU	-	expression tag	UNP A0A0H3MWQ1
B	0	LEU	-	expression tag	UNP A0A0H3MWQ1
B	14	R1A	ALA	engineered mutation	UNP A0A0H3MWQ1
B	16	R1A	TYR	engineered mutation	UNP A0A0H3MWQ1
B	24	R1A	ALA	engineered mutation	UNP A0A0H3MWQ1
B	29	R1A	ALA	engineered mutation	UNP A0A0H3MWQ1
B	51	R1A	ASP	engineered mutation	UNP A0A0H3MWQ1
B	70	LEU	ILE	engineered mutation	UNP A0A0H3MWQ1
B	79	R1A	VAL	engineered mutation	UNP A0A0H3MWQ1
B	107	ASP	VAL	engineered mutation	UNP A0A0H3MWQ1
B	113	ALA	CYS	engineered mutation	UNP A0A0H3MWQ1
B	116	R1A	LEU	engineered mutation	UNP A0A0H3MWQ1
C	-9	MET	-	initiating methionine	UNP A0A0H3MWQ1
C	-8	GLY	-	expression tag	UNP A0A0H3MWQ1
C	-7	HIS	-	expression tag	UNP A0A0H3MWQ1
C	-6	HIS	-	expression tag	UNP A0A0H3MWQ1
C	-5	HIS	-	expression tag	UNP A0A0H3MWQ1
C	-4	HIS	-	expression tag	UNP A0A0H3MWQ1
C	-3	HIS	-	expression tag	UNP A0A0H3MWQ1
C	-2	HIS	-	expression tag	UNP A0A0H3MWQ1
C	-1	GLU	-	expression tag	UNP A0A0H3MWQ1
C	0	LEU	-	expression tag	UNP A0A0H3MWQ1
C	14	R1A	ALA	engineered mutation	UNP A0A0H3MWQ1
C	16	R1A	TYR	engineered mutation	UNP A0A0H3MWQ1
C	24	R1A	ALA	engineered mutation	UNP A0A0H3MWQ1
C	29	R1A	ALA	engineered mutation	UNP A0A0H3MWQ1
C	51	R1A	ASP	engineered mutation	UNP A0A0H3MWQ1
C	70	LEU	ILE	engineered mutation	UNP A0A0H3MWQ1
C	79	R1A	VAL	engineered mutation	UNP A0A0H3MWQ1
C	107	ASP	VAL	engineered mutation	UNP A0A0H3MWQ1
C	113	ALA	CYS	engineered mutation	UNP A0A0H3MWQ1
C	116	R1A	LEU	engineered mutation	UNP A0A0H3MWQ1

4 Residue-property plots


4.1 Average score per residue in the NMR ensemble

These plots are provided for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic is the same as shown in the summary in section 1 of this report. The second graphic shows the sequence where residues are colour-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. Stretches of 2 or more consecutive residues without any outliers are shown as green connectors. Residues which are classified as ill-defined in the NMR ensemble, are shown in cyan with an underline colour-coded according to the previous scheme. Residues which were present in the experimental sample, but not modelled in the final structure are shown in grey.


- Molecule 1: Diacylglycerol kinase



- Molecule 1: Diacylglycerol kinase



- Molecule 1: Diacylglycerol kinase




4.2 Scores per residue for each member of the ensemble

Colouring as in section 4.1 above.


4.2.1 Score per residue for model 1 (medoid)

- Molecule 1: Diacylglycerol kinase



- Molecule 1: Diacylglycerol kinase

Chain B:  74% 17% 8%



- Molecule 1: Diacylglycerol kinase


Chain C:  74% 17% 8%



4.2.2 Score per residue for model 2

- Molecule 1: Diacylglycerol kinase

Chain A:  75% 17% 8%




- Molecule 1: Diacylglycerol kinase

Chain B:  75% 17% 8%



- Molecule 1: Diacylglycerol kinase


Chain C:  75% 17% 8%




4.2.3 Score per residue for model 3

- Molecule 1: Diacylglycerol kinase


Chain A:  74% 17% 8%



- Molecule 1: Diacylglycerol kinase




- Molecule 1: Diacylglycerol kinase




4.2.4 Score per residue for model 4


- Molecule 1: Diacylglycerol kinase



- Molecule 1: Diacylglycerol kinase




- Molecule 1: Diacylglycerol kinase




4.2.5 Score per residue for model 5

- Molecule 1: Diacylglycerol kinase




- Molecule 1: Diacylglycerol kinase






- Molecule 1: Diacylglycerol kinase




4.2.6 Score per residue for model 6


- Molecule 1: Diacylglycerol kinase



- Molecule 1: Diacylglycerol kinase




- Molecule 1: Diacylglycerol kinase




4.2.7 Score per residue for model 7


- Molecule 1: Diacylglycerol kinase



- Molecule 1: Diacylglycerol kinase




- Molecule 1: Diacylglycerol kinase




4.2.8 Score per residue for model 8


- Molecule 1: Diacylglycerol kinase



- Molecule 1: Diacylglycerol kinase




- Molecule 1: Diacylglycerol kinase




4.2.9 Score per residue for model 9


- Molecule 1: Diacylglycerol kinase



- Molecule 1: Diacylglycerol kinase




- Molecule 1: Diacylglycerol kinase




4.2.10 Score per residue for model 10


- Molecule 1: Diacylglycerol kinase



- Molecule 1: Diacylglycerol kinase



- Molecule 1: Diacylglycerol kinase



5 Refinement protocol and experimental data overview

The models were refined using the following method: *distance geometry*.

Of the 20000 calculated structures, 10 were deposited, based on the following criterion: *structures with the lowest energy*.

The following table shows the software used for structure solution, optimisation and refinement.

Software name	Classification	Version
CS-ROSETTA	structure calculation	

The following table shows chemical shift validation statistics as aggregates over all chemical shift files. Detailed validation can be found in section 7 of this report.

Chemical shift file(s)	working_cs.cif
Number of chemical shift lists	1
Total number of shifts	457
Number of shifts mapped to atoms	457
Number of unparsed shifts	0
Number of shifts with mapping errors	0
Number of shifts with mapping warnings	0
Assignment completeness (well-defined parts)	11%

Note: This is a solid-state NMR structure, where hydrogen atoms are typically not assigned a chemical shift value, which may lead to lower completeness of assignment measure.

6 Model quality [i](#)

6.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: R1A

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the (average) root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	#Z>5	RMSZ	#Z>5
1	A	0.58±0.00	0±0/772 (0.0± 0.0%)	1.04±0.01	1±1/1042 (0.1± 0.1%)
1	B	0.58±0.00	0±0/772 (0.0± 0.0%)	1.04±0.01	1±1/1042 (0.1± 0.1%)
1	C	0.58±0.00	0±0/772 (0.0± 0.0%)	1.04±0.01	1±1/1042 (0.1± 0.1%)
All	All	0.58	0/23160 (0.0%)	1.04	30/31260 (0.1%)

There are no bond-length outliers.

All unique angle outliers are listed below. They are sorted according to the Z-score of the worst occurrence in the ensemble.

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)	Models	
								Worst	Total
1	A	82	ILE	N-CA-C	6.72	116.86	110.74	6	6
1	B	82	ILE	N-CA-C	6.72	116.84	110.53	8	6
1	C	82	ILE	N-CA-C	6.72	116.84	110.53	8	6
1	B	31	PHE	N-CA-C	-5.36	107.39	114.04	3	1
1	A	31	PHE	N-CA-C	-5.36	107.39	114.04	3	1
1	C	31	PHE	N-CA-C	-5.33	107.42	114.04	3	1
1	A	54	THR	N-CA-C	-5.19	106.10	112.90	6	3
1	C	54	THR	N-CA-C	-5.19	106.10	112.90	3	3
1	B	54	THR	N-CA-C	-5.18	106.11	112.90	3	3

There are no chirality outliers.

There are no planarity outliers.

6.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in each chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes

averaged over the ensemble.

Mol	Chain	Non-H	H(model)	H(added)	Clashes
1	A	764	793	795	1±1
1	B	764	793	795	1±1
1	C	764	793	795	1±1
All	All	22920	23790	23850	21

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 0.

All unique clashes are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Clash(Å)	Distance(Å)	Models	
				Worst	Total
1:B:28:GLU:HG2	1:B:28:GLU:O	0.59	1.98	7	1
1:C:28:GLU:O	1:C:28:GLU:HG2	0.59	1.98	7	1
1:A:28:GLU:HG2	1:A:28:GLU:O	0.58	1.98	7	1
1:A:28:GLU:O	1:A:28:GLU:CG	0.48	2.61	7	1
1:B:28:GLU:O	1:B:28:GLU:CG	0.48	2.61	7	1
1:C:28:GLU:O	1:C:28:GLU:CG	0.48	2.61	7	1
1:C:92:ARG:HA	1:C:92:ARG:NE	0.45	2.27	10	1
1:B:92:ARG:HA	1:B:92:ARG:NE	0.44	2.27	10	1
1:A:92:ARG:NE	1:A:92:ARG:HA	0.44	2.27	10	1
1:C:115:LEU:C	1:C:115:LEU:HD12	0.43	2.38	5	1
1:B:115:LEU:C	1:B:115:LEU:HD12	0.43	2.38	5	1
1:B:32:ARG:O	1:B:32:ARG:HD3	0.42	2.14	9	1
1:C:32:ARG:O	1:C:32:ARG:HD3	0.42	2.14	9	1
1:A:115:LEU:HD12	1:A:115:LEU:C	0.42	2.39	5	1
1:A:39:LEU:C	1:A:39:LEU:HD23	0.42	2.39	1	1
1:C:39:LEU:HD23	1:C:39:LEU:C	0.41	2.40	1	1
1:B:39:LEU:HD23	1:B:39:LEU:C	0.41	2.39	1	1
1:A:32:ARG:O	1:A:32:ARG:HD3	0.41	2.14	9	1
1:B:32:ARG:NE	1:B:32:ARG:HA	0.40	2.31	9	1
1:A:32:ARG:NE	1:A:32:ARG:HA	0.40	2.31	9	1
1:C:32:ARG:NE	1:C:32:ARG:HA	0.40	2.31	9	1

6.3 Torsion angles [i](#)

6.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all NMR entries. The Analysed column shows the number of residues for which the backbone conformation was analysed and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	98/131 (75%)	95±2 (97±2%)	3±2 (3±2%)	0±0 (0±0%)	100	100
1	B	98/131 (75%)	95±2 (97±2%)	3±2 (3±2%)	0±0 (0±0%)	100	100
1	C	98/131 (75%)	95±2 (97±2%)	3±2 (3±2%)	0±0 (0±0%)	100	100
All	All	2940/3930 (75%)	2850 (97%)	90 (3%)	0 (0%)	100	100

There are no Ramachandran outliers.

6.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all NMR entries. The Analysed column shows the number of residues for which the sidechain conformation was analysed and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	79/98 (81%)	79±0 (100±0%)	0±0 (0±0%)	100	100
1	B	79/98 (81%)	79±0 (100±0%)	0±0 (0±0%)	100	100
1	C	79/98 (81%)	79±0 (100±0%)	0±0 (0±0%)	100	100
All	All	2370/2940 (81%)	2370 (100%)	0 (0%)	100	100

There are no protein residues with a non-rotameric sidechain to report.

6.3.3 RNA ⓘ

There are no RNA molecules in this entry.

6.4 Non-standard residues in protein, DNA, RNA chains ⓘ

24 non-standard protein/DNA/RNA residues are modelled in this entry.

In the following table, the Counts columns list the number of bonds for which Mogul statistics could be retrieved, the number of bonds that are observed in the model and the number of bonds that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length is the number of standard deviations the observed value is removed from the expected value. A bond length with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the average root-mean-square of all Z scores of the bond lengths.

Mol	Type	Chain	Res	Link	Bond lengths		
					Counts	RMSZ	#Z>2
1	R1A	C	29	1	15,18,19	0.81±0.00	0±0 (0±0%)
1	R1A	A	24	1	15,18,19	0.83±0.00	0±0 (0±0%)
1	R1A	C	116	1	15,18,19	0.85±0.01	0±0 (0±0%)
1	R1A	A	46	1	15,18,19	0.87±0.11	0±1 (2±5%)
1	R1A	B	24	1	15,18,19	0.82±0.00	0±0 (0±0%)
1	R1A	B	46	1	15,18,19	0.87±0.10	0±1 (2±5%)
1	R1A	B	116	1	15,18,19	0.85±0.01	0±0 (0±0%)
1	R1A	A	51	1	15,18,19	0.84±0.01	0±0 (0±0%)
1	R1A	A	14	1	15,18,19	0.88±0.10	0±1 (2±5%)
1	R1A	B	16	1	15,18,19	0.83±0.01	0±0 (0±0%)
1	R1A	A	29	1	15,18,19	0.82±0.01	0±0 (0±0%)
1	R1A	C	79	1	15,18,19	0.83±0.01	0±0 (0±0%)
1	R1A	A	79	1	15,18,19	0.83±0.00	0±0 (0±0%)
1	R1A	C	46	1	15,18,19	0.87±0.11	0±1 (2±5%)
1	R1A	B	51	1	15,18,19	0.84±0.01	0±0 (0±0%)
1	R1A	C	14	1	15,18,19	0.88±0.10	0±1 (2±5%)
1	R1A	B	14	1	15,18,19	0.88±0.11	0±1 (2±5%)
1	R1A	A	116	1	15,18,19	0.85±0.01	0±0 (0±0%)
1	R1A	A	16	1	15,18,19	0.83±0.01	0±0 (0±0%)
1	R1A	B	79	1	15,18,19	0.83±0.01	0±0 (0±0%)
1	R1A	C	51	1	15,18,19	0.84±0.01	0±0 (0±0%)
1	R1A	B	29	1	15,18,19	0.82±0.01	0±0 (0±0%)
1	R1A	C	24	1	15,18,19	0.83±0.01	0±0 (0±0%)
1	R1A	C	16	1	15,18,19	0.83±0.00	0±0 (0±0%)

In the following table, the Counts columns list the number of angles for which Mogul statistics could be retrieved, the number of angles that are observed in the model and the number of angles that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond angle is the number of standard deviations the observed value is removed from the expected value. A bond angle with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the average root-mean-square of all Z scores of the bond angles.

Mol	Type	Chain	Res	Link	Bond angles		
					Counts	RMSZ	#Z>2
1	R1A	C	29	1	15,27,29	1.03±0.01	1±0 (8±2%)
1	R1A	A	24	1	15,27,29	1.03±0.01	1±0 (7±2%)
1	R1A	C	116	1	15,27,29	1.02±0.01	1±0 (6±0%)
1	R1A	A	46	1	15,27,29	1.01±0.04	1±0 (6±2%)
1	R1A	B	24	1	15,27,29	1.02±0.01	1±0 (8±3%)
1	R1A	B	46	1	15,27,29	1.01±0.04	1±0 (7±2%)

Mol	Type	Chain	Res	Link	Counts	Bond angles	
						RMSZ	#Z>2
1	R1A	B	116	1	15,27,29	1.02±0.01	1±0 (6±0%)
1	R1A	A	51	1	15,27,29	1.03±0.01	1±0 (8±3%)
1	R1A	A	14	1	15,27,29	1.01±0.03	2±1 (10±4%)
1	R1A	B	16	1	15,27,29	1.02±0.01	1±0 (8±3%)
1	R1A	A	29	1	15,27,29	1.03±0.01	1±0 (7±2%)
1	R1A	C	79	1	15,27,29	1.03±0.01	1±0 (6±0%)
1	R1A	A	79	1	15,27,29	1.03±0.01	1±0 (8±2%)
1	R1A	C	46	1	15,27,29	1.01±0.04	1±0 (8±3%)
1	R1A	B	51	1	15,27,29	1.03±0.01	1±0 (8±2%)
1	R1A	C	14	1	15,27,29	1.01±0.02	1±1 (9±5%)
1	R1A	B	14	1	15,27,29	1.02±0.03	2±1 (11±4%)
1	R1A	A	116	1	15,27,29	1.02±0.01	1±0 (8±2%)
1	R1A	A	16	1	15,27,29	1.03±0.01	1±0 (7±2%)
1	R1A	B	79	1	15,27,29	1.03±0.01	1±0 (6±0%)
1	R1A	C	51	1	15,27,29	1.03±0.01	1±0 (6±0%)
1	R1A	B	29	1	15,27,29	1.03±0.00	1±0 (7±2%)
1	R1A	C	24	1	15,27,29	1.03±0.01	1±0 (7±2%)
1	R1A	C	16	1	15,27,29	1.03±0.01	1±0 (8±2%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
1	R1A	B	24	1	-	0±0,5,32,34	0±0,1,1,1
1	R1A	C	116	1	-	0±0,5,32,34	0±0,1,1,1
1	R1A	B	14	1	-	0±0,5,32,34	0±0,1,1,1
1	R1A	A	24	1	-	0±0,5,32,34	0±0,1,1,1
1	R1A	A	16	1	-	0±0,5,32,34	0±0,1,1,1
1	R1A	A	79	1	-	0±0,5,32,34	0±0,1,1,1
1	R1A	A	14	1	-	0±0,5,32,34	0±0,1,1,1
1	R1A	C	14	1	-	0±0,5,32,34	0±0,1,1,1
1	R1A	C	29	1	-	0±0,5,32,34	0±0,1,1,1
1	R1A	C	16	1	-	0±0,5,32,34	0±0,1,1,1
1	R1A	C	24	1	-	0±0,5,32,34	0±0,1,1,1
1	R1A	B	79	1	-	0±0,5,32,34	0±0,1,1,1
1	R1A	C	51	1	-	0±0,5,32,34	0±0,1,1,1

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
1	R1A	A	29	1	-	0±0,5,32,34	0±0,1,1,1
1	R1A	B	51	1	-	0±0,5,32,34	0±0,1,1,1
1	R1A	A	46	1	-	0±0,5,32,34	0±0,1,1,1
1	R1A	B	16	1	-	0±0,5,32,34	0±0,1,1,1
1	R1A	C	46	1	-	0±0,5,32,34	0±0,1,1,1
1	R1A	B	116	1	-	0±0,5,32,34	0±0,1,1,1
1	R1A	B	46	1	-	0±0,5,32,34	0±0,1,1,1
1	R1A	C	79	1	-	0±0,5,32,34	0±0,1,1,1
1	R1A	A	51	1	-	0±0,5,32,34	0±0,1,1,1
1	R1A	A	116	1	-	0±0,5,32,34	0±0,1,1,1
1	R1A	B	29	1	-	0±0,5,32,34	0±0,1,1,1

All unique bond outliers are listed below. They are sorted according to the Z-score of the worst occurrence in the ensemble.

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)	Models	
								Worst	Total
1	C	46	R1A	O1-N1	2.71	1.14	1.26	10	2
1	A	46	R1A	O1-N1	2.71	1.14	1.26	10	2
1	B	14	R1A	O1-N1	2.71	1.14	1.26	9	2
1	B	46	R1A	O1-N1	2.71	1.14	1.26	2	2
1	A	14	R1A	O1-N1	2.71	1.14	1.26	2	2
1	C	14	R1A	O1-N1	2.70	1.14	1.26	2	2
1	B	14	R1A	C4-C3	2.14	1.34	1.32	2	2
1	C	14	R1A	C4-C3	2.12	1.34	1.32	9	2
1	C	46	R1A	C4-C3	2.11	1.34	1.32	2	2
1	A	14	R1A	C4-C3	2.11	1.34	1.32	9	2
1	B	46	R1A	C4-C3	2.11	1.34	1.32	10	2
1	A	46	R1A	C4-C3	2.10	1.34	1.32	2	2

All unique angle outliers are listed below. They are sorted according to the Z-score of the worst occurrence in the ensemble.

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)	Models	
								Worst	Total
1	A	79	R1A	C5-C4-C3	2.83	110.93	113.48	4	10
1	B	79	R1A	C5-C4-C3	2.82	110.93	113.48	2	10
1	B	51	R1A	C5-C4-C3	2.82	110.93	113.48	5	10
1	B	46	R1A	C5-C4-C3	2.82	110.94	113.48	9	9
1	C	24	R1A	C5-C4-C3	2.82	110.94	113.48	5	10
1	A	16	R1A	C5-C4-C3	2.81	110.94	113.48	1	10
1	C	29	R1A	C5-C4-C3	2.81	110.94	113.48	3	10
1	C	14	R1A	C5-C4-C3	2.81	110.94	113.48	4	10

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)	Models	
								Worst	Total
1	C	51	R1A	C5-C4-C3	2.80	110.95	113.48	7	10
1	C	79	R1A	C5-C4-C3	2.80	110.95	113.48	8	10
1	B	14	R1A	C5-C4-C3	2.80	110.95	113.48	7	10
1	A	24	R1A	C5-C4-C3	2.80	110.95	113.48	6	10
1	B	116	R1A	C5-C4-C3	2.79	110.96	113.48	9	10
1	C	116	R1A	C5-C4-C3	2.79	110.96	113.48	4	10
1	A	14	R1A	C5-C4-C3	2.79	110.96	113.48	5	10
1	A	51	R1A	C5-C4-C3	2.79	110.96	113.48	8	10
1	C	46	R1A	C5-C4-C3	2.79	110.96	113.48	9	10
1	C	16	R1A	C5-C4-C3	2.79	110.96	113.48	7	10
1	B	16	R1A	C5-C4-C3	2.79	110.96	113.48	9	10
1	A	29	R1A	C5-C4-C3	2.78	110.97	113.48	2	10
1	B	29	R1A	C5-C4-C3	2.78	110.97	113.48	7	10
1	A	116	R1A	C5-C4-C3	2.78	110.97	113.48	9	10
1	A	46	R1A	C5-C4-C3	2.78	110.97	113.48	9	9
1	B	24	R1A	C5-C4-C3	2.78	110.97	113.48	10	10
1	B	14	R1A	C6-C5-C4	2.09	110.75	112.76	2	3
1	C	46	R1A	C7-C5-C4	2.07	110.76	112.76	10	3
1	C	14	R1A	C7-C5-C4	2.07	110.76	112.76	2	2
1	C	14	R1A	C6-C5-C4	2.05	110.78	112.76	9	2
1	A	14	R1A	C6-C5-C4	2.04	110.79	112.76	2	2
1	A	79	R1A	C7-C5-C4	2.04	110.79	112.76	4	2
1	A	116	R1A	C7-C5-C4	2.04	110.79	112.76	9	2
1	B	46	R1A	C7-C5-C4	2.03	110.79	112.76	10	2
1	A	51	R1A	C7-C5-C4	2.03	110.80	112.76	7	3
1	A	14	R1A	C7-C5-C4	2.03	110.80	112.76	9	3
1	B	51	R1A	C7-C5-C4	2.02	110.81	112.76	3	2
1	A	46	R1A	C7-C5-C4	2.02	110.81	112.76	6	1
1	B	14	R1A	C7-C5-C4	2.02	110.81	112.76	2	4
1	B	16	R1A	C7-C5-C4	2.02	110.81	112.76	5	3
1	A	16	R1A	C7-C5-C4	2.02	110.81	112.76	4	1
1	A	24	R1A	C7-C5-C4	2.01	110.81	112.76	3	1
1	C	16	R1A	C7-C5-C4	2.01	110.81	112.76	7	2
1	C	29	R1A	C7-C5-C4	2.01	110.81	112.76	9	2
1	B	29	R1A	C7-C5-C4	2.01	110.82	112.76	10	1
1	B	24	R1A	C7-C5-C4	2.01	110.82	112.76	7	3
1	A	29	R1A	C7-C5-C4	2.01	110.82	112.76	8	1
1	C	24	R1A	C7-C5-C4	2.00	110.83	112.76	10	1

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

6.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.6 Ligand geometry [i](#)

There are no ligands in this entry.

6.7 Other polymers [i](#)

There are no such molecules in this entry.

6.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

7 Chemical shift validation

The completeness of assignment taking into account all chemical shift lists is 11% for the well-defined parts and 10% for the entire structure.

7.1 Chemical shift list 1

File name: working_cs.cif

Chemical shift list name: *no_cys_DgkA-assignment.txt*

7.1.1 Bookkeeping

The following table shows the results of parsing the chemical shift list and reports the number of nuclei with statistically unusual chemical shifts.

Total number of shifts	457
Number of shifts mapped to atoms	457
Number of unparsed shifts	0
Number of shifts with mapping errors	0
Number of shifts with mapping warnings	0
Number of shift outliers (ShiftChecker)	0

7.1.2 Chemical shift referencing

The following table shows the suggested chemical shift referencing corrections.

Nucleus	# values	Correction $\pm$ precision, ppm	Suggested action
$^{13}\text{C}_\alpha$	87	-0.91 ± 0.15	Should be checked
$^{13}\text{C}_\beta$	81	0.06 ± 0.15	None needed (< 0.5 ppm)
$^{13}\text{C}'$	87	-0.32 ± 0.07	None needed (< 0.5 ppm)
^{15}N	87	0.86 ± 0.35	Should be applied

7.1.3 Completeness of resonance assignments

The following table shows the completeness of the chemical shift assignments for the well-defined regions of the structure. The overall completeness is 11%, i.e. 452 atoms were assigned a chemical shift out of a possible 4152. 0 out of 72 assigned methyl groups (LEU and VAL) were assigned stereospecifically.

	Total	^1H	^{13}C	^{15}N
Backbone	258/1503 (17%)	0/612 (0%)	172/594 (29%)	86/297 (29%)
Sidechain	192/2346 (8%)	0/1572 (0%)	192/714 (27%)	0/60 (0%)

Continued on next page...

Continued from previous page...

	Total	¹H	¹³C	¹⁵N
Aromatic	2/303 (1%)	0/156 (0%)	2/132 (2%)	0/15 (0%)
Overall	452/4152 (11%)	0/2340 (0%)	366/1440 (25%)	86/372 (23%)

Note: This is a solid-state NMR structure, where hydrogen atoms are typically not assigned a chemical shift value, which may lead to lower completeness of assignment measure.

The following table shows the completeness of the chemical shift assignments for the full structure. The overall completeness is 10%, i.e. 455 atoms were assigned a chemical shift out of a possible 4698. 0 out of 72 assigned methyl groups (LEU and VAL) were assigned stereospecifically.

	Total	¹H	¹³C	¹⁵N
Backbone	261/1719 (15%)	0/702 (0%)	174/678 (26%)	87/339 (26%)
Sidechain	192/2646 (7%)	0/1767 (0%)	192/801 (24%)	0/78 (0%)
Aromatic	2/333 (1%)	0/171 (0%)	2/147 (1%)	0/15 (0%)
Overall	455/4698 (10%)	0/2640 (0%)	368/1626 (23%)	87/432 (20%)

Note: This is a solid-state NMR structure, where hydrogen atoms are typically not assigned a chemical shift value, which may lead to lower completeness of assignment measure.

7.1.4 Statistically unusual chemical shifts [i](#)

There are no statistically unusual chemical shifts.

7.1.5 Random Coil Index (RCI) plots [i](#)

The image below reports *random coil index* values for the protein chains in the structure. The height of each bar gives a probability of a given residue to be disordered, as predicted from the available chemical shifts and the amino acid sequence. A value above 0.2 is an indication of significant predicted disorder. The colour of the bar shows whether the residue is in the well-defined core (black) or in the ill-defined residue ranges (cyan), as described in section 2 on ensemble composition. If well-defined core and ill-defined regions are not identified then it is shown as gray bars.

Random coil index (RCI) for chain A:

