



wwPDB EM Validation Summary Report ⓘ

Mar 9, 2026 – 04:49 PM UTC

PDB ID : 6DW0 / pdb_00006dw0
EMDB ID : EMD-8922
Title : Cryo-EM structure of the benzodiazepine-sensitive $\alpha 1\beta\gamma 2S$ tri-heteromeric GABAA receptor in complex with GABA (Whole map)
Authors : Phulera, S.; Zhu, H.; Yu, J.; Yoshioka, C.; Gouaux, E.
Deposited on : 2018-06-26
Resolution : 3.80 Å(reported)

This is a wwPDB EM Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/EMValidationReportHelp>
with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

EMDB validation analysis : 0.0.1.dev132
Mogul : 2022.3.0, CSD as543be (2022)
MolProbity : 4-5-2 with Phenix2.0
Buster-report : wwPDB partial adaption of 1.1.7 (2018)
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
EM percentile statistics : 202505.v01 (Using data in the EMDB archive up until May 2025)
MapQ : 1.9.13
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49



1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

ELECTRON MICROSCOPY

The reported resolution of this entry is 3.80 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.








Metric	Whole archive (#Entries)	EM structures (#Entries)	Similar EM resolution (#Entries, resolution range(Å))
Clashscore	229148	23984	-
Ramachandran outliers	224038	23583	-
Sidechain outliers	223484	23102	-
Q-score	-	25397	10198 (3.30 - 4.30)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the map. The red, orange, yellow and green segments of the bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the EM map (all-atom inclusion $< 40\%$). The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	D	490	15% 54% 7% 39%
2	A	402	7% 62% 11% 26%
2	C	402	14% 62% 12% 26%
3	B	384	14% 72% 11% 17%

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
3	E	384	
4	F	4	
5	G	7	
6	H	3	
6	J	3	
7	I	2	
7	K	2	

2 Entry composition

There are 8 unique types of molecules in this entry. The entry contains 11929 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Gamma-aminobutyric acid receptor subunit gamma-2.

Mol	Chain	Residues	Atoms					AltConf	Trace
1	D	299	Total	C	N	O	S	0	0
			2221	1428	370	413	10		

There are 24 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
D	429	LEU	-	expression tag	UNP P18508
D	430	VAL	-	expression tag	UNP P18508
D	431	PRO	-	expression tag	UNP P18508
D	432	ARG	-	expression tag	UNP P18508
D	433	GLY	-	expression tag	UNP P18508
D	434	SER	-	expression tag	UNP P18508
D	435	ARG	-	expression tag	UNP P18508
D	436	HIS	-	expression tag	UNP P18508
D	437	HIS	-	expression tag	UNP P18508
D	438	HIS	-	expression tag	UNP P18508
D	439	HIS	-	expression tag	UNP P18508
D	440	HIS	-	expression tag	UNP P18508
D	441	HIS	-	expression tag	UNP P18508
D	442	HIS	-	expression tag	UNP P18508
D	443	HIS	-	expression tag	UNP P18508
D	444	THR	-	expression tag	UNP P18508
D	445	GLU	-	expression tag	UNP P18508
D	446	THR	-	expression tag	UNP P18508
D	447	SER	-	expression tag	UNP P18508
D	448	GLN	-	expression tag	UNP P18508
D	449	VAL	-	expression tag	UNP P18508
D	450	ALA	-	expression tag	UNP P18508
D	451	PRO	-	expression tag	UNP P18508
D	452	ALA	-	expression tag	UNP P18508

- Molecule 2 is a protein called Gamma-aminobutyric acid receptor subunit alpha-1, Gamma-aminobutyric acid receptor subunit alpha-1.

Mol	Chain	Residues	Atoms					AltConf	Trace
2	C	296	Total	C	N	O	S	0	0
			2260	1441	389	417	13		
2	A	296	Total	C	N	O	S	0	0
			2229	1423	380	413	13		

There are 30 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
C	314	THR	-	linker	UNP P62813
C	362	LEU	-	expression tag	UNP P62813
C	363	VAL	-	expression tag	UNP P62813
C	364	PRO	-	expression tag	UNP P62813
C	365	ARG	-	expression tag	UNP P62813
C	366	GLY	-	expression tag	UNP P62813
C	367	SER	-	expression tag	UNP P62813
C	368	HIS	-	expression tag	UNP P62813
C	369	HIS	-	expression tag	UNP P62813
C	370	HIS	-	expression tag	UNP P62813
C	371	HIS	-	expression tag	UNP P62813
C	372	HIS	-	expression tag	UNP P62813
C	373	HIS	-	expression tag	UNP P62813
C	374	HIS	-	expression tag	UNP P62813
C	375	HIS	-	expression tag	UNP P62813
A	314	THR	-	linker	UNP P62813
A	362	LEU	-	expression tag	UNP P62813
A	363	VAL	-	expression tag	UNP P62813
A	364	PRO	-	expression tag	UNP P62813
A	365	ARG	-	expression tag	UNP P62813
A	366	GLY	-	expression tag	UNP P62813
A	367	SER	-	expression tag	UNP P62813
A	368	HIS	-	expression tag	UNP P62813
A	369	HIS	-	expression tag	UNP P62813
A	370	HIS	-	expression tag	UNP P62813
A	371	HIS	-	expression tag	UNP P62813
A	372	HIS	-	expression tag	UNP P62813
A	373	HIS	-	expression tag	UNP P62813
A	374	HIS	-	expression tag	UNP P62813
A	375	HIS	-	expression tag	UNP P62813


- Molecule 3 is a protein called Gamma-aminobutyric acid receptor subunit beta-1, Gamma-aminobutyric acid receptor subunit beta-1.

Mol	Chain	Residues	Atoms					AltConf	Trace
3	E	319	Total	C	N	O	S	0	0
			2459	1586	400	458	15		
3	B	319	Total	C	N	O	S	0	0
			2472	1603	400	457	12		

There are 32 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
E	309	GLY	-	linker	UNP P15431
E	310	THR	-	linker	UNP P15431
E	346	LEU	-	expression tag	UNP P15431
E	347	VAL	-	expression tag	UNP P15431
E	348	PRO	-	expression tag	UNP P15431
E	349	ARG	-	expression tag	UNP P15431
E	350	GLY	-	expression tag	UNP P15431
E	351	SER	-	expression tag	UNP P15431
E	352	HIS	-	expression tag	UNP P15431
E	353	HIS	-	expression tag	UNP P15431
E	354	HIS	-	expression tag	UNP P15431
E	355	HIS	-	expression tag	UNP P15431
E	356	HIS	-	expression tag	UNP P15431
E	357	HIS	-	expression tag	UNP P15431
E	358	HIS	-	expression tag	UNP P15431
E	359	HIS	-	expression tag	UNP P15431
B	309	GLY	-	linker	UNP P15431
B	310	THR	-	linker	UNP P15431
B	346	LEU	-	expression tag	UNP P15431
B	347	VAL	-	expression tag	UNP P15431
B	348	PRO	-	expression tag	UNP P15431
B	349	ARG	-	expression tag	UNP P15431
B	350	GLY	-	expression tag	UNP P15431
B	351	SER	-	expression tag	UNP P15431
B	352	HIS	-	expression tag	UNP P15431
B	353	HIS	-	expression tag	UNP P15431
B	354	HIS	-	expression tag	UNP P15431
B	355	HIS	-	expression tag	UNP P15431
B	356	HIS	-	expression tag	UNP P15431
B	357	HIS	-	expression tag	UNP P15431
B	358	HIS	-	expression tag	UNP P15431
B	359	HIS	-	expression tag	UNP P15431

- Molecule 4 is an oligosaccharide called alpha-D-mannopyranose-(1-3)-beta-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose.




Mol	Chain	Residues	Atoms				AltConf	Trace
4	F	4	Total	C	N	O	0	0
			50	28	2	20		

- Molecule 5 is an oligosaccharide called alpha-D-mannopyranose-(1-2)-alpha-D-mannopyranose-(1-3)-[alpha-D-mannopyranose-(1-3)-alpha-D-mannopyranose-(1-6)]beta-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose.



Mol	Chain	Residues	Atoms				AltConf	Trace
5	G	7	Total	C	N	O	0	0
			83	46	2	35		

- Molecule 6 is an oligosaccharide called beta-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose.




Mol	Chain	Residues	Atoms				AltConf	Trace
6	H	3	Total	C	N	O	0	0
			39	22	2	15		
6	J	3	Total	C	N	O	0	0
			39	22	2	15		

- Molecule 7 is an oligosaccharide called 2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose.




Mol	Chain	Residues	Atoms				AltConf	Trace
7	I	2	Total	C	N	O	0	0
			28	16	2	10		
7	K	2	Total	C	N	O	0	0
			28	16	2	10		


- Molecule 8 is GAMMA-AMINO-BUTANOIC ACID (CCD ID: ABU) (formula: $C_4H_9NO_2$) (labeled as "Ligand of Interest" by depositor).





Mol	Chain	Residues	Atoms				AltConf
8	A	1	Total	C	N	O	0
			7	4	1	2	
8	B	1	Total	C	N	O	0
			7	4	1	2	
8	B	1	Total	C	N	O	0
			7	4	1	2	




- Molecule 2: Gamma-aminobutyric acid receptor subunit alpha-1, Gamma-aminobutyric acid receptor subunit alpha-1




- Molecule 3: Gamma-aminobutyric acid receptor subunit beta-1, Gamma-aminobutyric acid receptor subunit beta-1



- Molecule 3: Gamma-aminobutyric acid receptor subunit beta-1, Gamma-aminobutyric acid receptor subunit beta-1





- Molecule 4: alpha-D-mannopyranose-(1-3)-beta-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose



- Molecule 5: alpha-D-mannopyranose-(1-2)-alpha-D-mannopyranose-(1-3)-[alpha-D-mannopyranose-(1-3)-alpha-D-mannopyranose-(1-6)]beta-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose



- Molecule 6: beta-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose



- Molecule 6: beta-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose



- Molecule 7: 2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose

Chain I:  100%

NAG1
NAG2

- Molecule 7: 2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose

Chain K:  50%
100%

NAG1
NAG2

4 Experimental information

Property	Value	Source
EM reconstruction method	SINGLE PARTICLE	Depositor
Imposed symmetry	POINT, C1	Depositor
Number of particles used	49417	Depositor
Resolution determination method	FSC 0.143 CUT-OFF	Depositor
CTF correction method	PHASE FLIPPING AND AMPLITUDE CORRECTION	Depositor
Microscope	FEI TITAN KRIOS	Depositor
Voltage (kV)	300	Depositor
Electron dose ($e^-/\text{\AA}^2$)	37	Depositor
Minimum defocus (nm)	-1200	Depositor
Maximum defocus (nm)	-2500	Depositor
Magnification	120000	Depositor
Image detector	FEI FALCON III (4k x 4k)	Depositor
Maximum map value	137.256	Depositor
Minimum map value	-158.255	Depositor
Average map value	-0.073	Depositor
Map value standard deviation	2.932	Depositor
Recommended contour level	17	Depositor
Map size (Å)	311.52, 311.52, 311.52	wwPDB
Map dimensions	240, 240, 240	wwPDB
Map angles (°)	90.0, 90.0, 90.0	wwPDB
Pixel spacing (Å)	1.298, 1.298, 1.298	Depositor

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: NAG, BMA, ABU, MAN

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	D	0.21	0/2270	0.53	0/3110
2	A	0.23	0/2275	0.54	2/3107 (0.1%)
2	C	0.23	0/2310	0.52	0/3150
3	B	0.21	0/2526	0.52	0/3445
3	E	0.20	0/2510	0.50	0/3421
All	All	0.22	0/11891	0.52	2/16233 (0.0%)

There are no bond length outliers.

All (2) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	111	MET	CA-C-N	5.18	131.44	121.54
2	A	111	MET	C-N-CA	5.18	131.44	121.54

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	D	2221	0	2069	20	0
2	A	2229	0	2162	27	0
2	C	2260	0	2181	27	0
3	B	2472	0	2413	26	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
3	E	2459	0	2400	27	0
4	F	50	0	43	0	0
5	G	83	0	70	0	0
6	H	39	0	34	1	0
6	J	39	0	34	0	0
7	I	28	0	25	0	0
7	K	28	0	25	0	0
8	A	7	0	0	0	0
8	B	14	0	0	0	0
All	All	11929	0	11456	117	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 5.

The worst 5 of 117 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
3:E:64:GLN:HE21	3:E:125:LEU:HD11	1.57	0.67
3:B:43:ASP:HB2	3:B:62:TYR:HB2	1.80	0.64
1:D:73:THR:HA	1:D:147:ILE:O	1.97	0.63
1:D:123:TRP:HA	1:D:127:PRO:HB3	1.79	0.63
1:D:43:ARG:NH2	1:D:45:ASP:O	2.35	0.59

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all EM entries.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	D	297/490 (61%)	272 (92%)	25 (8%)	0	100	100
2	A	294/402 (73%)	276 (94%)	18 (6%)	0	100	100

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
2	C	294/402 (73%)	273 (93%)	21 (7%)	0	100	100
3	B	315/384 (82%)	303 (96%)	12 (4%)	0	100	100
3	E	315/384 (82%)	304 (96%)	11 (4%)	0	100	100
All	All	1515/2062 (74%)	1428 (94%)	87 (6%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all EM entries.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	D	225/449 (50%)	225 (100%)	0	100	100
2	A	233/360 (65%)	233 (100%)	0	100	100
2	C	237/360 (66%)	237 (100%)	0	100	100
3	B	263/348 (76%)	262 (100%)	1 (0%)	84	83
3	E	263/348 (76%)	263 (100%)	0	100	100
All	All	1221/1865 (66%)	1220 (100%)	1 (0%)	87	89

All (1) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
3	B	264	ILE

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 13 such sidechains are listed below:

Mol	Chain	Res	Type
2	A	215	HIS
2	A	217	HIS
3	B	64	GLN
3	E	64	GLN
3	E	336	ASN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

21 monosaccharides are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
4	NAG	F	1	2,4	14,14,15	0.23	0	17,19,21	0.50	0
4	NAG	F	2	4	14,14,15	0.21	0	17,19,21	0.59	0
4	BMA	F	3	4	11,11,12	0.72	0	15,15,17	0.96	1 (6%)
4	MAN	F	4	4	11,11,12	1.28	2 (18%)	15,15,17	1.16	2 (13%)
5	NAG	G	1	2,5	14,14,15	0.20	0	17,19,21	0.68	1 (5%)
5	NAG	G	2	5	14,14,15	0.24	0	17,19,21	0.57	0
5	BMA	G	3	5	11,11,12	0.70	0	15,15,17	1.22	3 (20%)
5	MAN	G	4	5	11,11,12	0.96	0	15,15,17	1.57	4 (26%)
5	MAN	G	5	5	11,11,12	0.94	1 (9%)	15,15,17	0.85	1 (6%)
5	MAN	G	6	5	11,11,12	0.96	0	15,15,17	1.34	2 (13%)
5	MAN	G	7	5	11,11,12	0.69	0	15,15,17	1.16	2 (13%)
6	NAG	H	1	6,3	14,14,15	0.47	0	17,19,21	0.39	0
6	NAG	H	2	6	14,14,15	0.49	0	17,19,21	0.41	0
6	BMA	H	3	6	11,11,12	0.77	0	15,15,17	0.84	0
7	NAG	I	1	7,3	14,14,15	0.23	0	17,19,21	0.45	0
7	NAG	I	2	7	14,14,15	0.39	0	17,19,21	0.40	0
6	NAG	J	1	6,3	14,14,15	0.36	0	17,19,21	0.54	0
6	NAG	J	2	6	14,14,15	0.20	0	17,19,21	0.56	0
6	BMA	J	3	6	11,11,12	0.80	0	15,15,17	0.76	0
7	NAG	K	1	7,3	14,14,15	0.23	0	17,19,21	0.50	0
7	NAG	K	2	7	14,14,15	0.38	0	17,19,21	0.43	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	NAG	F	1	2,4	-	3/6/23/26	0/1/1/1
4	NAG	F	2	4	-	3/6/23/26	0/1/1/1
4	BMA	F	3	4	-	2/2/19/22	0/1/1/1
4	MAN	F	4	4	-	1/2/19/22	0/1/1/1
5	NAG	G	1	2,5	-	2/6/23/26	0/1/1/1
5	NAG	G	2	5	-	1/6/23/26	0/1/1/1
5	BMA	G	3	5	-	2/2/19/22	0/1/1/1
5	MAN	G	4	5	-	2/2/19/22	0/1/1/1
5	MAN	G	5	5	-	1/2/19/22	0/1/1/1
5	MAN	G	6	5	-	1/2/19/22	0/1/1/1
5	MAN	G	7	5	-	2/2/19/22	0/1/1/1
6	NAG	H	1	6,3	-	2/6/23/26	0/1/1/1
6	NAG	H	2	6	-	2/6/23/26	0/1/1/1
6	BMA	H	3	6	-	0/2/19/22	0/1/1/1
7	NAG	I	1	7,3	-	2/6/23/26	0/1/1/1
7	NAG	I	2	7	-	2/6/23/26	0/1/1/1
6	NAG	J	1	6,3	-	2/6/23/26	0/1/1/1
6	NAG	J	2	6	-	0/6/23/26	0/1/1/1
6	BMA	J	3	6	-	0/2/19/22	0/1/1/1
7	NAG	K	1	7,3	-	1/6/23/26	0/1/1/1
7	NAG	K	2	7	-	1/6/23/26	0/1/1/1

All (3) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
4	F	4	MAN	C2-C3	2.66	1.56	1.52
4	F	4	MAN	C4-C3	2.22	1.58	1.52
5	G	5	MAN	O5-C1	-2.05	1.40	1.43

The worst 5 of 16 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
5	G	4	MAN	C1-O5-C5	3.92	117.44	112.19
5	G	6	MAN	C1-O5-C5	3.81	117.30	112.19
5	G	7	MAN	C1-O5-C5	3.13	116.39	112.19
5	G	4	MAN	O2-C2-C3	-2.99	103.97	110.15

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
5	G	6	MAN	O2-C2-C3	-2.56	104.85	110.15

There are no chirality outliers.

5 of 32 torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
6	J	1	NAG	O5-C5-C6-O6
6	H	2	NAG	O5-C5-C6-O6
7	I	2	NAG	O5-C5-C6-O6
5	G	3	BMA	O5-C5-C6-O6
6	H	2	NAG	C4-C5-C6-O6


There are no ring outliers.



1 monomer is involved in 1 short contact:


Mol	Chain	Res	Type	Clashes	Symm-Clashes
6	H	1	NAG	1	0


The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for oligosaccharide.











5.6 Ligand geometry [i](#)

3 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
8	ABU	B	407	-	6,6,6	0.87	0	6,6,6	1.50	2 (33%)
8	ABU	B	401	-	6,6,6	0.87	0	6,6,6	1.45	1 (16%)
8	ABU	A	408	-	6,6,6	0.89	0	6,6,6	1.37	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
8	ABU	B	407	-	-	1/4/4/4	-
8	ABU	B	401	-	-	0/4/4/4	-
8	ABU	A	408	-	-	0/4/4/4	-

There are no bond length outliers.

All (3) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
8	B	407	ABU	CB-CG-C	-2.19	108.78	114.51
8	B	407	ABU	OXT-C-CG	2.03	120.43	114.00
8	B	401	ABU	CB-CG-C	-2.01	109.26	114.51

There are no chirality outliers.



All (1) torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
8	B	407	ABU	CD-CB-CG-C

There are no ring outliers.

No monomer is involved in short contacts.

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.





5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.


6 Map visualisation [i](#)

This section contains visualisations of the EMDB entry EMD-8922. These allow visual inspection of the internal detail of the map and identification of artifacts.


No raw map or half-maps were deposited for this entry and therefore no images, graphs, etc. pertaining to the raw map can be shown.

6.1 Orthogonal projections [i](#)


6.1.1 Primary map



X



Y




Z


The images above show the map projected in three orthogonal directions.

6.2 Central slices [i](#)


6.2.1 Primary map



X Index: 120



Y Index: 120




Z Index: 120


The images above show central slices of the map in three orthogonal directions.

6.3 Largest variance slices [i](#)


6.3.1 Primary map



X Index: 106



Y Index: 111




Z Index: 135


The images above show the largest variance slices of the map in three orthogonal directions.

6.4 Orthogonal standard-deviation projections (False-color) [i](#)


6.4.1 Primary map



X



Y



Z

The images above show the map standard deviation projections with false color in three orthogonal directions. Minimum values are shown in green, max in blue, and dark to light orange shades represent small to large values respectively.

6.5 Orthogonal surface views [i](#)

6.5.1 Primary map



X



Y



Z

The images above show the 3D surface view of the map at the recommended contour level 17.0. These images, in conjunction with the slice images, may facilitate assessment of whether an appropriate contour level has been provided.


6.6 Mask visualisation [i](#)

This section was not generated. No masks/segmentation were deposited.

7 Map analysis ⓘ


This section contains the results of statistical analysis of the map.

7.1 Map-value distribution ⓘ



The map-value distribution is plotted in 128 intervals along the x-axis. The y-axis is logarithmic. A spike in this graph at zero usually indicates that the volume has been masked.


7.2 Volume estimate [i](#)



The volume at the recommended contour level is 119 nm³; this corresponds to an approximate mass of 108 kDa.

The volume estimate graph shows how the enclosed volume varies with the contour level. The recommended contour level is shown as a vertical line and the intersection between the line and the curve gives the volume of the enclosed surface at the given level.

7.3 Rotationally averaged power spectrum ⓘ



*Reported resolution corresponds to spatial frequency of 0.263 Å⁻¹


8 Fourier-Shell correlation

This section was not generated. No FSC curve or half-maps provided.

9 Map-model fit [i](#)


This section contains information regarding the fit between EMDB map EMD-8922 and PDB model 6DW0. Per-residue inclusion information can be found in [section 3](#) on [page 9](#).

9.1 Map-model overlay [i](#)



The images above show the 3D surface view of the map at the recommended contour level 17.0 at 50% transparency in yellow overlaid with a ribbon representation of the model coloured in blue. These images allow for the visual assessment of the quality of fit between the atomic model and the map.

9.2 Q-score mapped to coordinate model [i](#)




The images above show the model with each residue coloured according to its Q-score. This shows their resolvability in the map with higher Q-score values reflecting better resolvability. Please note: Q-score is calculating the resolvability of atoms, and thus high values are only expected at resolutions at which atoms can be resolved. Low Q-score values may therefore be expected for many entries.

9.3 Atom inclusion mapped to coordinate model [i](#)

This section was not generated.

9.4 Atom inclusion [i](#)



At the recommended contour level, 84% of all backbone atoms, 59% of all non-hydrogen atoms, are inside the map.

9.5 Map-model fit summary ⓘ

The table lists the average atom inclusion at the recommended contour level (17) and Q-score for the entire model and for each chain.

Chain	Atom inclusion	Q-score
All	0.5920	0.3550
A	0.6260	0.3680
B	0.5900	0.3410
C	0.6070	0.3710
D	0.5630	0.3470
E	0.5790	0.3430
F	0.5000	0.4680
G	0.5900	0.4730
H	0.4620	0.4130
I	0.6790	0.4110
J	0.4870	0.4010
K	0.5000	0.3750

1.0

0.0

<0.0