



wwPDB EM Validation Summary Report ⓘ

Mar 6, 2026 – 03:53 PM UTC

PDB ID : 8DWO / pdb_00008dwo
EMDB ID : EMD-27757
Title : Cryo-EM Structure of Eastern Equine Encephalitis Virus in complex with SKE26 Fab
Authors : Pletnev, S.; Verardi, R.; Roedeger, M.; Kwong, P.
Deposited on : 2022-08-01
Resolution : 3.50 Å(reported)
Based on initial model : 6XO4

This is a wwPDB EM Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/EMValidationReportHelp>
with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

EMDB validation analysis : 0.0.1.dev132
MolProbity : 4-5-2 with Phenix2.0
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
EM percentile statistics : 202505.v01 (Using data in the EMDB archive up until May 2025)
MapQ : 1.9.13
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49



1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

ELECTRON MICROSCOPY

The reported resolution of this entry is 3.50 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.









Metric	Whole archive (#Entries)	EM structures (#Entries)	Similar EM resolution (#Entries, resolution range(Å))
Clashscore	229148	23984	-
Ramachandran outliers	224038	23583	-
Sidechain outliers	223484	23102	-
Q-score	-	25397	13950 (3.00 - 4.00)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the map. The red, orange, yellow and green segments of the bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the EM map (all-atom inclusion $< 40\%$). The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	441	
1	F	441	
1	J	441	
2	B	420	

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
2	G	420	
2	K	420	
3	H	233	
3	S	233	
3	U	233	
4	L	219	
4	T	219	
4	V	219	

2 Entry composition [i](#)

There are 4 unique types of molecules in this entry. The entry contains 22572 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Envelope glycoprotein E1.

Mol	Chain	Residues	Atoms					AltConf	Trace
1	A	392	Total 3000	C 1903	N 498	O 579	S 20	0	0
1	F	392	Total 3000	C 1903	N 498	O 579	S 20	0	0
1	J	392	Total 3000	C 1903	N 498	O 579	S 20	0	0

There are 3 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	89	TYR	TRP	conflict	UNP Q88678
F	89	TYR	TRP	conflict	UNP Q88678
J	89	TYR	TRP	conflict	UNP Q88678

- Molecule 2 is a protein called Envelope glycoprotein E2.

Mol	Chain	Residues	Atoms					AltConf	Trace
2	B	346	Total 2720	C 1706	N 503	O 495	S 16	0	0
2	G	346	Total 2720	C 1706	N 503	O 495	S 16	0	0
2	K	346	Total 2720	C 1706	N 503	O 495	S 16	0	0

- Molecule 3 is a protein called SKE26 Fab Heavy Chain.

Mol	Chain	Residues	Atoms					AltConf	Trace
3	H	122	Total 972	C 624	N 164	O 181	S 3	0	0
3	S	122	Total 972	C 624	N 164	O 181	S 3	0	0
3	U	122	Total 972	C 624	N 164	O 181	S 3	0	0


- Molecule 4 is a protein called SKE26 Fab Light Chain.

Mol	Chain	Residues	Atoms					AltConf	Trace
4	L	112	Total	C	N	O	S	0	0
			832	521	139	168	4		
4	T	112	Total	C	N	O	S	0	0
			832	521	139	168	4		
4	V	112	Total	C	N	O	S	0	0
			832	521	139	168	4		


3 Residue-property plots

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and atom inclusion in map density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red diamond above a residue indicates a poor fit to the EM map for this residue (all-atom inclusion < 40%). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


• Molecule 1: Envelope glycoprotein E1




• Molecule 1: Envelope glycoprotein E1




• Molecule 1: Envelope glycoprotein E1







• Molecule 2: Envelope glycoprotein E2



• Molecule 2: Envelope glycoprotein E2



• Molecule 2: Envelope glycoprotein E2



GLN
VAL
PRO
THR
ILE
LEU
LEU
ALA
LEU
CYS
CYS
ILE
LYS
PRO
THR
ARG
ALA

• Molecule 3: SKE26 Fab Heavy Chain

Chain H:  43% 9% 48%



SER ARG SER THR GLU SER THR ALA LEU GLY CYS LEU VAL LYS ASP TYR PHE GLU PRO VAL THR VAL SER TRP VAL SER THR ASN CYS GLY SER LEU THR GLY VAL HIS THR PHE PRO ALA VAL G100C ALA LEU GLN SER S113 ALA SER THR TYR SER LYS LEU SER GLY PRO SER SER VAL VAL PHE THR VAL PRO LEU ALA PRO SER SER

SER LEU GLY THR GLN THR TYR VAL CYS ASN VAL ASN HIS LYS PRO SER LYS ASP THR LYS VAL ASP PRO VAL ARG VAL GLU THR LYS THR CYS GLY SER LEU VAL LEU PHE GLN

• Molecule 3: SKE26 Fab Heavy Chain

Chain S:  43% 9% 48%



ARG SER THR GLU THR SER VAL CYS ASN VAL ASN HIS LYS PRO SER LYS ASP THR PHE PRO GLU PRO VAL THR VAL SER TRP ASN THR CYS GLY SER LEU THR GLY VAL HIS THR PHE PRO ALA VAL G100C ALA LEU GLN SER SER GLY ALA SER THR TYR SER LYS LEU SER SER VAL VAL PHE THR VAL PRO ALA PRO SER SER

LEU GLY THR GLN THR TYR VAL CYS ASN VAL ASN HIS LYS PRO SER LYS ASP THR PHE PRO VAL ASP PRO VAL ARG VAL GLU THR LYS THR CYS GLY SER LEU VAL LEU PHE GLN

• Molecule 3: SKE26 Fab Heavy Chain


Chain U:  42% 11% 48%



LEU ALA PRO SER SER ARG SER THR THR GLN THR TYR VAL CYS ASN VAL ASN HIS LYS PRO SER LYS ASP THR PHE PRO VAL ASP PRO VAL ARG VAL THR VAL SER TRP ASN THR CYS GLY SER LEU VAL LEU PHE GLN

VAL PRO SER SER SER GLY THR THR GLN THR TYR VAL CYS ASN VAL VAL ASN HIS LYS PRO SER SER LYS ASP THR PHE PRO VAL ARG VAL THR VAL SER TRP ASN THR CYS GLY SER LEU VAL LEU PHE GLN

• Molecule 4: SKE26 Fab Light Chain


Chain L:  41% 10% 49%



SER VAL VAL CYS LEU LEU ASN ASN PHE THR THR GLN THR TYR VAL CYS ASN VAL VAL ASN HIS LYS PRO SER SER LYS ASP THR PHE PRO VAL ARG VAL THR VAL SER TRP ASN THR CYS GLY SER LEU VAL LEU PHE GLN

VAL
TYR
ALA
CYS
GLU
VAL
THR
HIS
GLN
GLY
LEU
SER
SER
SER
PRO
VAL
LYS
THR
PHE
ASN
ARG
GLY
GLU
CYS


● Molecule 4: SKE26 Fab Light Chain



PHE
TYR
PRO
ARG
GLU
SER
ALA
VAL
SER
VAL
LYS
TRP
LYS
PHE
VAL
ASN
ASP
GLY
ALA
LEU
LYS
THR
GLY
ASN
SER
GLN
GLU
SER
VAL
THR
GLN
ASP
SER
LYS
ASN
VAL
THR
TYR
SER
LEU
SER
SER
THR
LEU
THR
LEU
SER
SER
THR
GLU
TYR
GLN
SER
HIS
LYS
VAL
TYR
VAL
ALA
CYS
GLU
VAL
THR
HIS

GLN
GLY
LEU
SER
SER
PRO
VAL
THR
LYS
SER
PHE
ASN
ARG
GLY
GLU
CYS

● Molecule 4: SKE26 Fab Light Chain



GLY
THR
VAL
SER
VAL
VAL
CYS
LEU
LEU
ASN
ASN
PHE
GLN
TYR
PRO
ARG
GLU
ALA
SER
VAL
LYS
TRP
SER
LYS
VAL
VAL
ASP
GLY
ALA
LEU
LYS
THR
GLY
ASN
SER
GLN
GLU
SER
VAL
THR
THR
GLU
GLN
ASP
SER
LYS
ASP
ASN
THR
TYR
SER
SER
SER
SER
THR
THR
THR
SER
SER
THR
GLU
TYR
GLN

SER
HIS
LYS
VAL
TYR
ALA
CYS
GLU
VAL
THR
HIS
GLN
GLY
LEU
SER
SER
PRO
VAL
THR
LYS
SER
PHE
ASN
ARG
GLY
GLU
CYS

4 Experimental information

Property	Value	Source
EM reconstruction method	SINGLE PARTICLE	Depositor
Imposed symmetry	POINT, Not provided	
Number of particles used	3071400	Depositor
Resolution determination method	FSC 0.143 CUT-OFF	Depositor
CTF correction method	PHASE FLIPPING AND AMPLITUDE CORRECTION	Depositor
Microscope	FEI TITAN KRIOS	Depositor
Voltage (kV)	300	Depositor
Electron dose ($e^-/\text{\AA}^2$)	58.06	Depositor
Minimum defocus (nm)	700	Depositor
Maximum defocus (nm)	2000	Depositor
Magnification	Not provided	
Image detector	GATAN K3 (6k x 4k)	Depositor
Maximum map value	5.102	Depositor
Minimum map value	-3.035	Depositor
Average map value	0.000	Depositor
Map value standard deviation	0.061	Depositor
Recommended contour level	0.4	Depositor
Map size (Å)	1162.0, 1162.0, 1162.0	wwPDB
Map dimensions	700, 700, 700	wwPDB
Map angles (°)	90.0, 90.0, 90.0	wwPDB
Pixel spacing (Å)	1.66, 1.66, 1.66	Depositor

5 Model quality [i](#)

5.1 Standard geometry [i](#)

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.24	0/3080	0.53	0/4197
1	F	0.24	0/3080	0.52	0/4197
1	J	0.25	0/3080	0.52	1/4197 (0.0%)
2	B	0.22	0/2799	0.48	0/3808
2	G	0.21	0/2799	0.49	0/3808
2	K	0.22	0/2799	0.50	0/3808
3	H	0.20	0/1001	0.48	0/1363
3	S	0.19	0/1001	0.43	0/1363
3	U	0.21	0/1001	0.45	0/1363
4	L	0.22	0/850	0.47	0/1159
4	T	0.22	0/850	0.49	0/1159
4	V	0.24	0/850	0.52	0/1159
All	All	0.23	0/23190	0.50	1/31581 (0.0%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	1

There are no bond length outliers.

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	J	154	PRO	CA-N-CD	-6.14	103.40	112.00

There are no chirality outliers.

All (1) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	62	CYS	Peptide

5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3000	0	2900	35	0
1	F	3000	0	2900	44	0
1	J	3000	0	2902	29	0
2	B	2720	0	2655	27	0
2	G	2720	0	2655	45	0
2	K	2720	0	2655	44	0
3	H	972	0	936	14	0
3	S	972	0	936	14	0
3	U	972	0	936	17	0
4	L	832	0	811	12	0
4	T	832	0	811	11	0
4	V	832	0	811	15	0
All	All	22572	0	21908	290	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 7.

The worst 5 of 290 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:J:135:ILE:HD12	1:J:157:ILE:HD13	1.59	0.84
3:U:51:ILE:HD13	3:U:57:ILE:HD13	1.65	0.78
3:S:56:ASP:OD2	3:S:58:ARG:NH2	2.20	0.75
1:J:92:ALA:O	2:K:224:ARG:NH1	2.18	0.74
1:J:75:ASP:OD2	1:J:218:THR:OG1	2.05	0.74

There are no symmetry-related clashes.

5.3 Torsion angles

5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all EM entries.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	388/441 (88%)	359 (92%)	29 (8%)	0	100	100
1	F	388/441 (88%)	359 (92%)	29 (8%)	0	100	100
1	J	388/441 (88%)	355 (92%)	33 (8%)	0	100	100
2	B	344/420 (82%)	308 (90%)	36 (10%)	0	100	100
2	G	344/420 (82%)	310 (90%)	34 (10%)	0	100	100
2	K	344/420 (82%)	306 (89%)	38 (11%)	0	100	100
3	H	120/233 (52%)	116 (97%)	4 (3%)	0	100	100
3	S	120/233 (52%)	117 (98%)	3 (2%)	0	100	100
3	U	120/233 (52%)	118 (98%)	2 (2%)	0	100	100
4	L	110/219 (50%)	107 (97%)	3 (3%)	0	100	100
4	T	110/219 (50%)	106 (96%)	4 (4%)	0	100	100
4	V	110/219 (50%)	107 (97%)	3 (3%)	0	100	100
All	All	2886/3939 (73%)	2668 (92%)	218 (8%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all EM entries.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	333/371 (90%)	332 (100%)	1 (0%)	86	83
1	F	333/371 (90%)	333 (100%)	0	100	100

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	J	333/371 (90%)	333 (100%)	0	100	100
2	B	300/367 (82%)	299 (100%)	1 (0%)	86	83
2	G	300/367 (82%)	297 (99%)	3 (1%)	68	75
2	K	300/367 (82%)	298 (99%)	2 (1%)	76	78
3	H	106/204 (52%)	106 (100%)	0	100	100
3	S	106/204 (52%)	106 (100%)	0	100	100
3	U	106/204 (52%)	106 (100%)	0	100	100
4	L	95/192 (50%)	89 (94%)	6 (6%)	16	42
4	T	95/192 (50%)	95 (100%)	0	100	100
4	V	95/192 (50%)	94 (99%)	1 (1%)	65	74
All	All	2502/3402 (74%)	2488 (99%)	14 (1%)	76	79

5 of 14 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
4	L	27	GLN
4	L	27(A)	SER
4	V	33	LEU
4	L	27(D)	ASP
4	L	27(E)	SER

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 17 such sidechains are listed below:

Mol	Chain	Res	Type
2	K	160	GLN
4	V	30	ASN
2	G	126	HIS
2	G	285	HIS
1	J	134	ASN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

There are no ligands in this entry.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.


6 Map visualisation [i](#)

This section contains visualisations of the EMDB entry EMD-27757. These allow visual inspection of the internal detail of the map and identification of artifacts.


Images derived from a raw map, generated by summing the deposited half-maps, are presented below the corresponding image components of the primary map to allow further visual inspection and comparison with those of the primary map.

6.1 Orthogonal projections [i](#)


6.1.1 Primary map



X




Y




Z


6.1.2 Raw map



X



Y




Z


The images above show the map projected in three orthogonal directions.

6.2 Central slices [i](#)


6.2.1 Primary map



X Index: 350




Y Index: 350




Z Index: 350


6.2.2 Raw map



X Index: 350



Y Index: 350




Z Index: 350


The images above show central slices of the map in three orthogonal directions.

6.3 Largest variance slices [i](#)


6.3.1 Primary map



X Index: 336




Y Index: 489




Z Index: 473


6.3.2 Raw map



X Index: 336



Y Index: 472




Z Index: 473


The images above show the largest variance slices of the map in three orthogonal directions.

6.4 Orthogonal standard-deviation projections (False-color) [i](#)


6.4.1 Primary map



X




Y




Z


6.4.2 Raw map



X



Y




Z

The images above show the map standard deviation projections with false color in three orthogonal directions. Minimum values are shown in green, max in blue, and dark to light orange shades represent small to large values respectively.

6.5 Orthogonal surface views [i](#)

6.5.1 Primary map



The images above show the 3D surface view of the map at the recommended contour level 0.4. These images, in conjunction with the slice images, may facilitate assessment of whether an appropriate contour level has been provided.

6.5.2 Raw map



These images show the 3D surface of the raw map. The raw map's contour level was selected so that its surface encloses the same volume as the primary map does at its recommended contour level.


6.6 Mask visualisation [i](#)

This section shows the 3D surface view of the primary map at 50% transparency overlaid with the specified mask at 0% transparency


A mask typically either:

- Encompasses the whole structure
- Separates out a domain, a functional unit, a monomer or an area of interest from a larger structure


6.6.1 emd_27757_msk_1.map [i](#)



X



Y




Z

7 Map analysis [i](#)


This section contains the results of statistical analysis of the map.

7.1 Map-value distribution [i](#)



The map-value distribution is plotted in 128 intervals along the x-axis. The y-axis is logarithmic. A spike in this graph at zero usually indicates that the volume has been masked.


7.2 Volume estimate [i](#)



The volume at the recommended contour level is 818 nm³; this corresponds to an approximate mass of 739 kDa.

The volume estimate graph shows how the enclosed volume varies with the contour level. The recommended contour level is shown as a vertical line and the intersection between the line and the curve gives the volume of the enclosed surface at the given level.

7.3 Rotationally averaged power spectrum ⓘ




*Reported resolution corresponds to spatial frequency of 0.286 $\text{\AA}^{-1}$

8 Fourier-Shell correlation [i](#)

Fourier-Shell Correlation (FSC) is the most commonly used method to estimate the resolution of single-particle and subtomogram-averaged maps. The shape of the curve depends on the imposed symmetry, mask and whether or not the two 3D reconstructions used were processed from a common reference. The reported resolution is shown as a black line. A curve is displayed for the half-bit criterion in addition to lines showing the 0.143 gold standard cut-off and 0.5 cut-off.

8.1 FSC [i](#)



*Reported resolution corresponds to spatial frequency of 0.286 Å⁻¹

8.2 Resolution estimates [i](#)


Resolution estimate (Å)	Estimation criterion (FSC cut-off)		
	0.143	0.5	Half-bit
Reported by author	3.50	-	-
Author-provided FSC curve	3.36	3.65	3.37
Unmasked-calculated*	4.97	8.06	5.34

*Resolution estimate based on FSC curve calculated by comparison of deposited half-maps. The value from deposited half-maps intersecting FSC 0.143 CUT-OFF 4.97 differs from the reported value 3.5 by more than 10 %

9 Map-model fit [i](#)

This section contains information regarding the fit between EMDB map EMD-27757 and PDB model 8DWO. Per-residue inclusion information can be found in [section 3](#) on [page 6](#).

9.1 Map-model overlay [i](#)



The images above show the 3D surface view of the map at the recommended contour level 0.4 at 50% transparency in yellow overlaid with a ribbon representation of the model coloured in blue. These images allow for the visual assessment of the quality of fit between the atomic model and the map.

9.2 Q-score mapped to coordinate model [i](#)




The images above show the model with each residue coloured according to its Q-score. This shows their resolvability in the map with higher Q-score values reflecting better resolvability. Please note: Q-score is calculating the resolvability of atoms, and thus high values are only expected at resolutions at which atoms can be resolved. Low Q-score values may therefore be expected for many entries.

9.3 Atom inclusion mapped to coordinate model [i](#)



The images above show the model with each residue coloured according to its atom inclusion. This shows to what extent they are inside the map at the recommended contour level (0.4).

9.4 Atom inclusion [i](#)



At the recommended contour level, 93% of all backbone atoms, 88% of all non-hydrogen atoms, are inside the map.

9.5 Map-model fit summary ⓘ

The table lists the average atom inclusion at the recommended contour level (0.4) and Q-score for the entire model and for each chain.

Chain	Atom inclusion	Q-score
All	0.8850	0.4830
A	0.8920	0.4950
B	0.8990	0.4990
F	0.8930	0.4980
G	0.9000	0.4990
H	0.8620	0.4320
J	0.8630	0.4740
K	0.9000	0.4970
L	0.8500	0.4540
S	0.8910	0.4500
T	0.8400	0.4610
U	0.8810	0.4480
V	0.8540	0.4480

1.0

0.0

<0.0