



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 10, 2026 – 04:38 PM UTC

PDB ID : 2DX8 / pdb_00002dx8
Title : Crystal Structure Analysis of the PHD domain of the Transcription Coactivator Pygophus
Authors : Padmanabhan, B.; Yokoyama, S.; RIKEN Structural Genomics/Proteomics Initiative (RSGI)
Deposited on : 2006-08-24
Resolution : 2.70 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Xtriage (Phenix)	:	NOT EXECUTED
EDS	:	NOT EXECUTED
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.70 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
Clashscore	190562	3843 (2.70-2.70)
Ramachandran outliers	187476	3778 (2.70-2.70)
Sidechain outliers	187428	3778 (2.70-2.70)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$.

Note EDS was not executed.

Mol	Chain	Length	Quality of chain
1	A	67	
1	B	67	

2 Entry composition

There are 3 unique types of molecules in this entry. The entry contains 971 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Pygopus homolog 1.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	60	Total	C	N	O	S	0	0	0
			450	276	72	93	9			
1	B	60	Total	C	N	O	S	0	0	0
			450	276	72	93	9			

- Molecule 2 is ZINC ION (CCD ID: ZN) (formula: Zn).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	A	2	Total	Zn	0	0
			2	2		
2	B	2	Total	Zn	0	0
			2	2		

- Molecule 3 is water.


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	32	Total	O	0	0
			32	32		
3	B	35	Total	O	0	0
			35	35		


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

Note EDS was not executed.

• Molecule 1: Pygopus homolog 1

Chain A: 



• Molecule 1: Pygopus homolog 1

Chain B: 



4 Data and refinement statistics

Xtriage (Phenix) and EDS were not executed - this section is therefore incomplete.

Property	Value	Source
Space group	P 43 21 2	Depositor
Cell constants a, b, c, α , β , γ	60.08 Å 60.08 Å 95.24 Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	20.00 – 2.70	Depositor
% Data completeness (in resolution range)	96.9 (20.00-2.70)	Depositor
R_{merge}	0.09	Depositor
R_{sym}	(Not available)	Depositor
Refinement program	REFMAC 5.2.0005	Depositor
R, R_{free}	0.197 , 0.255	Depositor
Estimated twinning fraction	No twinning to report.	Xtriage
Total number of atoms	971	wwPDB-VP
Average B, all atoms (Å ²)	52.0	wwPDB-VP

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: ZN

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	1.89	12/459 (2.6%)	1.24	1/624 (0.2%)
1	B	1.93	9/459 (2.0%)	1.36	8/624 (1.3%)
All	All	1.91	21/918 (2.3%)	1.30	9/1248 (0.7%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	1

All (21) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	338	VAL	N-CA	8.95	1.57	1.46
1	B	338	VAL	CA-CB	8.94	1.64	1.54
1	B	362	GLN	CA-C	-8.09	1.43	1.53
1	B	355	ILE	C-O	-7.38	1.16	1.24
1	A	338	VAL	CA-CB	6.90	1.62	1.54
1	A	366	HIS	CA-C	-6.49	1.44	1.52
1	A	367	ARG	C-O	-6.30	1.16	1.24
1	B	349	ASN	CA-C	-5.87	1.45	1.53
1	A	354	ALA	CA-C	5.58	1.59	1.52
1	A	356	LEU	CA-C	-5.57	1.45	1.52
1	B	383	GLU	N-CA	5.41	1.53	1.46
1	B	366	HIS	CA-C	-5.39	1.45	1.52
1	A	379	LEU	N-CA	-5.36	1.39	1.46
1	A	369	CYS	CB-SG	-5.32	1.63	1.81
1	A	374	GLU	C-O	-5.31	1.17	1.24
1	B	361	CYS	CA-C	-5.27	1.45	1.52

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	369	CYS	C-O	-5.23	1.17	1.24
1	A	355	ILE	CA-CB	-5.22	1.48	1.54
1	A	382	ALA	CA-CB	-5.18	1.45	1.53
1	B	337	PRO	CA-C	5.09	1.63	1.52
1	B	337	PRO	N-CA	5.01	1.54	1.47

All (9) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	338	VAL	N-CA-C	11.06	120.80	110.74
1	B	361	CYS	CA-C-N	-8.12	110.73	122.67
1	B	361	CYS	C-N-CA	-8.12	110.73	122.67
1	B	383	GLU	N-CA-C	6.21	119.21	109.96
1	B	361	CYS	N-CA-C	-5.31	99.49	110.80
1	B	339	TYR	CA-C-N	-5.25	114.20	120.13
1	B	339	TYR	C-N-CA	-5.25	114.20	120.13
1	B	369	CYS	N-CA-C	5.15	118.35	111.75
1	B	366	HIS	N-CA-C	-5.06	102.37	109.96

There are no chirality outliers.

All (1) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	337	PRO	Peptide

5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	450	0	404	11	0
1	B	450	0	404	14	0
2	A	2	0	0	0	0
2	B	2	0	0	0	0
3	A	32	0	0	2	0
3	B	35	0	0	3	0
All	All	971	0	808	25	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 15.

All (25) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:373:THR:OG1	3:B:52:HOH:O	1.52	1.21
1:A:377:TYR:O	1:A:381:THR:HG23	1.67	0.95
1:B:344:CYS:O	1:B:345:THR:HB	1.64	0.95
1:A:394:MET:HG2	3:B:27:HOH:O	1.82	0.78
1:B:373:THR:CB	3:B:52:HOH:O	2.22	0.74
1:B:360:SER:HB3	1:B:393:CYS:SG	2.32	0.70
1:A:396:ASP:HB3	3:A:12:HOH:O	1.93	0.67
1:B:360:SER:C	1:B:361:CYS:O	2.31	0.64
1:A:358:GLU:HG3	3:A:10:HOH:O	2.02	0.60
1:B:344:CYS:O	1:B:345:THR:CB	2.42	0.56
1:A:366:HIS:O	1:A:370:THR:HG23	2.07	0.54
1:B:344:CYS:SG	1:B:346:ASN:ND2	2.82	0.52
1:B:379:LEU:O	1:B:379:LEU:HD22	2.10	0.52
1:B:360:SER:O	1:B:361:CYS:O	2.29	0.50
1:B:359:ALA:HB2	1:B:394:MET:HE3	1.94	0.49
1:B:345:THR:HG22	1:B:345:THR:O	2.13	0.48
1:A:353:ASP:HB3	1:A:367:ARG:HB3	1.94	0.48
1:A:359:ALA:O	1:A:360:SER:HB2	2.14	0.47
1:B:359:ALA:CB	1:B:394:MET:HE3	2.44	0.47
1:A:349:ASN:HB2	1:A:351:ASP:OD2	2.14	0.47
1:B:345:THR:O	1:B:345:THR:CG2	2.62	0.47
1:A:380:LEU:HD11	1:A:388:TRP:HB2	1.97	0.46
1:A:383:GLU:HA	1:A:383:GLU:OE1	2.15	0.46
1:A:360:SER:O	1:A:362:GLN:HG2	2.18	0.43
1:B:360:SER:O	1:B:361:CYS:C	2.65	0.40

There are no symmetry-related clashes.

5.3 Torsion angles ⓘ

5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was

analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	58/67 (87%)	52 (90%)	4 (7%)	2 (3%)	3	7
1	B	58/67 (87%)	52 (90%)	3 (5%)	3 (5%)	1	3
All	All	116/134 (87%)	104 (90%)	7 (6%)	5 (4%)	2	4

All (5) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	360	SER
1	B	361	CYS
1	B	382	ALA
1	A	353	ASP
1	B	360	SER

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	49/55 (89%)	38 (78%)	11 (22%)	1	3
1	B	49/55 (89%)	45 (92%)	4 (8%)	10	27
All	All	98/110 (89%)	83 (85%)	15 (15%)	3	7

All (15) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	345	THR
1	A	349	ASN
1	A	350	ASP
1	A	355	ILE
1	A	367	ARG
1	A	368	ILE
1	A	379	LEU
1	A	380	LEU
1	A	381	THR

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	A	391	ASP
1	A	394	MET
1	B	355	ILE
1	B	356	LEU
1	B	379	LEU
1	B	383	GLU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (1) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	349	ASN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 4 ligands modelled in this entry, 4 are monoatomic - leaving 0 for Mogul analysis.

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers

There are no such residues in this entry.

5.8 Polymer linkage issues

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

EDS was not executed - this section is therefore empty.

6.2 Non-standard residues in protein, DNA, RNA chains

EDS was not executed - this section is therefore empty.

6.3 Carbohydrates

EDS was not executed - this section is therefore empty.

6.4 Ligands

EDS was not executed - this section is therefore empty.

6.5 Other polymers

EDS was not executed - this section is therefore empty.