



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 5, 2026 – 05:56 AM UTC

PDB ID : 4DX8 / pdb_00004dx8
Title : ICAP1 in complex with KRIT1 N-terminus
Authors : Liu, W.; Draheim, K.; Zhang, R.; Calderwood, D.A.; Boggon, T.J.
Deposited on : 2012-02-27
Resolution : 2.54 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.54 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	1091 (2.54-2.54)
Clashscore	190562	1120 (2.54-2.54)
Ramachandran outliers	187476	1106 (2.54-2.54)
Sidechain outliers	187428	1106 (2.54-2.54)
RSRZ outliers	180081	1091 (2.54-2.54)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	154	5% 82% 8% 10%
1	B	154	5% 80% 7% 12%
1	D	154	8% 75% 10% 14%
1	E	154	6% 77% 8% 14%
2	H	203	8% 69% 8% 22%

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
2	I	203	10% 48% 6% 45%
2	J	203	16% 51% 7% 41%
2	K	203	16% 37% 5% 57%

2 Entry composition

There are 4 unique types of molecules in this entry. The entry contains 8128 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Integrin beta-1-binding protein 1.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	139	Total	C	N	O	S	0	0	0
			1080	690	172	212	6			
1	B	136	Total	C	N	O	S	0	0	0
			1061	680	168	207	6			
1	D	133	Total	C	N	O	S	0	0	0
			1038	665	164	203	6			
1	E	133	Total	C	N	O	S	0	0	0
			1038	665	164	203	6			

There are 8 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	47	GLY	-	expression tag	UNP O14713
A	48	SER	-	expression tag	UNP O14713
B	47	GLY	-	expression tag	UNP O14713
B	48	SER	-	expression tag	UNP O14713
D	47	GLY	-	expression tag	UNP O14713
D	48	SER	-	expression tag	UNP O14713
E	47	GLY	-	expression tag	UNP O14713
E	48	SER	-	expression tag	UNP O14713

- Molecule 2 is a protein called Krev interaction trapped protein 1.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	H	159	Total	C	N	O	S	0	0	0
			1278	825	221	225	7			
2	I	111	Total	C	N	O	S	0	0	0
			906	590	159	152	5			
2	J	120	Total	C	N	O	S	0	0	0
			979	634	166	176	3			
2	K	88	Total	C	N	O	S	0	0	0
			718	465	126	125	2			

There are 20 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
H	-4	GLY	-	expression tag	UNP O00522
H	-3	PRO	-	expression tag	UNP O00522
H	-2	LEU	-	expression tag	UNP O00522
H	-1	GLY	-	expression tag	UNP O00522
H	0	SER	-	expression tag	UNP O00522
I	-4	GLY	-	expression tag	UNP O00522
I	-3	PRO	-	expression tag	UNP O00522
I	-2	LEU	-	expression tag	UNP O00522
I	-1	GLY	-	expression tag	UNP O00522
I	0	SER	-	expression tag	UNP O00522
J	-4	GLY	-	expression tag	UNP O00522
J	-3	PRO	-	expression tag	UNP O00522
J	-2	LEU	-	expression tag	UNP O00522
J	-1	GLY	-	expression tag	UNP O00522
J	0	SER	-	expression tag	UNP O00522
K	-4	GLY	-	expression tag	UNP O00522
K	-3	PRO	-	expression tag	UNP O00522
K	-2	LEU	-	expression tag	UNP O00522
K	-1	GLY	-	expression tag	UNP O00522
K	0	SER	-	expression tag	UNP O00522

- Molecule 3 is BROMIDE ION (CCD ID: BR) (formula: Br).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	A	3	Total Br 3 3	0	0
3	B	3	Total Br 3 3	0	0
3	D	4	Total Br 4 4	0	0
3	E	3	Total Br 3 3	0	0
3	H	3	Total Br 3 3	0	0

- Molecule 4 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
4	A	4	Total O 4 4	0	0
4	D	1	Total O 1 1	0	0

Continued on next page...


Continued from previous page...

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	E	7	Total 7	O 7	0	0
4	H	1	Total 1	O 1	0	0
4	J	1	Total 1	O 1	0	0


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.



- Molecule 1: Integrin beta-1-binding protein 1





- Molecule 1: Integrin beta-1-binding protein 1



- Molecule 1: Integrin beta-1-binding protein 1




- Molecule 1: Integrin beta-1-binding protein 1




- Molecule 2: Krev interaction trapped protein 1








● Molecule 2: Krev interaction trapped protein 1



● Molecule 2: Krev interaction trapped protein 1



● Molecule 2: Krev interaction trapped protein 1



4 Data and refinement statistics

Property	Value	Source
Space group	C 2 2 21	Depositor
Cell constants a, b, c, α , β , γ	153.90Å 157.98Å 152.01Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	50.00 – 2.54 50.00 – 2.54	Depositor EDS
% Data completeness (in resolution range)	100.0 (50.00-2.54) 99.6 (50.00-2.54)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	0.09	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.59 (at 2.54Å)	Xtriage
Refinement program	REFMAC 5.6.0117	Depositor
R, R_{free}	0.220 , 0.252 0.218 , 0.249	Depositor DCC
R_{free} test set	3101 reflections (5.06%)	wwPDB-VP
Wilson B-factor (Å ²)	83.6	Xtriage
Anisotropy	0.053	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.32 , 58.6	EDS
L-test for twinning ²	$\langle L \rangle = 0.47$, $\langle L^2 \rangle = 0.30$	Xtriage
Estimated twinning fraction	0.034 for -k,-h,-l	Xtriage
F_o, F_c correlation	0.94	EDS
Total number of atoms	8128	wwPDB-VP
Average B, all atoms (Å ²)	102.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.34% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: BR

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.55	0/1098	0.75	0/1486
1	B	0.54	0/1079	0.75	0/1460
1	D	0.56	0/1056	0.75	0/1430
1	E	0.55	0/1056	0.71	0/1430
2	H	0.59	0/1299	0.74	0/1750
2	I	0.56	1/920 (0.1%)	0.73	0/1239
2	J	0.51	0/992	0.73	2/1334 (0.1%)
2	K	0.58	0/728	0.76	0/980
All	All	0.56	1/8228 (0.0%)	0.74	2/11109 (0.0%)

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	I	22	SER	CB-OG	5.05	1.52	1.42

All (2) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	J	113	VAL	CA-C-N	5.05	124.81	119.76
2	J	113	VAL	C-N-CA	5.05	124.81	119.76

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within

the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1080	0	1070	6	0
1	B	1061	0	1057	8	0
1	D	1038	0	1028	10	0
1	E	1038	0	1028	11	0
2	H	1278	0	1322	20	0
2	I	906	0	953	6	0
2	J	979	0	1010	9	0
2	K	718	0	740	7	0
3	A	3	0	0	1	0
3	B	3	0	0	1	0
3	D	4	0	0	1	0
3	E	3	0	0	1	0
3	H	3	0	0	0	0
4	A	4	0	0	0	0
4	D	1	0	0	0	0
4	E	7	0	0	0	0
4	H	1	0	0	0	0
4	J	1	0	0	0	0
All	All	8128	0	8208	67	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 4.

All (67) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:H:100:ASP:HB3	2:H:101:GLY:CA	1.77	1.15
2:H:100:ASP:HB3	2:H:101:GLY:HA3	1.30	1.09
2:H:147:THR:HA	2:H:148:HIS:HB2	1.48	0.93
1:D:140:ARG:NH1	3:D:302:BR:BR	2.61	0.88
1:E:140:ARG:NH1	3:E:302:BR:BR	2.62	0.87
2:J:94:MET:HE3	2:J:112:ILE:HD11	1.62	0.79
1:A:140:ARG:NH1	3:A:302:BR:BR	2.73	0.76
2:H:147:THR:CA	2:H:148:HIS:HB2	2.17	0.73
2:H:100:ASP:HB3	2:H:101:GLY:HA2	1.71	0.73
1:B:140:ARG:NH1	3:B:302:BR:BR	2.77	0.72
2:H:100:ASP:CB	2:H:101:GLY:CA	2.61	0.72
1:D:190:ALA:O	1:D:194:VAL:HG13	1.90	0.71
1:D:141:MET:HE3	1:D:184:CYS:SG	2.31	0.71
1:D:143:CYS:HB2	1:D:155:LEU:HD23	1.76	0.67

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:E:93:ASP:OD2	2:K:185:ARG:NH2	2.31	0.64
1:E:160:THR:HG21	1:E:164:ASN:OD1	1.99	0.63
2:H:83:ASN:N	2:J:62:SER:HG	1.97	0.63
2:I:170:GLN:O	2:I:171:SER:HB3	2.02	0.60
2:H:170:GLN:O	2:H:171:SER:HB3	2.02	0.60
1:A:134:ALA:HB1	1:E:134:ALA:HB1	1.84	0.59
2:H:100:ASP:CB	2:H:101:GLY:HA3	2.18	0.59
1:B:145:ASP:O	2:H:185:ARG:HD2	2.01	0.59
1:D:93:ASP:OD2	2:I:185:ARG:NH2	2.35	0.59
1:B:59:THR:HG21	1:B:116:LYS:HE2	1.83	0.59
2:H:94:MET:HE3	2:H:112:ILE:HD11	1.86	0.57
1:B:160:THR:HG21	1:B:164:ASN:OD1	2.04	0.57
2:H:94:MET:HE1	2:H:160:LEU:CD1	2.38	0.54
1:E:145:ASP:O	2:K:185:ARG:HD2	2.08	0.54
1:A:145:ASP:O	2:J:185:ARG:HD2	2.08	0.53
1:D:143:CYS:HB2	1:D:155:LEU:CD2	2.37	0.53
1:D:145:ASP:O	2:I:185:ARG:HD2	2.08	0.53
1:B:141:MET:HE3	1:B:184:CYS:SG	2.49	0.53
1:A:93:ASP:OD2	2:J:185:ARG:NH2	2.41	0.52
2:H:147:THR:HA	2:H:148:HIS:CB	2.27	0.52
1:A:141:MET:HE1	1:A:183:ILE:HG22	1.90	0.52
1:D:135:LEU:CD2	1:D:194:VAL:HG11	2.41	0.51
2:J:170:GLN:O	2:J:171:SER:HB3	2.11	0.50
2:H:163:TRP:O	2:H:167:ARG:HG2	2.12	0.50
1:E:141:MET:HE3	1:E:184:CYS:SG	2.53	0.49
2:J:94:MET:HE1	2:J:160:LEU:HA	1.94	0.49
1:E:93:ASP:CG	2:K:185:ARG:HH22	2.21	0.49
1:E:139:ILE:HD11	1:E:160:THR:HG23	1.95	0.48
2:J:89:LYS:HE2	2:J:116:VAL:HG23	1.95	0.48
1:B:141:MET:HE1	1:B:183:ILE:HG22	1.95	0.48
2:I:65:THR:HG23	2:I:91:VAL:HB	1.95	0.48
2:K:171:SER:C	2:K:173:PHE:H	2.23	0.47
1:D:135:LEU:HD23	1:D:194:VAL:HG11	1.95	0.47
2:H:94:MET:HE1	2:H:160:LEU:HA	1.95	0.47
2:H:94:MET:HE1	2:H:160:LEU:HD12	1.95	0.47
1:D:82:LEU:HD12	1:D:87:ASP:HB3	1.99	0.45
2:H:76:THR:HG22	2:H:76:THR:O	2.18	0.44
2:H:76:THR:HG21	2:H:86:ILE:HG12	2.00	0.44
2:H:76:THR:HG21	2:H:86:ILE:CG1	2.48	0.44
2:H:170:GLN:O	2:H:171:SER:CB	2.65	0.43
2:I:94:MET:HE1	2:I:160:LEU:HA	2.00	0.43

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:76:LEU:HD13	1:B:165:GLU:HA	2.01	0.42
2:K:94:MET:HE1	2:K:160:LEU:HD12	2.01	0.42
2:J:98:PRO:HA	2:J:107:GLU:HG2	2.01	0.42
2:I:129:CYS:HB3	2:I:130:PRO:HD3	2.00	0.42
2:K:174:ILE:HB	2:K:175:PRO:HD3	2.01	0.42
1:A:56:ASN:HA	1:A:57:SER:HA	1.91	0.42
2:J:54:LEU:HB3	2:J:71:TYR:HE2	1.85	0.42
2:K:110:LEU:HD22	2:K:156:MET:HB3	2.01	0.42
1:E:141:MET:HE1	1:E:183:ILE:HG22	2.01	0.42
1:E:160:THR:HG22	1:E:166:GLU:O	2.19	0.41
1:B:111:ILE:HD12	1:B:126:GLN:HG3	2.03	0.41
1:E:178:GLU:CD	1:E:178:GLU:H	2.28	0.41

There are no symmetry-related clashes.

5.3 Torsion angles ⓘ

5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	135/154 (88%)	130 (96%)	5 (4%)	0	100	100
1	B	132/154 (86%)	128 (97%)	4 (3%)	0	100	100
1	D	129/154 (84%)	126 (98%)	2 (2%)	1 (1%)	16	22
1	E	129/154 (84%)	127 (98%)	2 (2%)	0	100	100
2	H	145/203 (71%)	135 (93%)	9 (6%)	1 (1%)	18	25
2	I	97/203 (48%)	94 (97%)	2 (2%)	1 (1%)	12	17
2	J	104/203 (51%)	96 (92%)	7 (7%)	1 (1%)	12	17
2	K	74/203 (36%)	71 (96%)	2 (3%)	1 (1%)	9	11
All	All	945/1428 (66%)	907 (96%)	33 (4%)	5 (0%)	24	34

All (5) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	D	126	GLN
2	K	171	SER
2	H	100	ASP
2	I	171	SER
2	J	171	SER

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	119/132 (90%)	115 (97%)	4 (3%)	32	48
1	B	117/132 (89%)	115 (98%)	2 (2%)	53	72
1	D	114/132 (86%)	109 (96%)	5 (4%)	25	38
1	E	114/132 (86%)	109 (96%)	5 (4%)	25	38
2	H	142/179 (79%)	137 (96%)	5 (4%)	32	48
2	I	102/179 (57%)	98 (96%)	4 (4%)	28	43
2	J	109/179 (61%)	107 (98%)	2 (2%)	51	70
2	K	79/179 (44%)	76 (96%)	3 (4%)	29	44
All	All	896/1244 (72%)	866 (97%)	30 (3%)	33	50

All (30) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	76	LEU
1	A	130	LEU
1	A	136	TYR
1	A	175	ASN
1	B	59	THR
1	B	160	THR
1	D	59	THR
1	D	125	ASP
1	D	130	LEU
1	D	157	LEU
1	D	194	VAL

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	E	125	ASP
1	E	135	LEU
1	E	136	TYR
1	E	160	THR
1	E	178	GLU
2	H	23	LEU
2	H	56	THR
2	H	68	ILE
2	H	79	ILE
2	H	171	SER
2	I	23	LEU
2	I	76	THR
2	I	191	ILE
2	I	197	THR
2	J	8	GLU
2	J	56	THR
2	K	116	VAL
2	K	168	HIS
2	K	197	THR

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (9) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	181	GLN
1	B	175	ASN
1	D	175	ASN
1	D	181	GLN
1	E	175	ASN
1	E	181	GLN
2	H	170	GLN
2	I	170	GLN
2	K	189	ASN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 16 ligands modelled in this entry, 16 are monoatomic - leaving 0 for Mogul analysis.

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ > 2		OWAB(Å ²)	Q < 0.9
1	A	139/154 (90%)	0.41	8 (5%)	29 28	58, 73, 129, 152	0
1	B	136/154 (88%)	0.25	8 (5%)	28 27	57, 72, 118, 149	0
1	D	133/154 (86%)	0.44	12 (9%)	15 14	59, 73, 113, 158	0
1	E	133/154 (86%)	0.31	9 (6%)	23 22	58, 76, 113, 150	0
2	H	159/203 (78%)	0.64	17 (10%)	11 10	57, 91, 140, 162	0
2	I	111/203 (54%)	1.18	21 (18%)	3 2	57, 143, 208, 254	0
2	J	120/203 (59%)	1.37	33 (27%)	1 1	58, 143, 197, 219	0
2	K	88/203 (43%)	1.52	32 (36%)	1 0	58, 151, 210, 256	0
All	All	1019/1428 (71%)	0.71	140 (13%)	6 5	57, 87, 186, 256	0

All (140) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
2	I	24	ASN	6.7
2	J	115	SER	6.5
2	K	33	TYR	6.4
1	A	76	LEU	6.2
1	A	55	ASN	5.7
2	H	117	VAL	5.2
2	J	64	ILE	4.7
2	I	78	PRO	4.6
2	K	13	ALA	4.6
2	J	58	LEU	4.6
1	D	150	ALA	4.5
2	J	12	VAL	4.3
2	K	110	LEU	4.3
2	J	116	VAL	4.1
1	A	75	LYS	4.0
2	I	36	LEU	4.0

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
2	H	147	THR	3.9
2	I	79	ILE	3.9
2	K	92	VAL	3.7
2	K	72	VAL	3.7
2	K	23	LEU	3.7
2	J	73	VAL	3.7
2	K	73	VAL	3.6
2	J	26	ARG	3.5
2	K	94	MET	3.5
2	J	36	LEU	3.5
1	B	194	VAL	3.4
2	I	64	ILE	3.4
2	I	117	VAL	3.4
2	J	68	ILE	3.4
2	K	64	ILE	3.4
2	K	197	THR	3.3
1	D	128	ASP	3.3
1	D	60	CYS	3.3
1	B	76	LEU	3.3
1	B	81	GLY	3.3
1	D	127	TYR	3.2
2	K	30	ALA	3.2
2	J	54	LEU	3.2
2	K	69	LEU	3.2
2	J	14	VAL	3.2
2	J	72	VAL	3.2
1	D	81	GLY	3.1
2	K	76	THR	3.1
2	J	10	ALA	3.1
2	J	53	LEU	3.1
2	I	111	PHE	3.1
2	H	79	ILE	3.1
2	K	116	VAL	3.1
2	J	154	ALA	3.0
2	J	157	LEU	3.0
1	D	59	THR	3.0
2	J	197	THR	3.0
2	K	115	SER	3.0
2	K	71	TYR	3.0
2	H	103	LYS	3.0
1	E	60	CYS	3.0
2	H	101	GLY	3.0

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
2	J	110	LEU	2.9
2	K	157	LEU	2.9
1	D	196	THR	2.9
1	E	196	THR	2.9
1	E	59	THR	2.8
2	H	99	LEU	2.8
2	J	111	PHE	2.8
2	H	104	MET	2.8
2	K	112	ILE	2.8
2	I	25	SER	2.8
2	H	142	CYS	2.8
2	H	105	GLY	2.7
2	J	11	TYR	2.7
1	B	196	THR	2.7
2	K	93	LEU	2.7
2	H	40	VAL	2.7
2	J	153	THR	2.7
2	J	166	GLU	2.7
2	J	158	ILE	2.7
2	J	24	ASN	2.7
2	I	194	ALA	2.7
2	J	18	LYS	2.7
2	I	23	LEU	2.7
2	H	195	TYR	2.7
2	I	18	LYS	2.6
2	I	68	ILE	2.6
2	K	162	LYS	2.6
2	H	197	THR	2.6
2	J	65	THR	2.6
1	D	195	LEU	2.6
2	I	69	LEU	2.6
1	A	196	THR	2.6
2	K	154	ALA	2.5
2	H	100	ASP	2.5
2	K	14	VAL	2.5
1	A	77	SER	2.5
1	A	81	GLY	2.5
1	D	149	GLY	2.5
2	K	158	ILE	2.5
2	K	75	THR	2.5
2	K	68	ILE	2.5
1	E	100	LYS	2.5

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	59	THR	2.5
2	H	63	GLU	2.4
1	E	195	LEU	2.4
2	I	33	TYR	2.4
2	K	91	VAL	2.4
2	I	94	MET	2.4
2	J	22	SER	2.4
1	D	74	LEU	2.4
2	I	136	GLN	2.4
1	A	56	ASN	2.4
2	H	129	CYS	2.4
2	J	156	MET	2.4
2	K	89	LYS	2.4
2	K	26	ARG	2.3
2	K	18	LYS	2.3
1	B	150	ALA	2.3
2	J	106	ARG	2.3
2	I	71	TYR	2.3
2	J	137	ASP	2.3
1	D	143	CYS	2.2
1	E	127	TYR	2.2
1	E	74	LEU	2.2
1	E	129	VAL	2.2
1	B	195	LEU	2.2
2	J	93	LEU	2.2
1	E	150	ALA	2.1
2	K	66	GLN	2.1
1	B	130	LEU	2.1
2	I	75	THR	2.1
2	I	12	VAL	2.1
2	J	62	SER	2.1
2	J	138	ILE	2.1
2	I	168	HIS	2.1
1	D	146	ASP	2.1
2	H	41	PRO	2.1
2	K	111	PHE	2.0
2	H	58	LEU	2.0
2	K	163	TRP	2.0
1	B	77	SER	2.0
2	I	17	PRO	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
3	BR	A	303	1/1	0.87	0.12	162,162,162,162	0
3	BR	E	303	1/1	0.89	0.17	139,139,139,139	1
3	BR	B	303	1/1	0.92	0.09	146,146,146,146	0
3	BR	H	303	1/1	0.92	0.09	161,161,161,161	0
3	BR	H	302	1/1	0.94	0.08	146,146,146,146	0
3	BR	H	301	1/1	0.94	0.07	135,135,135,135	0
3	BR	B	302	1/1	0.95	0.07	105,105,105,105	0
3	BR	D	304	1/1	0.95	0.11	166,166,166,166	0
3	BR	D	301	1/1	0.96	0.07	120,120,120,120	0
3	BR	E	301	1/1	0.96	0.06	127,127,127,127	0
3	BR	D	302	1/1	0.96	0.07	95,95,95,95	0
3	BR	D	303	1/1	0.97	0.07	133,133,133,133	0
3	BR	E	302	1/1	0.97	0.07	92,92,92,92	0
3	BR	A	302	1/1	0.98	0.08	97,97,97,97	0
3	BR	A	301	1/1	0.98	0.04	119,119,119,119	0
3	BR	B	301	1/1	0.98	0.05	121,121,121,121	0

6.5 Other polymers [i](#)

There are no such residues in this entry.