



wwPDB X-ray Structure Validation Summary Report ⓘ

Apr 25, 2026 – 01:04 PM EDT

PDB ID : 5DX8 / pdb_00005dx8
Title : Crystal structure of CARM1, sinefungin, and methylated PABP1 peptide (R455)
Authors : Boriack-Sjodin, P.A.
Deposited on : 2015-09-23
Resolution : 1.94 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0
Mogul : 2022.3.0, CSD as543be (2022)
Xtriage (Phenix) : 2.0
EDS : 3.0
Buster-report : wwPDB partial adaption of 1.1.7 (2018)
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4 : 9.0.010 (Gargrove)
Density-Fitness : 1.0.12
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.94 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	1452 (1.94-1.94)
Clashscore	190562	1494 (1.94-1.94)
Ramachandran outliers	187476	1479 (1.94-1.94)
Sidechain outliers	187428	1479 (1.94-1.94)
RSRZ outliers	180081	1453 (1.94-1.94)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	349	2% 89% 9%
1	B	349	% 90% 8%
1	C	349	3% 91% 7%
1	D	349	5% 90% 8%
2	E	19	11% 37% 11% 53%

Continued on next page...

Mol	Chain	Length	Quality of chain
2	F	19	11% 42% 5% 53%
2	G	19	32% 68%
2	H	19	5% 42% 58%

2 Entry composition

There are 4 unique types of molecules in this entry. The entry contains 11891 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Histone-arginine methyltransferase CARM1.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	344	Total	C	N	O	S	0	10	0
			2823	1821	466	520	16			
1	B	344	Total	C	N	O	S	0	8	0
			2802	1810	461	516	15			
1	C	343	Total	C	N	O	S	0	5	0
			2782	1796	461	509	16			
1	D	343	Total	C	N	O	S	0	5	0
			2787	1796	462	514	15			

There are 12 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	131	SER	-	expression tag	UNP Q86X55
A	132	ILE	-	expression tag	UNP Q86X55
A	133	ALA	-	expression tag	UNP Q86X55
B	131	SER	-	expression tag	UNP Q86X55
B	132	ILE	-	expression tag	UNP Q86X55
B	133	ALA	-	expression tag	UNP Q86X55
C	131	SER	-	expression tag	UNP Q86X55
C	132	ILE	-	expression tag	UNP Q86X55
C	133	ALA	-	expression tag	UNP Q86X55
D	131	SER	-	expression tag	UNP Q86X55
D	132	ILE	-	expression tag	UNP Q86X55
D	133	ALA	-	expression tag	UNP Q86X55

- Molecule 2 is a protein called methylated PABP1 peptide.

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
2	E	9	Total	C	N	O	0	0	0
			65	41	15	9			
2	F	9	Total	C	N	O	0	0	0
			68	44	15	9			

Continued on next page...


Continued from previous page...

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
2	G	6	Total	C	N	O	0	0	0
			44	29	9	6			
2	H	8	Total	C	N	O	0	0	0
			63	41	14	8			

There are 4 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
E	19	NH2	-	amidation	UNP P11940
F	19	NH2	-	amidation	UNP P11940
G	19	NH2	-	amidation	UNP P11940
H	19	NH2	-	amidation	UNP P11940


- Molecule 3 is SINEFUNGIN (CCD ID: SFG) (formula: $C_{15}H_{23}N_7O_5$).





Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
3	A	1	Total	C	N	O	0	0
			27	15	7	5		
3	B	1	Total	C	N	O	0	0
			27	15	7	5		
3	C	1	Total	C	N	O	0	0
			27	15	7	5		
3	D	1	Total	C	N	O	0	0
			27	15	7	5		

- Molecule 4 is water.


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	97	Total 97	O 97	0	0
4	B	105	Total 105	O 105	0	0
4	C	62	Total 62	O 62	0	0
4	D	80	Total 80	O 80	0	0
4	E	1	Total 1	O 1	0	0
4	F	3	Total 3	O 3	0	0
4	G	1	Total 1	O 1	0	0



- Molecule 2: methylated PABP1 peptide




- Molecule 2: methylated PABP1 peptide



- Molecule 2: methylated PABP1 peptide



- Molecule 2: methylated PABP1 peptide



4 Data and refinement statistics

Property	Value	Source
Space group	P 2 ₁ 2 ₁ 2	Depositor
Cell constants a, b, c, α , β , γ	74.86Å 98.27Å 207.54Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	50.00 – 1.94 50.00 – 1.94	Depositor EDS
% Data completeness (in resolution range)	90.1 (50.00-1.94) 90.1 (50.00-1.94)	Depositor EDS
R_{merge}	0.12	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.13 (at 1.94Å)	Xtriage
Refinement program	REFMAC 5.8.0073	Depositor
R, R_{free}	0.204 , 0.246 0.210 , 0.250	Depositor DCC
R_{free} test set	5156 reflections (4.53%)	wwPDB-VP
Wilson B-factor (Å ²)	22.6	Xtriage
Anisotropy	0.095	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.37 , 37.0	EDS
L-test for twinning ²	$\langle L \rangle = 0.46$, $\langle L^2 \rangle = 0.28$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.94	EDS
Total number of atoms	11891	wwPDB-VP
Average B, all atoms (Å ²)	26.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The analyses of the Patterson function reveals a significant off-origin peak that is 38.30 % of the origin peak, indicating pseudo-translational symmetry. The chance of finding a peak of this or larger height randomly in a structure without pseudo-translational symmetry is equal to 3.7649e-04. The detected translational NCS is most likely also responsible for the elevated intensity ratio.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: SFG, NMM

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.76	0/2915	0.87	1/3947 (0.0%)
1	B	0.76	0/2897	0.87	1/3926 (0.0%)
1	C	0.68	0/2867	0.81	0/3882
1	D	0.69	0/2860	0.83	0/3874
2	E	0.67	0/41	0.91	0/54
2	F	0.51	0/44	0.67	0/57
2	G	0.65	0/32	0.89	0/42
2	H	0.59	0/39	0.62	0/50
All	All	0.72	0/11695	0.84	2/15832 (0.0%)

There are no bond length outliers.

All (2) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	197	ILE	N-CA-C	5.63	116.28	110.36
1	A	197	ILE	N-CA-C	5.21	116.57	110.62

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2823	0	2780	25	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	B	2802	0	2762	16	0
1	C	2782	0	2744	20	0
1	D	2787	0	2735	25	0
2	E	65	0	66	2	0
2	F	68	0	71	2	0
2	G	44	0	46	0	0
2	H	63	0	66	0	0
3	A	27	0	22	0	0
3	B	27	0	22	0	0
3	C	27	0	22	0	0
3	D	27	0	22	0	0
4	A	97	0	0	7	0
4	B	105	0	0	6	0
4	C	62	0	0	5	0
4	D	80	0	0	12	0
4	E	1	0	0	0	0
4	F	3	0	0	0	0
4	G	1	0	0	0	0
All	All	11891	0	11358	84	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 4.

The worst 5 of 84 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:C:349:MET:HE1	1:C:384:VAL:HG22	1.68	0.75
1:A:364:GLU:OE1	4:A:601:HOH:O	2.03	0.75
1:C:284[B]:ASN:ND2	4:C:601:HOH:O	2.19	0.75
1:A:304:MET:HE1	1:C:304:MET:HE2	1.69	0.73
1:A:336:ARG:HD3	4:A:692:HOH:O	1.89	0.73

There are no symmetry-related clashes.

5.3 Torsion angles ⓘ

5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	352/349 (101%)	340 (97%)	12 (3%)	0	100	100
1	B	350/349 (100%)	339 (97%)	9 (3%)	2 (1%)	21	11
1	C	346/349 (99%)	332 (96%)	13 (4%)	1 (0%)	36	30
1	D	346/349 (99%)	334 (96%)	12 (4%)	0	100	100
2	E	6/19 (32%)	4 (67%)	2 (33%)	0	100	100
2	F	5/19 (26%)	4 (80%)	1 (20%)	0	100	100
2	G	3/19 (16%)	3 (100%)	0	0	100	100
2	H	4/19 (21%)	4 (100%)	0	0	100	100
All	All	1412/1472 (96%)	1360 (96%)	49 (4%)	3 (0%)	43	37

All (3) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	B	266	GLU
1	B	258	PRO
1	C	258	PRO

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	310/303 (102%)	305 (98%)	5 (2%)	55	46
1	B	308/303 (102%)	304 (99%)	4 (1%)	61	54
1	C	304/303 (100%)	302 (99%)	2 (1%)	76	74
1	D	304/303 (100%)	300 (99%)	4 (1%)	61	54
2	E	3/12 (25%)	3 (100%)	0	100	100
2	F	4/12 (33%)	4 (100%)	0	100	100
2	G	3/12 (25%)	3 (100%)	0	100	100
2	H	4/12 (33%)	4 (100%)	0	100	100

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
All	All	1240/1260 (98%)	1225 (99%)	15 (1%)	63 57

5 of 15 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	B	353	VAL
1	D	286	PHE
1	B	424	SER
1	D	424	SER
1	D	139	GLU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 15 such sidechains are listed below:

Mol	Chain	Res	Type
1	B	180	HIS
1	D	220	GLN
1	B	465	ASN
1	D	306	GLN
1	C	160	GLN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

7 non-standard protein/DNA/RNA residues are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
2	NMM	F	7	2	8,11,12	0.54	0	7,12,14	0.96	0
2	NMM	G	7	2	8,11,12	0.42	0	7,12,14	0.59	0
2	NMM	F	12	2	8,11,12	0.49	0	7,12,14	1.66	2 (28%)

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
2	NMM	E	7	2	8,11,12	0.63	0	7,12,14	0.74	0
2	NMM	H	12	2	8,11,12	0.51	0	7,12,14	0.51	0
2	NMM	E	12	2	8,11,12	0.51	0	7,12,14	2.20	3 (42%)
2	NMM	H	7	2	8,11,12	0.54	0	7,12,14	0.53	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	NMM	F	7	2	-	0/9/11/13	-
2	NMM	G	7	2	-	1/9/11/13	-
2	NMM	F	12	2	-	3/9/11/13	-
2	NMM	E	7	2	-	1/9/11/13	-
2	NMM	H	12	2	-	3/9/11/13	-
2	NMM	E	12	2	-	2/9/11/13	-
2	NMM	H	7	2	-	0/9/11/13	-

There are no bond length outliers.

All (5) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	E	12	NMM	NE-CZ-NH2	4.66	123.75	119.48
2	F	12	NMM	CD-NE-CZ	3.08	129.11	123.46
2	F	12	NMM	CG-CD-NE	2.55	119.35	112.20
2	E	12	NMM	NE-CZ-NH1	-2.45	115.74	120.26
2	E	12	NMM	CD-NE-CZ	2.28	127.64	123.46

There are no chirality outliers.

5 of 10 torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	E	12	NMM	O-C-CA-CB
2	F	12	NMM	CG-CD-NE-CZ
2	H	12	NMM	O-C-CA-CB
2	E	12	NMM	CA-CB-CG-CD
2	H	12	NMM	NE-CD-CG-CB

There are no ring outliers.

1 monomer is involved in 2 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	F	12	NMM	2	0

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

4 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# $ Z > 2$	Counts	RMSZ	# $ Z > 2$
3	SFG	B	501	-	28,29,29	1.48	4 (14%)	34,42,42	1.86	9 (26%)
3	SFG	A	501	-	28,29,29	1.43	5 (17%)	34,42,42	1.94	10 (29%)
3	SFG	C	501	-	28,29,29	1.37	4 (14%)	34,42,42	2.05	11 (32%)
3	SFG	D	501	-	28,29,29	1.45	6 (21%)	34,42,42	2.00	9 (26%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	SFG	B	501	-	-	4/17/33/33	0/3/3/3
3	SFG	A	501	-	-	2/17/33/33	0/3/3/3
3	SFG	C	501	-	-	3/17/33/33	0/3/3/3
3	SFG	D	501	-	-	2/17/33/33	0/3/3/3

The worst 5 of 19 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	D	501	SFG	C5-C4	4.76	1.47	1.39
3	B	501	SFG	C5-C4	4.54	1.47	1.39
3	A	501	SFG	C5-C4	4.25	1.46	1.39
3	C	501	SFG	C5-C4	4.11	1.46	1.39
3	D	501	SFG	C5-N7	-2.86	1.33	1.39

The worst 5 of 39 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	C	501	SFG	C5-C4-N3	-4.96	119.89	126.72
3	C	501	SFG	N3-C4-N9	4.89	135.49	127.17
3	D	501	SFG	N3-C4-N9	4.85	135.42	127.17
3	D	501	SFG	C5-C4-N3	-4.85	120.04	126.72
3	A	501	SFG	C4-N9-C8	4.59	110.56	105.74

There are no chirality outliers.




5 of 11 torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
3	A	501	SFG	NE-CD-CG-CB
3	A	501	SFG	C5'-CD-CG-CB
3	B	501	SFG	NE-CD-CG-CB
3	B	501	SFG	C5'-CD-CG-CB
3	C	501	SFG	NE-CD-CG-CB

There are no ring outliers.

No monomer is involved in short contacts.

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.





5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	344/349 (98%)	0.14	7 (2%) 65 72	8, 22, 35, 64	10 (2%)
1	B	344/349 (98%)	0.07	3 (0%) 81 85	9, 20, 31, 60	8 (2%)
1	C	343/349 (98%)	0.48	11 (3%) 50 56	15, 28, 39, 61	5 (1%)
1	D	343/349 (98%)	0.64	17 (4%) 34 38	12, 28, 43, 65	5 (1%)
2	E	7/19 (36%)	1.07	2 (28%) 1 1	21, 23, 39, 50	0
2	F	7/19 (36%)	1.44	2 (28%) 1 1	19, 23, 34, 54	0
2	G	5/19 (26%)	0.70	0 100 100	26, 28, 30, 34	0
2	H	6/19 (31%)	1.49	1 (16%) 4 4	26, 32, 37, 58	0
All	All	1399/1472 (95%)	0.35	43 (3%) 51 57	8, 25, 39, 65	28 (2%)

The worst 5 of 43 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
2	H	13	PRO	4.2
2	F	5	ALA	3.6
1	B	477	THR	3.4
1	C	314	TYR	3.4
2	E	4	GLY	3.4

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
2	NMM	E	12	12/13	0.64	0.23	40,59,75,76	0

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
2	NMM	H	12	12/13	0.66	0.20	47,61,71,71	0
2	NMM	F	12	12/13	0.73	0.18	40,59,74,75	0
2	NMM	G	7	12/13	0.92	0.09	20,22,27,27	0
2	NMM	E	7	12/13	0.93	0.08	19,21,23,24	0
2	NMM	H	7	12/13	0.94	0.09	22,25,26,28	0
2	NMM	F	7	12/13	0.96	0.06	17,19,21,22	0

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)




In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q < 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
3	SFG	C	501	27/27	0.93	0.07	22,24,30,30	0
3	SFG	D	501	27/27	0.94	0.07	21,24,31,33	0
3	SFG	A	501	27/27	0.96	0.05	14,15,18,22	0
3	SFG	B	501	27/27	0.96	0.06	13,14,21,23	0




The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.

Electron density around SFG C 501:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)




**Electron density around SFG D 501:**

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around SFG A 501:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)

**Electron density around SFG B 501:**

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



6.5 Other polymers [i](#)

There are no such residues in this entry.