



Full wwPDB X-ray Structure Validation Report ⓘ

Apr 25, 2026 – 11:45 AM EDT

PDB ID : 6DXZ / pdb_00006dxz
Title : Rabbit N-acyl ethanolamine-hydrolyzing acid amidase (NAAA) in complex with non-covalent benzothiazole-piperazine inhibitor ARN19702, in presence of Triton X-100
Authors : Gorelik, A.; Gebai, A.; Illes, K.; Piomelli, D.; Nagar, B.
Deposited on : 2018-07-01
Resolution : 2.70 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Buster-report	:	wwPDB partial adaption of 1.1.7 (2018)
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION


The reported resolution of this entry is 2.70 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	3538 (2.70-2.70)
Clashscore	190562	3843 (2.70-2.70)
Ramachandran outliers	187476	3778 (2.70-2.70)
Sidechain outliers	187428	3778 (2.70-2.70)
RSRZ outliers	180081	3538 (2.70-2.70)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	107	
2	B	232	

2 Entry composition

There are 9 unique types of molecules in this entry. The entry contains 5728 atoms, of which 2799 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called N-acylethanolamine acid amidase alpha-subunit.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
1	A	95	Total	C	H	N	O	S	0	1	0
			1542	499	785	127	129	2			


There are 11 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	21	ASP	-	expression tag	UNP G1T7U7
A	22	ARG	-	expression tag	UNP G1T7U7
A	23	HIS	-	expression tag	UNP G1T7U7
A	24	HIS	-	expression tag	UNP G1T7U7
A	25	HIS	-	expression tag	UNP G1T7U7
A	26	HIS	-	expression tag	UNP G1T7U7
A	27	HIS	-	expression tag	UNP G1T7U7
A	28	HIS	-	expression tag	UNP G1T7U7
A	29	LYS	-	expression tag	UNP G1T7U7
A	30	LEU	-	expression tag	UNP G1T7U7
A	48	LEU	-	insertion	UNP G1T7U7

- Molecule 2 is a protein called N-acylethanolamine acid amidase beta-subunit.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
2	B	230	Total	C	H	N	O	S	0	3	0
			3707	1204	1840	321	339	3			

- Molecule 3 is 2-acetamido-2-deoxy-beta-D-glucopyranose (CCD ID: NAG) (formula: C₈H₁₅NO₆).




Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
3	A	1	Total	C	H	N	O	0	0
			28	8	14	1	5		
3	A	1	Total	C	H	N	O	0	0
			28	8	14	1	5		

- Molecule 4 is POTASSIUM ION (CCD ID: K) (formula: K).


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	1	Total	K	0	0
			1	1		

- Molecule 5 is FRAGMENT OF TRITON X-100 (CCD ID: TRT) (formula: C₂₁H₃₆O₄).




Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
5	A	1	Total	C	H	O	0	0
			61	21	36	4		
5	A	1	Total	C	H	O	0	0
			61	21	36	4		

- Molecule 6 is 4-(2,4,4-trimethylpentan-2-yl)phenol (CCD ID: 27L) (formula: C₁₄H₂₂O).




Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
6	A	1	Total	C	H	O	0	0
			37	14	22	1		

- Molecule 7 is [2-(ethylsulfonyl)phenyl][(2S)-4-(6-fluoro-1,3-benzothiazol-2-yl)-2-methylpiperazin-1-yl]methanone (CCD ID: WTF) (formula: C₂₁H₂₂FN₃O₃S₂).



Mol	Chain	Residues	Atoms						ZeroOcc	AltConf
7	B	1	Total	C	F	H	N	O	S	
			52	21	1	22	3	3	2	

- Molecule 8 is 2-{2-[4-(1,1,3,3-TETRAMETHYLBUTYL)PHENOXY]ETHOXY}ETHANOL (CCD ID: TON) (formula: C₁₈H₃₀O₃).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
8	B	1	Total	C	H	O		
			51	18	30	3	0	0



- Molecule 9 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
9	A	52	Total 52	O 52	0	0
9	B	108	Total 108	O 108	0	0

3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: N-acylethanolamine acid amidase alpha-subunit



- Molecule 2: N-acylethanolamine acid amidase beta-subunit



4 Data and refinement statistics

Property	Value	Source
Space group	P 43 3 2	Depositor
Cell constants a, b, c, α , β , γ	159.14Å 159.14Å 159.14Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	45.94 – 2.70 45.94 – 2.70	Depositor EDS
% Data completeness (in resolution range)	98.2 (45.94-2.70) 95.9 (45.94-2.70)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.85 (at 2.69Å)	Xtriage
Refinement program	PHENIX (1.13_2998)	Depositor
R, R_{free}	0.182 , 0.222 0.183 , 0.223	Depositor DCC
R_{free} test set	1917 reflections (10.00%)	wwPDB-VP
Wilson B-factor (Å ²)	41.6	Xtriage
Anisotropy	0.000	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.39 , 49.0	EDS
L-test for twinning ²	$\langle L \rangle = 0.48$, $\langle L^2 \rangle = 0.31$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.94	EDS
Total number of atoms	5728	wwPDB-VP
Average B, all atoms (Å ²)	50.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 2.18% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: K, NAG, 27L, WTF, TRT, TON

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.20	0/780	0.32	0/1064
2	B	0.14	0/1930	0.30	0/2636
All	All	0.16	0/2710	0.31	0/3700

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	757	785	785	4	0
2	B	1867	1840	1840	7	0
3	A	28	28	26	0	0
4	A	1	0	0	0	0
5	A	50	72	72	3	0
6	A	15	22	22	5	0
7	B	30	22	0	0	0
8	B	21	30	30	0	0
9	A	52	0	0	0	0
9	B	108	0	0	1	0
All	All	2929	2799	2775	17	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 3.

All (17) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
6:A:206:27L:H13	6:A:206:27L:H9	1.83	0.61
6:A:206:27L:CAD	6:A:206:27L:H1	2.35	0.57
2:B:128:CYS:N	9:B:501:HOH:O	2.38	0.57
2:B:319:SER:O	2:B:323:LEU:HB2	2.07	0.53
2:B:288:THR:OG1	2:B:289:ASN:N	2.42	0.46
5:A:204:TRT:H3C2	5:A:204:TRT:C9	2.46	0.45
1:A:95:PRO:N	1:A:96:PRO:HD2	2.31	0.45
1:A:122:TYR:O	2:B:200:ARG:HD2	2.16	0.44
1:A:123:GLU:OE1	1:A:123:GLU:N	2.49	0.44
6:A:206:27L:H1	6:A:206:27L:H14	2.01	0.43
2:B:129:THR:O	2:B:144:ARG:HA	2.19	0.42
6:A:206:27L:H17	6:A:206:27L:CAC	2.49	0.42
6:A:206:27L:H13	6:A:206:27L:H1	2.00	0.42
5:A:204:TRT:H2C3	5:A:204:TRT:H8C3	2.03	0.41
1:A:65:ALA:HB1	2:B:153:ILE:HG21	2.04	0.40
5:A:205:TRT:C10	5:A:205:TRT:H3C2	2.51	0.40
2:B:155[A]:ARG:NE	2:B:356:ARG:O	2.54	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	94/107 (88%)	92 (98%)	2 (2%)	0	100	100
2	B	231/232 (100%)	219 (95%)	12 (5%)	0	100	100
All	All	325/339 (96%)	311 (96%)	14 (4%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	81/92 (88%)	80 (99%)	1 (1%)	63	84
2	B	200/199 (100%)	198 (99%)	2 (1%)	68	86
All	All	281/291 (97%)	278 (99%)	3 (1%)	65	85

All (3) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	93	LEU
2	B	144	ARG
2	B	323	LEU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (1) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	56	HIS

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry

Of 8 ligands modelled in this entry, 1 is monoatomic - leaving 7 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
6	27L	A	206	-	15,15,15	0.52	0	23,23,23	0.64	0
3	NAG	A	201	1	14,14,15	0.28	0	17,19,21	0.56	0
8	TON	B	402	-	21,21,21	0.92	2 (9%)	29,29,29	0.64	0
7	WTF	B	401	-	32,33,33	3.14	17 (53%)	47,49,49	3.27	15 (31%)
3	NAG	A	202	1	14,14,15	0.27	0	17,19,21	0.37	0
5	TRT	A	204	-	25,25,25	1.12	2 (8%)	33,33,33	0.56	0
5	TRT	A	205	-	25,25,25	1.08	2 (8%)	33,33,33	0.58	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
6	27L	A	206	-	-	2/12/12/12	0/1/1/1
3	NAG	A	201	1	-	3/6/23/26	0/1/1/1
8	TON	B	402	-	-	2/19/19/19	0/1/1/1
7	WTF	B	401	-	-	0/21/34/34	0/4/4/4
3	NAG	A	202	1	-	2/6/23/26	0/1/1/1
5	TRT	A	204	-	-	5/23/23/23	0/1/1/1
5	TRT	A	205	-	-	9/23/23/23	0/1/1/1

All (23) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
7	B	401	WTF	C12-N14	6.83	1.50	1.34
7	B	401	WTF	C07-C12	6.20	1.59	1.50
7	B	401	WTF	C21-N17	5.43	1.46	1.34
7	B	401	WTF	C18-N17	5.42	1.53	1.46

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
7	B	401	WTF	C21-N22	5.23	1.43	1.30
7	B	401	WTF	C30-C23	4.70	1.47	1.40
7	B	401	WTF	C23-N22	4.17	1.48	1.39
7	B	401	WTF	C06-S03	3.57	1.86	1.78
7	B	401	WTF	C29-C27	3.35	1.43	1.37
7	B	401	WTF	C19-N14	-3.34	1.44	1.47
7	B	401	WTF	C26-C24	3.29	1.44	1.39
7	B	401	WTF	C16-N17	3.28	1.52	1.47
5	A	204	TRT	O15-C12	3.15	1.44	1.37
7	B	401	WTF	C26-C27	3.11	1.42	1.37
5	A	205	TRT	O15-C12	3.01	1.44	1.37
5	A	204	TRT	C6-C9	2.81	1.58	1.53
8	B	402	TON	O1-C15	2.75	1.43	1.37
5	A	205	TRT	C6-C9	2.74	1.58	1.53
7	B	401	WTF	C15-N14	-2.69	1.42	1.47
7	B	401	WTF	C21-S25	-2.33	1.70	1.75
7	B	401	WTF	C24-S25	-2.31	1.69	1.74
8	B	402	TON	C5-C18	2.08	1.57	1.53
7	B	401	WTF	O13-C12	-2.05	1.18	1.22

All (15) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
7	B	401	WTF	O05-S03-O04	-15.18	102.18	118.45
7	B	401	WTF	C24-S25-C21	6.37	98.14	89.37
7	B	401	WTF	C16-C15-N14	6.12	121.39	110.33
7	B	401	WTF	C19-C18-N17	5.39	117.14	110.99
7	B	401	WTF	C02-S03-C06	5.35	114.17	105.55
7	B	401	WTF	S25-C21-N22	-5.17	109.17	116.92
7	B	401	WTF	C15-C16-N17	4.78	119.92	110.42
7	B	401	WTF	C07-C12-N14	3.21	122.24	117.85
7	B	401	WTF	C26-C24-C23	-3.17	119.63	121.99
7	B	401	WTF	S25-C21-N17	3.13	123.46	119.32
7	B	401	WTF	C18-C19-N14	2.88	115.22	109.34
7	B	401	WTF	C26-C24-S25	2.65	132.68	126.07
7	B	401	WTF	C24-C26-C27	2.60	120.30	116.75
7	B	401	WTF	C29-C27-C26	-2.12	120.43	123.23
7	B	401	WTF	O05-S03-C02	2.03	112.24	108.31

There are no chirality outliers.

All (23) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
5	A	205	TRT	O15-C16-C17-O18
5	A	205	TRT	C4-C1-C5-C6
3	A	202	NAG	C4-C5-C6-O6
5	A	204	TRT	O21-C22-C23-O24
5	A	205	TRT	C2-C1-C5-C6
8	B	402	TON	O2-C3-C4-O3
3	A	202	NAG	O5-C5-C6-O6
5	A	205	TRT	C11-C12-O15-C16
5	A	205	TRT	C13-C12-O15-C16
5	A	205	TRT	C3-C1-C5-C6
5	A	204	TRT	O18-C19-C20-O21
5	A	204	TRT	O15-C16-C17-O18
3	A	201	NAG	C4-C5-C6-O6
5	A	205	TRT	C17-C16-O15-C12
6	A	206	27L	CAN-CAK-CAO-CAD
3	A	201	NAG	O5-C5-C6-O6
6	A	206	27L	CAN-CAK-CAO-CAM
5	A	205	TRT	C16-C17-O18-C19
5	A	205	TRT	C22-C23-O24-C25
8	B	402	TON	C4-C3-O2-C2
3	A	201	NAG	C1-C2-N2-C7
5	A	204	TRT	C22-C23-O24-C25
5	A	204	TRT	C20-C19-O18-C17


There are no ring outliers.


3 monomers are involved in 8 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
6	A	206	27L	5	0
5	A	204	TRT	2	0
5	A	205	TRT	1	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient

equivalents in the CSD to analyse the geometry.





5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2		OWAB(Å ²)	Q<0.9
1	A	95/107 (88%)	-0.10	1 (1%)	78 76	25, 45, 88, 112	1 (1%)
2	B	230/232 (99%)	-0.24	4 (1%)	69 67	21, 43, 80, 111	3 (1%)
All	All	325/339 (95%)	-0.20	5 (1%)	72 70	21, 44, 82, 112	4 (1%)

All (5) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	33	PRO	6.1
2	B	201	GLY	3.7
2	B	357	ASN	3.1
2	B	298	GLU	2.5
2	B	233[A]	GLU	2.1

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)


There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.




Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
3	NAG	A	202	14/15	0.61	0.27	104,145,174,180	0
3	NAG	A	201	14/15	0.72	0.20	70,92,115,121	0
6	27L	A	206	15/15	0.76	0.30	66,92,115,126	0
4	K	A	203	1/1	0.89	0.19	82,82,82,82	0
5	TRT	A	205	25/25	0.90	0.18	30,51,89,99	0
8	TON	B	402	21/21	0.90	0.16	55,79,118,123	0
5	TRT	A	204	25/25	0.92	0.14	36,58,100,133	0
7	WTF	B	401	30/30	0.96	0.10	23,36,44,56	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.






Electron density around TON B 402:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)




**Electron density around TRT A 204:**

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



Electron density around WTF B 401:

$2mF_o - DF_c$ (at 0.7 rmsd) in gray
 $mF_o - DF_c$ (at 3 rmsd) in purple (negative)
and green (positive)



6.5 Other polymers [i](#)

There are no such residues in this entry.