



wwPDB X-ray Structure Validation Summary Report ⓘ

Mar 5, 2026 – 03:46 PM UTC

PDB ID : 8DX4 / pdb_00008dx4
Title : Clostridioides difficile R20291 minor pilin - PilW fused with Maltose Binding Protein
Authors : Ronish, L.A.; Piepenbrink, K.
Deposited on : 2022-08-02
Resolution : 2.49 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0
Mogul : 2022.3.0, CSD as543be (2022)
Xtriage (Phenix) : 2.0
EDS : 3.0
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4 : 9.0.010 (Gargrove)
Density-Fitness : 1.0.12
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.49 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	7589 (2.50-2.46)
Clashscore	190562	8295 (2.50-2.46)
Ramachandran outliers	187476	8164 (2.50-2.46)
Sidechain outliers	187428	8166 (2.50-2.46)
RSRZ outliers	180081	7593 (2.50-2.46)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	512	
1	B	512	
1	C	512	
1	D	512	

2 Entry composition

There are 6 unique types of molecules in this entry. The entry contains 15965 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Maltodextrin-binding protein, Putative pilin protein.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	497	Total	C	N	O	S	5	3	0
			3778	2427	610	731	10			
1	B	497	Total	C	N	O	S	0	5	0
			3818	2453	617	738	10			
1	C	497	Total	C	N	O	S	0	4	0
			3845	2471	622	743	9			
1	D	493	Total	C	N	O	S	0	9	0
			3795	2440	610	734	11			

There are 68 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	0	MET	-	initiating methionine	UNP A0A4P1LXE0
A	82	ALA	ASP	conflict	UNP A0A4P1LXE0
A	83	ALA	LYS	conflict	UNP A0A4P1LXE0
A	172	ALA	GLU	conflict	UNP A0A4P1LXE0
A	173	ALA	ASN	conflict	UNP A0A4P1LXE0
A	239	ALA	LYS	conflict	UNP A0A4P1LXE0
A	359	ALA	GLU	conflict	UNP A0A4P1LXE0
A	362	ALA	LYS	conflict	UNP A0A4P1LXE0
A	363	ALA	ASP	conflict	UNP A0A4P1LXE0
A	1159	LEU	-	expression tag	UNP A0A0H3N4W9
A	1160	GLU	-	expression tag	UNP A0A0H3N4W9
A	1161	HIS	-	expression tag	UNP A0A0H3N4W9
A	1162	HIS	-	expression tag	UNP A0A0H3N4W9
A	1163	HIS	-	expression tag	UNP A0A0H3N4W9
A	1164	HIS	-	expression tag	UNP A0A0H3N4W9
A	1165	HIS	-	expression tag	UNP A0A0H3N4W9
A	1166	HIS	-	expression tag	UNP A0A0H3N4W9
B	0	MET	-	initiating methionine	UNP A0A4P1LXE0
B	82	ALA	ASP	conflict	UNP A0A4P1LXE0
B	83	ALA	LYS	conflict	UNP A0A4P1LXE0
B	172	ALA	GLU	conflict	UNP A0A4P1LXE0

Continued on next page...

Continued from previous page...


Chain	Residue	Modelled	Actual	Comment	Reference
B	173	ALA	ASN	conflict	UNP A0A4P1LXE0
B	239	ALA	LYS	conflict	UNP A0A4P1LXE0
B	359	ALA	GLU	conflict	UNP A0A4P1LXE0
B	362	ALA	LYS	conflict	UNP A0A4P1LXE0
B	363	ALA	ASP	conflict	UNP A0A4P1LXE0
B	1159	LEU	-	expression tag	UNP A0A0H3N4W9
B	1160	GLU	-	expression tag	UNP A0A0H3N4W9
B	1161	HIS	-	expression tag	UNP A0A0H3N4W9
B	1162	HIS	-	expression tag	UNP A0A0H3N4W9
B	1163	HIS	-	expression tag	UNP A0A0H3N4W9
B	1164	HIS	-	expression tag	UNP A0A0H3N4W9
B	1165	HIS	-	expression tag	UNP A0A0H3N4W9
B	1166	HIS	-	expression tag	UNP A0A0H3N4W9
C	0	MET	-	initiating methionine	UNP A0A4P1LXE0
C	82	ALA	ASP	conflict	UNP A0A4P1LXE0
C	83	ALA	LYS	conflict	UNP A0A4P1LXE0
C	172	ALA	GLU	conflict	UNP A0A4P1LXE0
C	173	ALA	ASN	conflict	UNP A0A4P1LXE0
C	239	ALA	LYS	conflict	UNP A0A4P1LXE0
C	359	ALA	GLU	conflict	UNP A0A4P1LXE0
C	362	ALA	LYS	conflict	UNP A0A4P1LXE0
C	363	ALA	ASP	conflict	UNP A0A4P1LXE0
C	1159	LEU	-	expression tag	UNP A0A0H3N4W9
C	1160	GLU	-	expression tag	UNP A0A0H3N4W9
C	1161	HIS	-	expression tag	UNP A0A0H3N4W9
C	1162	HIS	-	expression tag	UNP A0A0H3N4W9
C	1163	HIS	-	expression tag	UNP A0A0H3N4W9
C	1164	HIS	-	expression tag	UNP A0A0H3N4W9
C	1165	HIS	-	expression tag	UNP A0A0H3N4W9
C	1166	HIS	-	expression tag	UNP A0A0H3N4W9
D	0	MET	-	initiating methionine	UNP A0A4P1LXE0
D	82	ALA	ASP	conflict	UNP A0A4P1LXE0
D	83	ALA	LYS	conflict	UNP A0A4P1LXE0
D	172	ALA	GLU	conflict	UNP A0A4P1LXE0
D	173	ALA	ASN	conflict	UNP A0A4P1LXE0
D	239	ALA	LYS	conflict	UNP A0A4P1LXE0
D	359	ALA	GLU	conflict	UNP A0A4P1LXE0
D	362	ALA	LYS	conflict	UNP A0A4P1LXE0
D	363	ALA	ASP	conflict	UNP A0A4P1LXE0
D	1159	LEU	-	expression tag	UNP A0A0H3N4W9
D	1160	GLU	-	expression tag	UNP A0A0H3N4W9
D	1161	HIS	-	expression tag	UNP A0A0H3N4W9

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
D	1162	HIS	-	expression tag	UNP A0A0H3N4W9
D	1163	HIS	-	expression tag	UNP A0A0H3N4W9
D	1164	HIS	-	expression tag	UNP A0A0H3N4W9
D	1165	HIS	-	expression tag	UNP A0A0H3N4W9
D	1166	HIS	-	expression tag	UNP A0A0H3N4W9

- Molecule 2 is 1,2-ETHANEDIOL (CCD ID: EDO) (formula: C₂H₆O₂).




Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
2	A	1	Total	C	O	0	0
			4	2	2		
2	A	1	Total	C	O	0	0
			4	2	2		
2	A	1	Total	C	O	0	0
			4	2	2		
2	A	1	Total	C	O	0	0
			4	2	2		
2	B	1	Total	C	O	0	0
			4	2	2		
2	B	1	Total	C	O	0	0
			4	2	2		
2	B	1	Total	C	O	0	0
			4	2	2		
2	C	1	Total	C	O	0	0
			4	2	2		

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
2	C	1	Total	C	O	0	0
			4	2	2		
2	C	1	Total	C	O	0	0
			4	2	2		
2	D	1	Total	C	O	0	0
			4	2	2		
2	D	1	Total	C	O	0	0
			4	2	2		
2	D	1	Total	C	O	0	0
			4	2	2		
2	D	1	Total	C	O	0	0
			4	2	2		
2	D	1	Total	C	O	0	0
			4	2	2		

- Molecule 3 is 4-(2-HYDROXYETHYL)-1-PIPERAZINE ETHANESULFONIC ACID (CCD ID: EPE) (formula: $C_8H_{18}N_2O_4S$).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
3	C	1	Total	C	N	O	S	0	0
			15	8	2	4	1		

- Molecule 4 is TRIETHYLENE GLYCOL (CCD ID: PGE) (formula: $C_6H_{14}O_4$).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
4	D	1	Total	C	O	0	0
			10	6	4		

- Molecule 5 is DI(HYDROXYETHYL)ETHER (CCD ID: PEG) (formula: $C_4H_{10}O_3$).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
5	D	1	Total	C	O	0	0
			7	4	3		

- Molecule 6 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
6	A	88	Total 88	O 88	0	0
6	B	201	Total 201	O 201	0	0
6	C	165	Total 165	O 165	0	0
6	D	171	Total 171	O 171	0	0



4 Data and refinement statistics

Property	Value	Source
Space group	P 1	Depositor
Cell constants a, b, c, α , β , γ	65.33Å 81.80Å 102.96Å 92.44° 90.95° 113.37°	Depositor
Resolution (Å)	39.73 – 2.49 39.73 – 2.49	Depositor EDS
% Data completeness (in resolution range)	93.5 (39.73-2.49) 91.1 (39.73-2.49)	Depositor EDS
R_{merge}	0.08	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.44 (at 2.48Å)	Xtriage
Refinement program	PHENIX 1.12_2829	Depositor
R, R_{free}	0.186 , 0.218 0.192 , 0.192	Depositor DCC
R_{free} test set	3179 reflections (4.63%)	wwPDB-VP
Wilson B-factor (Å ²)	39.0	Xtriage
Anisotropy	0.637	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.34 , 46.2	EDS
L-test for twinning ²	$\langle L \rangle = 0.48$, $\langle L^2 \rangle = 0.31$	Xtriage
Estimated twinning fraction	0.207 for h,-h-k,-l	Xtriage
F_o, F_c correlation	0.96	EDS
Total number of atoms	15965	wwPDB-VP
Average B, all atoms (Å ²)	61.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.99% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: EDO, EPE, PGE, PEG

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.19	0/3871	0.45	1/5263 (0.0%)
1	B	0.18	0/3917	0.40	0/5316
1	C	0.19	0/3938	0.46	0/5339
1	D	0.21	0/3895	0.45	0/5287
All	All	0.19	0/15621	0.44	1/21205 (0.0%)

There are no bond length outliers.

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	29	LYS	CD-CE-NZ	-7.63	87.49	111.90

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3778	0	3708	34	0
1	B	3818	0	3785	34	0
1	C	3845	0	3839	31	1
1	D	3795	0	3749	37	1
2	A	16	0	24	1	0
2	B	16	0	24	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
2	C	12	0	18	3	0
2	D	28	0	42	0	0
3	C	15	0	17	2	0
4	D	10	0	14	1	0
5	D	7	0	10	0	0
6	A	88	0	0	0	0
6	B	201	0	0	2	0
6	C	165	0	0	3	0
6	D	171	0	0	0	0
All	All	15965	0	15230	130	1

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 4.

The worst 5 of 130 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:1134:ASN:ND2	1:B:367:ASN:OD1	2.10	0.85
1:D:1094:TYR:CZ	1:D:1120:ILE:HD11	2.15	0.80
1:D:1094:TYR:CE1	1:D:1120:ILE:HD11	2.16	0.79
1:C:365:GLN:HG3	2:C:1202:EDO:H12	1.66	0.77
1:B:1078:PHE:HB3	1:B:1082:PRO:HD2	1.66	0.76

All (1) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:C:337:SER:OG	1:D:41[B]:ASP:OD2[1_445]	2.17	0.03

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	498/512 (97%)	466 (94%)	25 (5%)	7 (1%)	9	15
1	B	500/512 (98%)	476 (95%)	24 (5%)	0	100	100
1	C	499/512 (98%)	480 (96%)	18 (4%)	1 (0%)	43	61
1	D	498/512 (97%)	478 (96%)	16 (3%)	4 (1%)	16	28
All	All	1995/2048 (97%)	1900 (95%)	83 (4%)	12 (1%)	21	35

5 of 12 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	1082	PRO
1	A	1087	ILE
1	A	1086	LYS
1	C	1108	LYS
1	D	1110	LEU

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	383/410 (93%)	376 (98%)	7 (2%)	51	74
1	B	392/410 (96%)	385 (98%)	7 (2%)	51	74
1	C	398/410 (97%)	391 (98%)	7 (2%)	51	74
1	D	388/410 (95%)	382 (98%)	6 (2%)	57	78
All	All	1561/1640 (95%)	1534 (98%)	27 (2%)	51	75

5 of 27 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	C	1049	SER
1	C	1088	GLU
1	D	1078	PHE
1	C	1086	LYS
1	C	1113	SER

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 11 such sidechains are listed below:

Mol	Chain	Res	Type
1	D	1095	ASN
1	D	1140	ASN
1	D	1146	ASN
1	D	1142	GLN
1	D	39	HIS

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

21 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
2	EDO	B	1203	-	3,3,3	0.42	0	2,2,2	0.38	0
2	EDO	D	1203	-	3,3,3	0.42	0	2,2,2	0.47	0
2	EDO	D	1208	-	3,3,3	0.43	0	2,2,2	0.37	0
2	EDO	D	1205	-	3,3,3	0.42	0	2,2,2	0.37	0
2	EDO	A	1203	-	3,3,3	0.44	0	2,2,2	0.32	0
4	PGE	D	1202	-	9,9,9	0.31	0	8,8,8	0.38	0
3	EPE	C	1203	-	15,15,15	0.91	1 (6%)	19,20,20	1.63	4 (21%)

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
2	EDO	B	1201	-	3,3,3	0.43	0	2,2,2	0.42	0
2	EDO	B	1204	-	3,3,3	0.43	0	2,2,2	0.37	0
2	EDO	D	1207	-	3,3,3	0.45	0	2,2,2	0.40	0
2	EDO	C	1204	-	3,3,3	0.74	0	2,2,2	0.45	0
2	EDO	A	1202	-	3,3,3	0.42	0	2,2,2	0.39	0
5	PEG	D	1206	-	6,6,6	0.51	0	5,5,5	0.27	0
2	EDO	D	1209	-	3,3,3	0.52	0	2,2,2	1.12	0
2	EDO	C	1201	-	3,3,3	0.42	0	2,2,2	0.37	0
2	EDO	B	1202	-	3,3,3	0.44	0	2,2,2	0.31	0
2	EDO	A	1201	-	3,3,3	0.41	0	2,2,2	0.51	0
2	EDO	D	1204	-	3,3,3	0.41	0	2,2,2	0.31	0
2	EDO	C	1202	-	3,3,3	0.47	0	2,2,2	0.40	0
2	EDO	D	1201	-	3,3,3	0.41	0	2,2,2	0.23	0
2	EDO	A	1204	-	3,3,3	0.41	0	2,2,2	0.41	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	EDO	B	1203	-	-	0/1/1/1	-
2	EDO	D	1203	-	-	1/1/1/1	-
2	EDO	D	1208	-	-	0/1/1/1	-
2	EDO	D	1205	-	-	0/1/1/1	-
2	EDO	A	1203	-	-	0/1/1/1	-
4	PGE	D	1202	-	-	3/7/7/7	-
3	EPE	C	1203	-	-	2/9/19/19	0/1/1/1
2	EDO	B	1201	-	-	1/1/1/1	-
2	EDO	B	1204	-	-	1/1/1/1	-
2	EDO	D	1207	-	-	1/1/1/1	-
2	EDO	C	1204	-	-	1/1/1/1	-
2	EDO	A	1202	-	-	0/1/1/1	-
5	PEG	D	1206	-	-	2/4/4/4	-
2	EDO	D	1209	-	-	0/1/1/1	-
2	EDO	C	1201	-	-	0/1/1/1	-
2	EDO	B	1202	-	-	0/1/1/1	-
2	EDO	A	1201	-	-	1/1/1/1	-
2	EDO	D	1204	-	-	1/1/1/1	-
2	EDO	C	1202	-	-	0/1/1/1	-
2	EDO	D	1201	-	-	1/1/1/1	-
2	EDO	A	1204	-	-	1/1/1/1	-

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	C	1203	EPE	C10-S	3.04	1.81	1.77

All (4) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	C	1203	EPE	C5-N4-C3	3.51	116.40	108.84
3	C	1203	EPE	C7-N4-C3	3.09	119.47	111.24
3	C	1203	EPE	C7-N4-C5	2.91	119.00	111.24
3	C	1203	EPE	O2S-S-C10	2.33	110.24	106.73

There are no chirality outliers.

5 of 16 torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
3	C	1203	EPE	C8-C7-N4-C3
4	D	1202	PGE	O2-C3-C4-O3
5	D	1206	PEG	O2-C3-C4-O4
2	A	1201	EDO	O1-C1-C2-O2
2	C	1204	EDO	O1-C1-C2-O2

There are no ring outliers.

5 monomers are involved in 7 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
4	D	1202	PGE	1	0
3	C	1203	EPE	2	0
2	C	1201	EDO	1	0
2	A	1201	EDO	1	0
2	C	1202	EDO	2	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	497/512 (97%)	-1.15	0 100 100	26, 67, 127, 163	3 (0%)
1	B	497/512 (97%)	-1.29	0 100 100	16, 48, 126, 155	5 (1%)
1	C	497/512 (97%)	-1.35	0 100 100	18, 48, 116, 142	4 (0%)
1	D	493/512 (96%)	-1.36	0 100 100	19, 47, 123, 174	9 (1%)
All	All	1984/2048 (96%)	-1.29	0 100 100	16, 53, 123, 174	21 (1%)

There are no RSRZ outliers to report.

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
2	EDO	D	1209	4/4	0.95	0.06	67,68,73,88	0
2	EDO	A	1204	4/4	0.96	0.06	71,71,72,73	0
2	EDO	D	1207	4/4	0.97	0.04	66,67,67,68	0
2	EDO	D	1203	4/4	0.97	0.06	88,89,89,89	0

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
2	EDO	A	1202	4/4	0.98	0.08	65,65,67,68	0
2	EDO	D	1205	4/4	0.98	0.06	68,70,72,73	0
2	EDO	B	1204	4/4	0.98	0.05	81,81,81,82	0
2	EDO	D	1208	4/4	0.98	0.04	72,72,73,75	0
2	EDO	C	1202	4/4	0.98	0.06	54,55,55,62	0
2	EDO	A	1201	4/4	0.99	0.06	54,55,55,57	0
2	EDO	C	1204	4/4	0.99	0.06	72,72,80,83	0
2	EDO	D	1201	4/4	0.99	0.06	50,54,57,61	0
2	EDO	B	1201	4/4	0.99	0.05	65,67,71,75	0
2	EDO	D	1204	4/4	0.99	0.05	62,63,63,64	0
2	EDO	B	1202	4/4	0.99	0.05	60,62,65,69	0
2	EDO	B	1203	4/4	0.99	0.03	45,50,50,51	0
2	EDO	A	1203	4/4	0.99	0.04	75,75,76,78	0
2	EDO	C	1201	4/4	0.99	0.05	53,54,56,57	0
3	EPE	C	1203	15/15	0.99	0.05	86,89,108,108	0
4	PGE	D	1202	10/10	0.99	0.05	59,62,68,68	0
5	PEG	D	1206	7/7	0.99	0.04	74,80,82,83	0

6.5 Other polymers [i](#)

There are no such residues in this entry.