



wwPDB X-ray Structure Validation Summary Report ⓘ

Mar 8, 2026 – 11:40 AM UTC

PDB ID : 2DYY / pdb_00002dyd
Title : Crystal structure of putative translation initiation inhibitor PH0854 from *Pyrococcus horikoshii*
Authors : Ihsanawati; Kishishita, S.; Murayama, K.; Chen, L.; Liu, Z.J.; Wang, B.C.; Shirouzu, M.; Bessho, Y.; Yokoyama, S.; RIKEN Structural Genomics/Proteomics Initiative (RSGI)
Deposited on : 2006-09-19
Resolution : 2.60 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.60 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	4008 (2.60-2.60)
Clashscore	190562	4347 (2.60-2.60)
Ramachandran outliers	187476	4277 (2.60-2.60)
Sidechain outliers	187428	4277 (2.60-2.60)
RSRZ outliers	180081	4008 (2.60-2.60)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	126	 2% 60% 37% .
1	B	126	 62% 29% • 6%
1	C	126	 65% 29% • 6%
1	D	126	 74% 25% •
1	E	126	 67% 26% • 5%

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
1	F	126	2% 65% 24% 5% 6%
1	G	126	75% 20% 5%
1	H	126	6% 60% 33%
1	I	126	% 51% 37% 8%
1	J	126	6% 56% 38%
1	K	126	2% 63% 29% 5%
1	L	126	% 63% 31% 6%

2 Entry composition [i](#)

There are 2 unique types of molecules in this entry. The entry contains 11461 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called UPF0076 protein PH0854.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	125	Total	C	N	O	S	0	0	0
			965	623	155	185	2			
1	B	119	Total	C	N	O	S	0	0	0
			928	601	149	177	1			
1	C	119	Total	C	N	O	S	0	0	0
			921	596	147	177	1			
1	D	126	Total	C	N	O	S	0	0	0
			970	626	156	186	2			
1	E	120	Total	C	N	O	S	0	0	0
			922	598	146	177	1			
1	F	118	Total	C	N	O		0	0	0
			905	586	145	174				
1	G	125	Total	C	N	O	S	0	0	0
			971	628	155	187	1			
1	H	122	Total	C	N	O	S	0	0	0
			930	599	147	182	2			
1	I	116	Total	C	N	O		0	0	0
			895	580	142	173				
1	J	124	Total	C	N	O	S	0	0	0
			957	619	154	182	2			
1	K	123	Total	C	N	O	S	0	0	0
			944	609	151	183	1			
1	L	119	Total	C	N	O	S	0	0	0
			920	595	147	177	1			

- Molecule 2 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	A	19	Total	O	0	0
			19	19		
2	B	27	Total	O	0	0
			27	27		

Continued on next page...


Continued from previous page...

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	C	22	Total 22	O 22	0	0
2	D	22	Total 22	O 22	0	0
2	E	21	Total 21	O 21	0	0
2	F	14	Total 14	O 14	0	0
2	G	29	Total 29	O 29	0	0
2	H	19	Total 19	O 19	0	0
2	I	10	Total 10	O 10	0	0
2	J	19	Total 19	O 19	0	0
2	K	24	Total 24	O 24	0	0
2	L	7	Total 7	O 7	0	0


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: UPF0076 protein PH0854




- Molecule 1: UPF0076 protein PH0854



- Molecule 1: UPF0076 protein PH0854




- Molecule 1: UPF0076 protein PH0854




- Molecule 1: UPF0076 protein PH0854

Chain E: 



- Molecule 1: UPF0076 protein PH0854

Chain F: 




- Molecule 1: UPF0076 protein PH0854

Chain G: 




- Molecule 1: UPF0076 protein PH0854

Chain H: 




- Molecule 1: UPF0076 protein PH0854

Chain I: 




- Molecule 1: UPF0076 protein PH0854

Chain J:  6% 56% 38% . .



- Molecule 1: UPF0076 protein PH0854

Chain K:  2% 63% 29% 5% . .



- Molecule 1: UPF0076 protein PH0854

Chain L:  % 63% 31% . 6%



4 Data and refinement statistics

Property	Value	Source
Space group	P 1 2 1	Depositor
Cell constants a, b, c, α , β , γ	92.49Å 43.09Å 170.21Å 90.00° 91.51° 90.00°	Depositor
Resolution (Å)	29.73 – 2.60 29.73 – 2.60	Depositor EDS
% Data completeness (in resolution range)	96.7 (29.73-2.60) 96.7 (29.73-2.60)	Depositor EDS
R_{merge}	0.10	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	6.47 (at 2.61Å)	Xtriage
Refinement program	CNS 1.1	Depositor
R, R_{free}	0.209 , 0.295 0.209 , 0.294	Depositor DCC
R_{free} test set	2048 reflections (5.01%)	wwPDB-VP
Wilson B-factor (Å ²)	40.0	Xtriage
Anisotropy	0.587	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.36 , 59.3	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	0.017 for h,-k,-l	Xtriage
F_o, F_c correlation	0.94	EDS
Total number of atoms	11461	wwPDB-VP
Average B, all atoms (Å ²)	42.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The analyses of the Patterson function reveals a significant off-origin peak that is 34.61 % of the origin peak, indicating pseudo-translational symmetry. The chance of finding a peak of this or larger height randomly in a structure without pseudo-translational symmetry is equal to 6.6413e-04. The detected translational NCS is most likely also responsible for the elevated intensity ratio.*

¹ Intensities estimated from amplitudes.

² Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.48	0/980	0.84	1/1324 (0.1%)
1	B	0.52	0/943	0.93	0/1274
1	C	0.50	0/936	0.95	2/1267 (0.2%)
1	D	0.49	0/986	0.91	1/1334 (0.1%)
1	E	0.47	0/936	0.86	0/1267
1	F	0.45	0/920	0.88	2/1248 (0.2%)
1	G	0.52	0/988	0.93	1/1337 (0.1%)
1	H	0.46	0/944	0.88	4/1280 (0.3%)
1	I	0.40	0/910	0.81	0/1234
1	J	0.46	0/972	0.88	1/1313 (0.1%)
1	K	0.45	0/959	0.92	4/1299 (0.3%)
1	L	0.44	0/935	0.85	2/1266 (0.2%)
All	All	0.47	0/11409	0.89	18/15443 (0.1%)

There are no bond length outliers.

The worst 5 of 18 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	J	43	VAL	N-CA-C	9.21	116.06	106.21
1	G	117	ILE	N-CA-C	6.96	117.48	107.88
1	C	103	ARG	N-CA-C	6.67	120.27	109.40
1	K	100	LYS	CA-C-N	5.81	125.79	120.21
1	K	100	LYS	C-N-CA	5.81	125.79	120.21

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen

atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	965	0	983	47	0
1	B	928	0	957	42	0
1	C	921	0	937	28	0
1	D	970	0	986	32	0
1	E	922	0	939	30	0
1	F	905	0	910	29	0
1	G	971	0	983	32	0
1	H	930	0	921	40	0
1	I	895	0	904	53	0
1	J	957	0	979	46	0
1	K	944	0	947	49	0
1	L	920	0	935	41	0
2	A	19	0	0	3	0
2	B	27	0	0	0	0
2	C	22	0	0	2	0
2	D	22	0	0	0	0
2	E	21	0	0	3	0
2	F	14	0	0	1	0
2	G	29	0	0	2	0
2	H	19	0	0	0	0
2	I	10	0	0	0	0
2	J	19	0	0	0	0
2	K	24	0	0	1	0
2	L	7	0	0	1	0
All	All	11461	0	11381	421	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 19.

The worst 5 of 421 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:C:89:ASN:HD22	1:C:91:VAL:HG23	1.22	1.02
1:C:12:LYS:HD2	1:C:13:PRO:HD2	1.45	0.97
1:J:106:VAL:HG23	1:K:111:LEU:HD23	1.51	0.92
1:H:76:THR:HB	1:H:120:GLU:HG2	1.50	0.90
1:A:36:ASP:HB3	1:A:39:THR:HG22	1.56	0.87

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	121/126 (96%)	110 (91%)	10 (8%)	1 (1%)	16	34
1	B	115/126 (91%)	113 (98%)	2 (2%)	0	100	100
1	C	115/126 (91%)	111 (96%)	4 (4%)	0	100	100
1	D	124/126 (98%)	115 (93%)	9 (7%)	0	100	100
1	E	114/126 (90%)	110 (96%)	4 (4%)	0	100	100
1	F	114/126 (90%)	101 (89%)	10 (9%)	3 (3%)	4	7
1	G	123/126 (98%)	115 (94%)	7 (6%)	1 (1%)	16	34
1	H	116/126 (92%)	101 (87%)	11 (10%)	4 (3%)	3	4
1	I	112/126 (89%)	98 (88%)	12 (11%)	2 (2%)	6	14
1	J	120/126 (95%)	103 (86%)	14 (12%)	3 (2%)	4	8
1	K	119/126 (94%)	105 (88%)	11 (9%)	3 (2%)	4	8
1	L	115/126 (91%)	104 (90%)	11 (10%)	0	100	100
All	All	1408/1512 (93%)	1286 (91%)	105 (8%)	17 (1%)	10	23

5 of 17 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	44	LYS
1	F	90	GLU
1	H	115	VAL
1	J	39	THR
1	F	43	VAL

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	101/106 (95%)	96 (95%)	5 (5%)	22	46
1	B	99/106 (93%)	94 (95%)	5 (5%)	21	45
1	C	97/106 (92%)	96 (99%)	1 (1%)	68	86
1	D	101/106 (95%)	99 (98%)	2 (2%)	48	74
1	E	97/106 (92%)	89 (92%)	8 (8%)	10	24
1	F	94/106 (89%)	90 (96%)	4 (4%)	26	51
1	G	102/106 (96%)	96 (94%)	6 (6%)	18	38
1	H	96/106 (91%)	95 (99%)	1 (1%)	68	86
1	I	94/106 (89%)	89 (95%)	5 (5%)	20	43
1	J	100/106 (94%)	93 (93%)	7 (7%)	14	31
1	K	98/106 (92%)	93 (95%)	5 (5%)	21	45
1	L	97/106 (92%)	97 (100%)	0	100	100
All	All	1176/1272 (92%)	1127 (96%)	49 (4%)	26	52

5 of 49 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	G	84	ASP
1	I	107	GLU
1	G	113	LYS
1	I	25	ASN
1	J	1	MET

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 28 such sidechains are listed below:

Mol	Chain	Res	Type
1	G	9	ASN
1	L	89	ASN
1	H	70	ASN
1	K	9	ASN
1	H	9	ASN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

There are no ligands in this entry.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	125/126 (99%)	-0.06	2 (1%) 70 66	19, 35, 65, 73	0
1	B	119/126 (94%)	-0.17	0 100 100	17, 33, 56, 68	0
1	C	119/126 (94%)	-0.19	1 (0%) 82 80	20, 33, 57, 65	0
1	D	126/126 (100%)	-0.12	1 (0%) 82 80	17, 36, 62, 73	0
1	E	120/126 (95%)	-0.15	1 (0%) 82 80	19, 35, 59, 78	0
1	F	118/126 (93%)	0.10	2 (1%) 69 64	20, 41, 68, 80	0
1	G	125/126 (99%)	-0.21	0 100 100	18, 31, 53, 57	0
1	H	122/126 (96%)	0.15	7 (5%) 29 24	16, 41, 80, 89	0
1	I	116/126 (92%)	0.39	1 (0%) 81 78	27, 52, 79, 81	0
1	J	124/126 (98%)	0.17	7 (5%) 30 24	24, 42, 76, 78	0
1	K	123/126 (97%)	0.31	3 (2%) 59 54	27, 51, 75, 88	0
1	L	119/126 (94%)	0.16	1 (0%) 82 80	25, 45, 73, 83	0
All	All	1456/1512 (96%)	0.03	26 (1%) 67 63	16, 39, 71, 89	0

The worst 5 of 26 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	K	84	ASP	4.4
1	A	85	PHE	3.9
1	J	97	GLY	3.3
1	J	14	ILE	2.9
1	E	43	VAL	2.8

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

There are no ligands in this entry.

6.5 Other polymers [i](#)

There are no such residues in this entry.