



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 10, 2026 – 12:03 AM UTC

PDB ID : 7DYY / pdb_00007dyy
Title : Crystal structure of VIM-2 MBL in complex with 1-((2-aminobenzo[d]thiazol-6-yl)methyl)-1H-imidazole-2-carboxylic acid
Authors : Yan, Y.-H.; Li, G.-B.
Deposited on : 2021-01-23
Resolution : 1.80 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Buster-report	:	wwPDB partial adaption of 1.1.7 (2018)
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.80 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	7662 (1.80-1.80)
Clashscore	190562	8479 (1.80-1.80)
Ramachandran outliers	187476	8391 (1.80-1.80)
Sidechain outliers	187428	8390 (1.80-1.80)
RSRZ outliers	180081	7663 (1.80-1.80)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	231	19% 75% 23% .
1	B	231	19% 74% 25% .

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
3	HQF	A	303	-	X	-	-

2 Entry composition [i](#)

There are 5 unique types of molecules in this entry. The entry contains 3651 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called Beta-lactamase class B VIM-2.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	231	Total	C	N	O	S	0	0	0
			1733	1094	299	339	1			
1	B	231	Total	C	N	O	S	0	0	0
			1733	1094	299	339	1			

- Molecule 2 is ZINC ION (CCD ID: ZN) (formula: Zn).


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	A	2	Total	Zn	0	0
			2	2		
2	B	2	Total	Zn	0	0
			2	2		

- Molecule 3 is 1-[(2-azanyl-1,3-benzothiazol-6-yl)methyl]imidazole-2-carboxylic acid (CCD ID: HQF) (formula: C₁₂H₁₀N₄O₂S) (labeled as "Ligand of Interest" by depositor).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
3	A	1	Total	C	N	O	S	0	0
			19	12	4	2	1		
3	B	1	Total	C	N	O	S	0	0
			19	12	4	2	1		

- Molecule 4 is DI(HYDROXYETHYL)ETHER (CCD ID: PEG) (formula: C₄H₁₀O₃).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
4	A	1	Total	C	O	0	0
			7	4	3		
4	B	1	Total	C	O	0	0
			7	4	3		


- Molecule 5 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	67	Total	O	0	0
			67	67		
5	B	62	Total	O	0	0
			62	62		


3 Residue-property plots

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: Beta-lactamase class B VIM-2



• Molecule 1: Beta-lactamase class B VIM-2



4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	61.12Å 45.08Å 98.03Å 90.00° 90.06° 90.00°	Depositor
Resolution (Å)	51.84 – 1.80 51.84 – 1.80	Depositor EDS
% Data completeness (in resolution range)	53.6 (51.84-1.80) 53.6 (51.84-1.80)	Depositor EDS
R_{merge}	0.48	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.10 (at 1.79Å)	Xtriage
Refinement program	PHENIX 1.10.1_2155	Depositor
R, R_{free}	0.232 , 0.298 (Not available) , 0.286	Depositor DCC
R_{free} test set	1987 reflections (3.97%)	wwPDB-VP
Wilson B-factor (Å ²)	23.2	Xtriage
Anisotropy	0.054	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.37 , 41.8	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.33$	Xtriage
Estimated twinning fraction	0.349 for h,-k,-l	Xtriage
F_o, F_c correlation	0.90	EDS
Total number of atoms	3651	wwPDB-VP
Average B, all atoms (Å ²)	29.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 7.91% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: PEG, ZN, HQF

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.60	1/1774 (0.1%)	0.76	0/2429
1	B	0.60	0/1774	0.74	0/2429
All	All	0.60	1/3548 (0.0%)	0.75	0/4858

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	241	GLY	C-N	5.77	1.45	1.33

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1733	0	1676	27	0
1	B	1733	0	1676	33	0
2	A	2	0	0	0	0
2	B	2	0	0	0	0
3	A	19	0	0	1	0
3	B	19	0	0	0	0
4	A	7	0	10	1	0
4	B	7	0	10	1	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
5	A	67	0	0	4	0
5	B	62	0	0	8	0
All	All	3651	0	3372	60	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 9.

All (60) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:210:ASN:O	4:B:304:PEG:O4	2.00	0.78
1:B:118:ASP:O	5:B:401:HOH:O	2.03	0.77
3:A:303:HQF:N16	5:A:401:HOH:O	2.20	0.75
1:B:182:ASP:O	5:B:402:HOH:O	2.10	0.69
1:B:32:GLU:O	5:B:403:HOH:O	2.13	0.67
1:A:210:ASN:HB2	4:A:304:PEG:H41	1.80	0.62
1:A:113:THR:O	1:A:183:ASN:HA	2.01	0.60
1:B:227:GLN:HB2	1:B:249:LEU:HD23	1.84	0.60
1:A:88:GLY:HA3	5:A:455:HOH:O	2.03	0.57
1:B:166:ARG:HH22	1:B:189:PRO:HG2	1.70	0.56
1:A:42:GLY:HA2	1:A:91:ASN:OD1	2.07	0.55
1:B:247:ASP:OD1	1:B:247:ASP:N	2.40	0.54
1:B:124:ASP:OD1	1:B:127:ARG:NH2	2.41	0.54
1:B:180:SER:HB2	5:B:411:HOH:O	2.09	0.53
1:A:157:GLY:O	1:A:163:ASP:HB3	2.09	0.52
1:A:127:ARG:NH1	1:A:152:THR:OG1	2.43	0.52
1:A:247:ASP:OD1	1:A:247:ASP:N	2.41	0.52
1:B:219:TRP:HB3	1:B:220:PRO:HD3	1.92	0.51
1:A:232:GLU:O	1:A:233:ALA:C	2.54	0.50
1:B:134:TYR:CD2	1:B:153:HIS:HB2	2.46	0.50
1:B:236:VAL:HG12	1:B:248:LEU:HD12	1.94	0.48
1:A:215:ASP:OD1	1:A:218:GLU:HB3	2.14	0.48
1:B:201:TYR:HE2	1:B:208:ALA:HA	1.78	0.48
1:B:213:ASP:HB3	5:B:411:HOH:O	2.14	0.47
1:B:187:TYR:CG	1:B:230:TYR:CE1	3.03	0.47
1:A:92:THR:O	1:A:95:LEU:HB3	2.14	0.47
1:B:171:GLU:OE1	1:B:230:TYR:OH	2.32	0.47
1:B:107:VAL:O	1:B:131:VAL:HG22	2.14	0.47
1:B:158:LEU:HD21	1:B:165:VAL:HG22	1.96	0.46
1:B:187:TYR:CD1	1:B:230:TYR:CD1	3.03	0.46
1:B:232:GLU:O	1:B:233:ALA:C	2.56	0.46

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:127:ARG:NH1	1:B:152:THR:OG1	2.49	0.46
1:B:224:GLU:O	1:B:228:GLN:HB2	2.16	0.46
1:A:193:VAL:HG22	1:A:235:PHE:CD1	2.52	0.45
1:A:230:TYR:N	1:A:231:PRO:CD	2.80	0.44
1:B:109:ARG:HB2	5:B:443:HOH:O	2.15	0.44
1:B:261:ASN:N	1:B:261:ASN:OD1	2.50	0.44
1:A:135:ALA:HB2	1:A:151:PRO:HG2	1.99	0.44
1:A:249:LEU:O	1:A:253:THR:OG1	2.31	0.44
1:B:232:GLU:HB2	5:B:414:HOH:O	2.17	0.44
1:A:160:SER:O	1:A:161:SER:C	2.59	0.44
1:B:49:ILE:HD13	1:B:237:ILE:HD11	2.00	0.43
1:B:139:THR:OG1	1:B:182:ASP:OD2	2.36	0.43
1:A:219:TRP:O	1:A:219:TRP:HD1	2.01	0.43
1:A:39:ILE:HB	1:A:59:THR:HG21	1.99	0.43
1:A:137:PRO:HD3	1:A:155:LEU:O	2.18	0.43
1:B:33:TYR:CG	1:B:34:PRO:HD2	2.54	0.42
1:A:65:ALA:HB1	1:B:65:ALA:HB1	2.01	0.42
1:A:179:HIS:ND1	5:A:403:HOH:O	2.34	0.42
1:A:56:HIS:ND1	1:A:56:HIS:O	2.53	0.42
1:B:157:GLY:O	1:B:163:ASP:HB3	2.20	0.42
1:B:84:ASP:CG	1:B:119:ARG:HD3	2.45	0.42
1:A:114:HIS:HB3	1:A:183:ASN:HA	2.02	0.41
1:A:37:SER:HB3	5:A:441:HOH:O	2.21	0.41
1:A:233:ALA:HB3	1:A:246:LEU:HD21	2.02	0.41
1:A:223:ILE:HG23	1:A:249:LEU:HG	2.02	0.41
1:A:179:HIS:CD2	1:A:198:CYS:HB2	2.56	0.41
1:B:232:GLU:N	5:B:414:HOH:O	2.45	0.41
1:B:192:SER:CB	1:B:234:GLN:HG2	2.51	0.40
1:A:192:SER:CB	1:A:234:GLN:HG2	2.51	0.40

There are no symmetry-related clashes.

5.3 Torsion angles ⓘ

5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	229/231 (99%)	217 (95%)	10 (4%)	2 (1%)	14	5
1	B	229/231 (99%)	215 (94%)	11 (5%)	3 (1%)	9	3
All	All	458/462 (99%)	432 (94%)	21 (5%)	5 (1%)	11	3

All (5) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	233	ALA
1	B	178	ALA
1	A	178	ALA
1	B	204	SER
1	B	243	PRO

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	182/184 (99%)	164 (90%)	18 (10%)	7	2
1	B	182/184 (99%)	170 (93%)	12 (7%)	15	5
All	All	364/368 (99%)	334 (92%)	30 (8%)	10	3

All (30) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	61	SER
1	A	90	LYS
1	A	123	VAL
1	A	141	ARG
1	A	154	SER
1	A	156	GLU
1	A	159	SER
1	A	183	ASN
1	A	188	VAL
1	A	204	SER
1	A	206	THR

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	A	210	ASN
1	A	228	GLN
1	A	235	PHE
1	A	237	ILE
1	A	242	LEU
1	A	253	THR
1	A	260	THR
1	B	37	SER
1	B	90	LYS
1	B	117	ASP
1	B	136	SER
1	B	154	SER
1	B	183	ASN
1	B	185	VAL
1	B	218	GLU
1	B	236	VAL
1	B	242	LEU
1	B	260	THR
1	B	261	ASN

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (2) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	229	HIS
1	B	48	GLN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no oligosaccharides in this entry.

5.6 Ligand geometry

Of 8 ligands modelled in this entry, 4 are monoatomic - leaving 4 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
4	PEG	A	304	-	6,6,6	0.51	0	5,5,5	0.60	0
3	HQF	B	303	2	20,21,21	2.21	6 (30%)	27,30,30	4.33	11 (40%)
4	PEG	B	304	-	6,6,6	0.50	0	5,5,5	0.60	0
3	HQF	A	303	2	20,21,21	2.39	7 (35%)	27,30,30	5.28	18 (66%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	PEG	A	304	-	-	2/4/4/4	-
3	HQF	B	303	2	-	0/8/8/8	0/3/3/3
4	PEG	B	304	-	-	2/4/4/4	-
3	HQF	A	303	2	-	2/8/8/8	0/3/3/3

All (13) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	A	303	HQF	C14-N16	6.69	1.47	1.34
3	B	303	HQF	C14-N16	5.69	1.45	1.34
3	A	303	HQF	C02-N03	-3.93	1.31	1.37
3	B	303	HQF	C14-S15	-3.55	1.72	1.76
3	B	303	HQF	C02-N03	-3.34	1.32	1.37
3	B	303	HQF	C10-N13	-3.06	1.33	1.39
3	A	303	HQF	C14-S15	3.05	1.79	1.76
3	A	303	HQF	C05-N01	-2.94	1.32	1.38
3	A	303	HQF	C04-N03	-2.67	1.32	1.38
3	B	303	HQF	C04-N03	-2.64	1.32	1.38
3	A	303	HQF	C10-N13	-2.61	1.34	1.39

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	A	303	HQF	O19-C17	2.59	1.29	1.22
3	B	303	HQF	C05-N01	-2.40	1.33	1.38

All (29) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	B	303	HQF	S15-C14-N13	-13.11	107.58	115.72
3	A	303	HQF	S15-C14-N13	-12.98	107.67	115.72
3	A	303	HQF	C09-S15-C14	12.59	94.84	88.89
3	B	303	HQF	C09-S15-C14	12.30	94.70	88.89
3	A	303	HQF	S15-C14-N16	10.58	131.63	120.13
3	B	303	HQF	C10-N13-C14	9.39	116.14	110.57
3	A	303	HQF	C06-C07-C12	-7.38	107.15	120.75
3	A	303	HQF	C07-C06-N03	6.74	123.30	112.84
3	A	303	HQF	C06-C07-C08	6.02	131.62	120.27
3	A	303	HQF	C10-N13-C14	5.43	113.79	110.57
3	A	303	HQF	C10-C09-S15	-5.09	102.23	109.75
3	B	303	HQF	S15-C14-N16	4.85	125.41	120.13
3	A	303	HQF	C08-C09-S15	4.79	138.03	126.07
3	A	303	HQF	N16-C14-N13	-4.35	120.70	124.13
3	A	303	HQF	C11-C10-N13	-3.70	117.69	128.21
3	B	303	HQF	N16-C14-N13	3.61	126.97	124.13
3	A	303	HQF	C09-C10-N13	3.60	121.47	115.20
3	A	303	HQF	C17-C02-N01	-3.54	120.07	124.59
3	B	303	HQF	C10-C09-S15	-3.20	105.02	109.75
3	A	303	HQF	C08-C09-C10	-3.04	119.73	121.99
3	B	303	HQF	C08-C09-S15	2.78	133.02	126.07
3	B	303	HQF	C04-C05-N01	-2.78	107.10	110.74
3	A	303	HQF	C05-N01-C02	2.76	107.81	105.08
3	A	303	HQF	C04-C05-N01	-2.60	107.33	110.74
3	B	303	HQF	C06-C07-C12	2.54	125.43	120.75
3	B	303	HQF	C06-C07-C08	-2.54	115.49	120.27
3	B	303	HQF	C05-N01-C02	2.52	107.57	105.08
3	A	303	HQF	C04-N03-C02	2.27	108.48	106.66
3	A	303	HQF	C09-C08-C07	-2.07	117.57	120.00

There are no chirality outliers.

All (6) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
4	A	304	PEG	O2-C3-C4-O4
4	B	304	PEG	O2-C3-C4-O4

Continued on next page...

Continued from previous page...


Mol	Chain	Res	Type	Atoms
4	A	304	PEG	C1-C2-O2-C3
4	B	304	PEG	C4-C3-O2-C2
3	A	303	HQF	N03-C06-C07-C08
3	A	303	HQF	N03-C06-C07-C12


There are no ring outliers.

3 monomers are involved in 3 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
4	A	304	PEG	1	0
4	B	304	PEG	1	0
3	A	303	HQF	1	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.





5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	231/231 (100%)	1.34	45 (19%) 3 2	19, 26, 43, 70	0
1	B	231/231 (100%)	1.38	45 (19%) 3 2	19, 27, 44, 70	0
All	All	462/462 (100%)	1.36	90 (19%) 3 2	19, 27, 44, 70	0

All (90) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	208	ALA	5.2
1	B	229	HIS	4.8
1	A	262	ARG	4.7
1	A	260	THR	4.2
1	B	260	THR	4.0
1	A	258	ALA	3.8
1	B	246	LEU	3.7
1	B	207	SER	3.7
1	B	259	HIS	3.6
1	B	233	ALA	3.4
1	B	258	ALA	3.4
1	B	203	LEU	3.4
1	B	262	ARG	3.4
1	A	200	ILE	3.4
1	A	212	ALA	3.3
1	B	261	ASN	3.3
1	A	164	ALA	3.3
1	A	53	VAL	3.3
1	B	209	GLY	3.2
1	A	150	ILE	3.2
1	A	259	HIS	3.0
1	B	145	VAL	3.0
1	A	237	ILE	3.0
1	A	181	THR	2.9

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	229	HIS	2.9
1	A	165	VAL	2.9
1	A	211	VAL	2.9
1	B	255	VAL	2.9
1	B	210	ASN	2.9
1	B	123	VAL	2.9
1	B	242	LEU	2.9
1	B	93	ALA	2.8
1	A	242	LEU	2.8
1	A	210	ASN	2.8
1	A	207	SER	2.8
1	A	145	VAL	2.8
1	A	203	LEU	2.8
1	A	250	LYS	2.8
1	B	237	ILE	2.7
1	B	208	ALA	2.7
1	A	62	PHE	2.6
1	B	170	VAL	2.6
1	A	93	ALA	2.6
1	A	201	TYR	2.6
1	A	246	LEU	2.5
1	B	110	ALA	2.5
1	B	211	VAL	2.5
1	A	147	GLY	2.5
1	B	89	ALA	2.5
1	A	41	VAL	2.5
1	B	228	GLN	2.4
1	A	54	TRP	2.4
1	A	167	PHE	2.4
1	B	128	ALA	2.4
1	B	132	ALA	2.4
1	A	170	VAL	2.4
1	A	151	PRO	2.4
1	B	37	SER	2.4
1	B	253	THR	2.4
1	A	99	ILE	2.4
1	B	188	VAL	2.3
1	A	261	ASN	2.3
1	B	113	THR	2.3
1	B	235	PHE	2.3
1	A	66	VAL	2.3
1	A	256	VAL	2.3

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	158	LEU	2.3
1	B	131	VAL	2.2
1	A	220	PRO	2.2
1	B	221	THR	2.2
1	A	120	VAL	2.2
1	B	250	LYS	2.2
1	B	127	ARG	2.1
1	A	111	VAL	2.1
1	B	135	ALA	2.1
1	B	256	VAL	2.1
1	A	117	ASP	2.1
1	A	138	SER	2.1
1	B	181	THR	2.1
1	A	231	PRO	2.1
1	A	255	VAL	2.0
1	B	36	VAL	2.0
1	B	165	VAL	2.0
1	B	185	VAL	2.0
1	A	206	THR	2.0
1	B	62	PHE	2.0
1	B	230	TYR	2.0
1	A	233	ALA	2.0
1	B	214	ALA	2.0
1	B	257	LYS	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)


There are no oligosaccharides in this entry.


6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
4	PEG	B	304	7/7	0.77	0.20	35,41,53,58	0
4	PEG	A	304	7/7	0.85	0.13	25,32,37,44	0
3	HQF	A	303	19/19	0.86	0.14	22,32,39,58	0
3	HQF	B	303	19/19	0.87	0.13	23,32,40,47	0
2	ZN	A	301	1/1	0.97	0.05	33,33,33,33	0
2	ZN	B	301	1/1	0.97	0.04	28,28,28,28	0
2	ZN	A	302	1/1	0.98	0.04	24,24,24,24	0
2	ZN	B	302	1/1	0.99	0.06	28,28,28,28	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.





6.5 Other polymers [i](#)

There are no such residues in this entry.