



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 8, 2026 – 09:04 AM UTC

PDB ID : 4DZ8 / pdb_00004dz8
Title : human IgG1 Fc fragment Heterodimer
Authors : Strop, P.
Deposited on : 2012-02-29
Resolution : 1.91 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0
Mogul : 2022.3.0, CSD as543be (2022)
Xtriage (Phenix) : 2.0
EDS : 3.0
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4 : 9.0.010 (Gargrove)
Density-Fitness : 1.0.12
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.91 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	1188 (1.92-1.92)
Clashscore	190562	1209 (1.92-1.92)
Ramachandran outliers	187476	1195 (1.92-1.92)
Sidechain outliers	187428	1195 (1.92-1.92)
RSRZ outliers	180081	1188 (1.92-1.92)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	225	4% 87% 6% • 6%
2	B	225	22% 76% 14% • 8%
3	C	8	62% 38%
3	D	8	38% 62%

2 Entry composition [i](#)

There are 4 unique types of molecules in this entry. The entry contains 3989 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Ig gamma-1 chain C region.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	209	Total	C	N	O	S	0	2	0
			1688	1072	286	324	6			

There are 3 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	228	ARG	PRO	engineered mutation	UNP P01857
A	368	GLU	LEU	microheterogeneity	UNP P01857
A	409	ARG	LYS	microheterogeneity	UNP P01857


- Molecule 2 is a protein called Ig gamma-1 chain C region.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	B	205	Total	C	N	O	S	0	2	0
			1667	1060	282	319	6			

There are 3 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
B	228	GLU	PRO	engineered mutation	UNP P01857
B	368	GLU	LEU	microheterogeneity	UNP P01857
B	409	ARG	LYS	microheterogeneity	UNP P01857

- Molecule 3 is an oligosaccharide called 2-acetamido-2-deoxy-beta-D-glucopyranose-(1-2)-alpha-D-mannopyranose-(1-3)-[2-acetamido-2-deoxy-beta-D-glucopyranose-(1-2)-alpha-D-mannopyranose-(1-6)]beta-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-[alpha-L-fucopyranose-(1-6)]2-acetamido-2-deoxy-beta-D-glucopyranose.



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
3	C	8	Total 99	C 56	N 4	O 39	0	0	0
3	D	8	Total 99	C 56	N 4	O 39	0	0	0



- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	269	Total 269	O 269	0	0
4	B	167	Total 167	O 167	0	0



3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Ig gamma-1 chain C region



- Molecule 2: Ig gamma-1 chain C region



- Molecule 3: 2-acetamido-2-deoxy-beta-D-glucopyranose-(1-2)-alpha-D-mannopyranose-(1-3)-[2-acetamido-2-deoxy-beta-D-glucopyranose-(1-2)-alpha-D-mannopyranose-(1-6)]beta-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-[alpha-L-fucopyranose-(1-6)]2-acetamido-2-deoxy-beta-D-glucopyranose



- Molecule 3: 2-acetamido-2-deoxy-beta-D-glucopyranose-(1-2)-alpha-D-mannopyranose-(1-3)-[2-acetamido-2-deoxy-beta-D-glucopyranose-(1-2)-alpha-D-mannopyranose-(1-6)]beta-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-[alpha-L-fucopyranose-(1-6)]2-acetamido-2-deoxy-beta-D-glucopyranose



4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	49.93Å 81.29Å 136.15Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	30.00 – 1.91 30.00 – 1.91	Depositor EDS
% Data completeness (in resolution range)	99.5 (30.00-1.91) 99.7 (30.00-1.91)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	0.08	Depositor
$\langle I/\sigma(I) \rangle$ ¹	3.50 (at 1.91Å)	Xtriage
Refinement program	REFMAC 5.2.0019	Depositor
R, R_{free}	0.205 , 0.237 0.211 , 0.244	Depositor DCC
R_{free} test set	2211 reflections (5.03%)	wwPDB-VP
Wilson B-factor (Å ²)	29.6	Xtriage
Anisotropy	0.050	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.30 , 40.1	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.95	EDS
Total number of atoms	3989	wwPDB-VP
Average B, all atoms (Å ²)	46.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 5.39% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: FUC, NAG, BMA, MAN

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.51	0/1732	0.66	0/2355
2	B	0.49	0/1710	0.66	0/2325
All	All	0.50	0/3442	0.66	0/4680

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1688	0	1648	8	0
2	B	1667	0	1630	22	0
3	C	99	0	85	0	0
3	D	99	0	85	0	0
4	A	269	0	0	1	0
4	B	167	0	0	1	0
All	All	3989	0	3448	30	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 5.

All (30) close contacts within the same asymmetric unit are listed below, sorted by their clash

magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:B:409[A]:ARG:HG2	2:B:409[A]:ARG:HH11	1.59	0.67
2:B:358:LEU:HD23	2:B:363:VAL:HG11	1.81	0.63
2:B:273:VAL:HG21	2:B:302:VAL:HG11	1.84	0.58
1:A:355:ARG:HA	1:A:358:LEU:HD12	1.87	0.57
1:A:429:HIS:CD2	1:A:431:ALA:H	2.22	0.56
2:B:259:VAL:HG13	2:B:336:ILE:HD11	1.88	0.55
2:B:409[A]:ARG:HH11	2:B:409[A]:ARG:CG	2.21	0.54
2:B:406:LEU:HD12	2:B:406:LEU:C	2.33	0.54
1:A:429:HIS:HD2	1:A:431:ALA:H	1.55	0.54
2:B:273:VAL:CG2	2:B:302:VAL:HG11	2.39	0.53
2:B:429:HIS:HD2	2:B:431:ALA:H	1.56	0.52
2:B:429:HIS:CD2	2:B:431:ALA:H	2.27	0.51
2:B:355:ARG:HA	2:B:358:LEU:HD12	1.95	0.49
1:A:374:PRO:O	1:A:429:HIS:HE1	1.96	0.48
2:B:263:VAL:HG22	2:B:323:VAL:HG21	1.96	0.48
2:B:287:ALA:HB2	2:B:306:LEU:HD13	1.96	0.47
2:B:242:LEU:HD13	2:B:336:ILE:HG12	1.96	0.47
2:B:311:GLN:NE2	4:B:764:HOH:O	2.46	0.47
2:B:258:GLU:HB3	2:B:305:VAL:HG13	1.95	0.47
2:B:330:ALA:HB1	2:B:331:PRO:HD2	1.97	0.46
1:A:406:LEU:C	1:A:406:LEU:HD12	2.42	0.45
1:A:272:GLU:HG2	4:A:754:HOH:O	2.16	0.44
2:B:242:LEU:HD13	2:B:336:ILE:HG23	2.00	0.44
2:B:301:ARG:NH1	2:B:303:VAL:HG21	2.33	0.43
2:B:339:ALA:HB3	2:B:374:PRO:HB3	2.01	0.43
1:A:325:ASN:HB3	1:A:328:LEU:HD22	2.00	0.43
1:A:252:MET:HG2	1:A:428:MET:HE1	2.01	0.42
2:B:418:GLN:HA	2:B:443:LEU:HD22	2.01	0.42
2:B:374:PRO:O	2:B:429:HIS:HE1	2.03	0.41
2:B:279:VAL:HG22	2:B:319:TYR:CD2	2.56	0.41

There are no symmetry-related clashes.

5.3 Torsion angles ⓘ

5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	205/225 (91%)	205 (100%)	0	0	100	100
2	B	201/225 (89%)	201 (100%)	0	0	100	100
All	All	406/450 (90%)	406 (100%)	0	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	196/208 (94%)	189 (96%)	7 (4%)	31	15
2	B	194/208 (93%)	186 (96%)	8 (4%)	27	12
All	All	390/416 (94%)	375 (96%)	15 (4%)	29	14

All (15) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	242	LEU
1	A	269	GLU
1	A	272	GLU
1	A	302	VAL
1	A	328	LEU
1	A	360	LYS
1	A	413	ASP
2	B	242	LEU
2	B	246	LYS
2	B	253	ILE
2	B	272	GLU
2	B	292	ARG
2	B	409[A]	ARG
2	B	409[B]	LYS
2	B	438	GLN

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (11) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	285	HIS
1	A	325	ASN
1	A	418	GLN
1	A	429	HIS
1	A	434	ASN
2	B	315	ASN
2	B	325	ASN
2	B	418	GLN
2	B	429	HIS
2	B	434	ASN
2	B	438	GLN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

16 monosaccharides are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
3	NAG	C	1	1,3	14,14,15	0.44	0	17,19,21	1.24	1 (5%)
3	NAG	C	2	3	14,14,15	0.64	0	17,19,21	1.00	1 (5%)
3	BMA	C	3	3	11,11,12	0.33	0	15,15,17	0.72	0
3	MAN	C	4	3	11,11,12	0.60	0	15,15,17	0.84	0
3	NAG	C	5	3	14,14,15	0.47	0	17,19,21	0.84	0

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
3	MAN	C	6	3	11,11,12	0.49	0	15,15,17	1.10	2 (13%)
3	NAG	C	7	3	14,14,15	0.53	0	17,19,21	0.78	0
3	FUC	C	8	3	10,10,11	0.52	0	14,14,16	0.79	0
3	NAG	D	1	3,2	14,14,15	0.57	0	17,19,21	0.83	0
3	NAG	D	2	3	14,14,15	0.59	0	17,19,21	1.07	1 (5%)
3	BMA	D	3	3	11,11,12	0.50	0	15,15,17	1.19	1 (6%)
3	MAN	D	4	3	11,11,12	0.59	0	15,15,17	0.97	1 (6%)
3	NAG	D	5	3	14,14,15	0.54	0	17,19,21	0.86	1 (5%)
3	MAN	D	6	3	11,11,12	0.48	0	15,15,17	0.76	0
3	NAG	D	7	3	14,14,15	0.51	0	17,19,21	0.72	0
3	FUC	D	8	3	10,10,11	0.57	0	14,14,16	1.63	3 (21%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	NAG	C	1	1,3	-	0/6/23/26	0/1/1/1
3	NAG	C	2	3	-	0/6/23/26	0/1/1/1
3	BMA	C	3	3	-	0/2/19/22	0/1/1/1
3	MAN	C	4	3	-	0/2/19/22	0/1/1/1
3	NAG	C	5	3	-	4/6/23/26	0/1/1/1
3	MAN	C	6	3	-	0/2/19/22	0/1/1/1
3	NAG	C	7	3	-	1/6/23/26	0/1/1/1
3	FUC	C	8	3	-	-	0/1/1/1
3	NAG	D	1	3,2	-	2/6/23/26	0/1/1/1
3	NAG	D	2	3	-	1/6/23/26	0/1/1/1
3	BMA	D	3	3	-	0/2/19/22	0/1/1/1
3	MAN	D	4	3	-	2/2/19/22	0/1/1/1
3	NAG	D	5	3	-	2/6/23/26	0/1/1/1
3	MAN	D	6	3	-	0/2/19/22	0/1/1/1
3	NAG	D	7	3	-	0/6/23/26	0/1/1/1
3	FUC	D	8	3	-	-	0/1/1/1

There are no bond length outliers.

All (11) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	C	1	NAG	C1-O5-C5	4.14	117.73	112.19

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	D	8	FUC	O5-C5-C4	3.92	116.60	109.55
3	D	3	BMA	C1-C2-C3	3.17	114.26	109.64
3	D	8	FUC	C1-O5-C5	2.97	119.97	112.97
3	D	2	NAG	O5-C1-C2	-2.86	106.87	111.29
3	D	8	FUC	O5-C1-C2	2.43	116.58	110.79
3	D	4	MAN	C1-O5-C5	2.40	115.40	112.19
3	C	6	MAN	O2-C2-C3	-2.25	105.49	110.15
3	C	6	MAN	C1-O5-C5	2.21	115.15	112.19
3	D	5	NAG	C1-O5-C5	2.15	115.06	112.19
3	C	2	NAG	O7-C7-C8	-2.06	118.38	122.05

There are no chirality outliers.


All (12) torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
3	C	5	NAG	C8-C7-N2-C2
3	C	5	NAG	O7-C7-N2-C2
3	D	1	NAG	C8-C7-N2-C2
3	D	1	NAG	O7-C7-N2-C2
3	D	5	NAG	C8-C7-N2-C2
3	D	5	NAG	O7-C7-N2-C2
3	C	5	NAG	O5-C5-C6-O6
3	C	5	NAG	C4-C5-C6-O6
3	D	4	MAN	C4-C5-C6-O6
3	C	7	NAG	C1-C2-N2-C7
3	D	2	NAG	C4-C5-C6-O6
3	D	4	MAN	O5-C5-C6-O6

There are no ring outliers.

No monomer is involved in short contacts.

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for oligosaccharide.





5.6 Ligand geometry [i](#)

There are no ligands in this entry.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	207/225 (92%)	0.17	6 (2%) 53 60	18, 29, 58, 67	0
2	B	203/225 (90%)	1.09	49 (24%) 2 2	20, 45, 148, 201	0
All	All	410/450 (91%)	0.63	55 (13%) 7 9	18, 34, 127, 201	0

All (55) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
2	B	328	LEU	5.6
2	B	273	VAL	5.0
2	B	264	VAL	4.8
2	B	300	TYR	4.6
2	B	271	PRO	4.6
2	B	266	VAL	4.5
2	B	291	PRO	4.3
2	B	296	TYR	4.2
2	B	292	ARG	3.9
1	A	444	SER	3.9
2	B	239	SER	3.9
2	B	323	VAL	3.9
2	B	275	PHE	3.8
2	B	268	HIS	3.6
2	B	262	VAL	3.6
2	B	299	THR	3.5
2	B	327	ALA	3.5
2	B	302	VAL	3.4
2	B	267	SER	3.2
2	B	332	ILE	3.1
2	B	301	ARG	3.1
1	A	236	GLY	3.1
2	B	325	ASN	3.0
2	B	287	ALA	2.9

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	358	LEU	2.9
2	B	326	LYS	2.8
2	B	324	SER	2.8
2	B	274	LYS	2.8
2	B	298	SER	2.8
2	B	303	VAL	2.8
2	B	269	GLU	2.7
2	B	294	GLU	2.7
2	B	261	CYS	2.7
2	B	331	PRO	2.6
2	B	329	PRO	2.5
2	B	278	TYR	2.5
2	B	358	LEU	2.5
2	B	263	VAL	2.5
2	B	276	ASN	2.5
2	B	336	ILE	2.4
2	B	285	HIS	2.4
2	B	272	GLU	2.3
2	B	361	ASN	2.3
2	B	385	GLY	2.3
2	B	284	VAL	2.3
2	B	295	GLN	2.3
2	B	277	TRP	2.3
2	B	282	VAL	2.3
1	A	269	GLU	2.2
1	A	327	ALA	2.2
1	A	361	ASN	2.2
2	B	286	ASN	2.1
2	B	297	ASN	2.1
2	B	356	ASP	2.1
2	B	359	THR	2.1

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)




In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.


Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
3	FUC	D	8	10/11	0.40	0.17	74,76,77,77	0
3	NAG	D	1	14/15	0.54	0.16	63,65,68,72	0
3	NAG	D	2	14/15	0.67	0.16	54,58,60,61	0
3	NAG	D	5	14/15	0.69	0.13	57,59,60,60	0
3	NAG	C	5	14/15	0.74	0.13	53,56,57,57	0
3	BMA	D	3	11/12	0.83	0.10	47,50,51,52	0
3	NAG	D	7	14/15	0.84	0.11	55,55,57,57	0
3	MAN	D	4	11/12	0.84	0.10	44,47,49,53	0
3	MAN	D	6	11/12	0.86	0.10	53,55,55,56	0
3	MAN	C	4	11/12	0.89	0.09	38,42,45,48	0
3	FUC	C	8	10/11	0.89	0.09	34,35,37,38	0
3	NAG	C	7	14/15	0.93	0.07	28,30,31,32	0
3	MAN	C	6	11/12	0.93	0.07	27,28,30,32	0
3	NAG	C	2	14/15	0.94	0.07	23,25,27,28	0
3	BMA	C	3	11/12	0.94	0.06	26,27,29,32	0
3	NAG	C	1	14/15	0.94	0.07	27,30,36,37	0

The following is a graphical depiction of the model fit to experimental electron density for oligosaccharide. Each fit is shown from different orientation to approximate a three-dimensional view.

Electron density around Chain C:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)





6.4 Ligands [i](#)

There are no ligands in this entry.

6.5 Other polymers [i](#)

There are no such residues in this entry.