



Full wwPDB X-ray Structure Validation Report ⓘ

Apr 24, 2026 – 09:49 PM EDT

PDB ID : 1E1H / pdb_00001e1h
Title : Crystal Structure of recombinant Botulinum Neurotoxin Type A Light Chain,
self-inhibiting Zn endopeptidase.
Authors : Knapp, M.; Rupp, B.
Deposited on : 2000-05-08
Resolution : 1.80 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.80 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
Clashscore	190562	8479 (1.80-1.80)
Ramachandran outliers	187476	8391 (1.80-1.80)
Sidechain outliers	187428	8390 (1.80-1.80)
RSRZ outliers	180081	7663 (1.80-1.80)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	287	5% 75% 7% 18%
1	C	287	4% 77% 5% 18%
2	B	174	10% 75% 16% 6%
2	D	174	9% 81% 11% 5%

2 Entry composition

There are 4 unique types of molecules in this entry. The entry contains 7230 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called BOTULINUM NEUROTOXIN TYPE A LIGHT CHAIN.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	235	Total	C	N	O	S	0	0	0
			1864	1193	309	357	5			
1	C	235	Total	C	N	O	S	0	0	0
			1864	1193	309	357	5			

- Molecule 2 is a protein called BOTULINUM NEUROTOXIN TYPE A LIGHT CHAIN.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	B	164	Total	C	N	O	S	0	0	0
			1360	875	222	260	3			
2	D	165	Total	C	N	O	S	0	0	0
			1372	884	223	262	3			

- Molecule 3 is ZINC ION (CCD ID: ZN) (formula: Zn).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	1	Total	Zn	0	0
			1	1		
3	C	1	Total	Zn	0	0
			1	1		


- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	215	Total	O	0	0
			215	215		
4	B	172	Total	O	0	0
			172	172		
4	C	221	Total	O	0	0
			221	221		
4	D	160	Total	O	0	0
			160	160		


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


• Molecule 1: BOTULINUM NEUROTOXIN TYPE A LIGHT CHAIN




• Molecule 1: BOTULINUM NEUROTOXIN TYPE A LIGHT CHAIN



• Molecule 2: BOTULINUM NEUROTOXIN TYPE A LIGHT CHAIN



• Molecule 2: BOTULINUM NEUROTOXIN TYPE A LIGHT CHAIN



HIS
HIS
HIS
HIS
HIS
HIS

4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	58.06Å 94.26Å 100.16Å 90.00° 103.52° 90.00°	Depositor
Resolution (Å)	19.34 – 1.80 19.34 – 1.80	Depositor EDS
% Data completeness (in resolution range)	95.2 (19.34-1.80) 94.8 (19.34-1.80)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	0.04	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.51 (at 1.80Å)	Xtriage
Refinement program	REFMAC	Depositor
R, R_{free}	0.196 , 0.237 0.190 , (Not available)	Depositor DCC
R_{free} test set	No test flags present.	wwPDB-VP
Wilson B-factor (Å ²)	23.3	Xtriage
Anisotropy	0.502	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.40 , 52.0	EDS
L-test for twinning ²	$\langle L \rangle = 0.48$, $\langle L^2 \rangle = 0.31$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.95	EDS
Total number of atoms	7230	wwPDB-VP
Average B, all atoms (Å ²)	30.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.51% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: ZN

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	1.04	1/1910 (0.1%)	1.26	1/2596 (0.0%)
1	C	1.03	0/1910	1.24	1/2596 (0.0%)
2	B	1.01	0/1385	1.32	8/1857 (0.4%)
2	D	0.96	1/1398 (0.1%)	1.21	4/1875 (0.2%)
All	All	1.02	2/6603 (0.0%)	1.26	14/8924 (0.2%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	1
1	C	0	1
All	All	0	2

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	D	378	VAL	CA-CB	-5.84	1.51	1.54
1	A	249	TYR	C-OXT	5.45	1.34	1.23

All (14) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	B	269	ASP	CA-CB-CG	17.67	130.27	112.60
2	B	268	HIS	CA-CB-CG	-7.40	106.40	113.80
2	B	362	ARG	CD-NE-CZ	6.38	133.34	124.40
2	D	315	VAL	N-CA-CB	6.37	119.20	110.54
2	D	376	ASN	CA-CB-CG	6.26	118.86	112.60

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	D	391	LEU	N-CA-CB	6.20	119.22	110.36
2	B	304	GLY	N-CA-C	-6.09	103.80	112.81
1	C	226	HIS	CE1-NE2-CD2	5.73	114.73	109.00
1	A	144	ARG	CA-CB-CG	5.45	125.01	114.10
2	B	315	VAL	N-CA-CB	5.42	117.91	110.54
2	B	361	ASN	CA-CB-CG	5.28	117.88	112.60
2	B	368	PHE	CA-CB-CG	-5.06	108.74	113.80
2	D	362	ARG	NE-CZ-NH2	5.05	123.75	119.20
2	B	268	HIS	N-CA-C	5.04	116.46	111.07

There are no chirality outliers.

All (2) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	20	TYR	Mainchain
1	C	20	TYR	Mainchain

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1864	0	1820	15	0
1	C	1864	0	1820	12	0
2	B	1360	0	1340	20	0
2	D	1372	0	1349	16	0
3	A	1	0	0	0	0
3	C	1	0	0	0	0
4	A	215	0	0	5	0
4	B	172	0	0	6	0
4	C	221	0	0	2	0
4	D	160	0	0	2	0
All	All	7230	0	6329	53	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 4.

All (53) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:197:GLU:HG2	2:B:360:ILE:HD11	1.43	0.99
1:C:197:GLU:HG2	2:D:360:ILE:HD11	1.48	0.95
2:B:269:ASP:HB2	4:B:2035:HOH:O	1.69	0.93
2:D:386:LYS:HZ3	2:D:386:LYS:H	1.28	0.82
1:C:63:GLU:HA	1:C:63:GLU:OE1	1.91	0.70
1:C:124:THR:HG21	4:C:2131:HOH:O	1.90	0.69
2:D:361:ASN:O	2:D:363:LYS:HE2	1.92	0.68
1:A:63:GLU:HA	1:A:63:GLU:OE1	1.94	0.66
1:A:122:ILE:HG22	1:A:124:THR:HG22	1.83	0.61
4:C:2104:HOH:O	2:D:252:MET:HE2	2.03	0.59
1:A:160:ILE:HG21	2:B:372:VAL:HG11	1.86	0.57
2:B:363:LYS:CE	4:B:2129:HOH:O	2.54	0.56
2:D:378:VAL:HB	2:D:379:PRO:HD3	1.88	0.56
2:B:386:LYS:H	2:B:386:LYS:NZ	2.05	0.55
2:B:273:ILE:HG22	2:B:278:GLU:HG3	1.89	0.54
2:B:384:THR:HB	2:B:386:LYS:HZ3	1.74	0.53
2:D:252:MET:HE3	4:D:2030:HOH:O	2.11	0.51
1:C:197:GLU:CG	2:D:360:ILE:HD11	2.31	0.49
1:A:21:ILE:C	1:A:21:ILE:HD12	2.38	0.49
2:B:367:ASN:H	1:C:246:THR:HG1	1.60	0.48
1:A:7:MET:N	4:A:2001:HOH:O	2.45	0.48
2:B:406:GLU:HG3	4:B:2161:HOH:O	2.13	0.47
1:A:24:PRO:HB2	4:A:2023:HOH:O	2.15	0.47
1:A:125:GLU:OE1	2:B:303:ILE:HD11	2.15	0.47
2:D:386:LYS:HE2	4:D:2146:HOH:O	2.15	0.47
1:A:240:ARG:HD3	4:A:2035:HOH:O	2.15	0.47
1:C:160:ILE:HG21	2:D:372:VAL:HG11	1.95	0.47
1:C:197:GLU:HG2	2:D:360:ILE:CD1	2.34	0.46
1:C:21:ILE:C	1:C:21:ILE:HD12	2.40	0.46
2:D:378:VAL:N	2:D:379:PRO:CD	2.79	0.45
1:C:24:PRO:O	1:C:25:ASN:HB2	2.17	0.44
1:A:24:PRO:O	1:A:25:ASN:HB2	2.16	0.44
2:B:378:VAL:HB	2:B:379:PRO:HD3	2.00	0.44
2:B:368:PHE:CZ	2:D:255:LEU:HG	2.52	0.43
1:C:28:GLN:H	1:C:28:GLN:HG2	1.57	0.43
2:B:334:LYS:HG3	4:B:2052:HOH:O	2.18	0.43
2:B:398:THR:O	2:B:401:ASN:ND2	2.51	0.43
2:B:319:LYS:HD3	2:B:320:TYR:CE2	2.52	0.43
1:A:52:ASN:HB3	1:A:55:GLU:HB2	2.00	0.43
1:A:127:LYS:NZ	4:A:2127:HOH:O	2.52	0.42
1:A:23:ILE:HA	1:A:24:PRO:HD3	1.92	0.42
2:D:319:LYS:HD3	2:D:320:TYR:CE2	2.53	0.42

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:D:398:THR:O	2:D:401:ASN:ND2	2.53	0.42
1:A:24:PRO:HD2	4:A:2023:HOH:O	2.19	0.42
2:D:386:LYS:HD3	2:D:386:LYS:N	2.35	0.42
1:C:97:ILE:O	1:C:103:GLY:HA3	2.20	0.41
1:A:197:GLU:O	1:A:198:SER:HB2	2.20	0.41
2:B:277:GLN:HE21	2:B:277:GLN:HB2	1.65	0.41
2:B:284:TYR:CZ	2:B:288:LYS:HE2	2.56	0.41
1:C:125:GLU:OE1	2:D:303:ILE:HD11	2.20	0.41
2:B:386:LYS:HE2	4:B:2153:HOH:O	2.20	0.41
2:B:263:ARG:HH11	2:B:263:ARG:HD3	1.74	0.41
2:B:414:ARG:NE	4:B:2170:HOH:O	2.54	0.40

There are no symmetry-related clashes.

5.3 Torsion angles

5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	231/287 (80%)	223 (96%)	7 (3%)	1 (0%)	30	19
1	C	231/287 (80%)	222 (96%)	8 (4%)	1 (0%)	30	19
2	B	160/174 (92%)	157 (98%)	3 (2%)	0	100	100
2	D	161/174 (92%)	159 (99%)	2 (1%)	0	100	100
All	All	783/922 (85%)	761 (97%)	20 (3%)	2 (0%)	36	25

All (2) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	27	GLY
1	C	27	GLY

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	205/249 (82%)	202 (98%)	3 (2%)	57	49
1	C	205/249 (82%)	203 (99%)	2 (1%)	68	64
2	B	151/160 (94%)	141 (93%)	10 (7%)	15	5
2	D	152/160 (95%)	142 (93%)	10 (7%)	15	5
All	All	713/818 (87%)	688 (96%)	25 (4%)	32	19

All (25) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	28	GLN
1	A	63	GLU
1	A	144	ARG
2	B	258	SER
2	B	271	LYS
2	B	277	GLN
2	B	314	ASN
2	B	315	VAL
2	B	334	LYS
2	B	335	LEU
2	B	386	LYS
2	B	391	LEU
2	B	406	GLU
1	C	28	GLN
1	C	63	GLU
2	D	258	SER
2	D	271	LYS
2	D	277	GLN
2	D	315	VAL
2	D	334	LYS
2	D	335	LEU
2	D	363	LYS
2	D	386	LYS
2	D	391	LEU
2	D	406	GLU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (15) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	25	ASN
1	A	28	GLN
1	A	66	GLN
1	A	85	ASN
1	A	135	ASN
2	B	277	GLN
2	B	287	ASN
2	B	314	ASN
2	B	376	ASN
1	C	28	GLN
1	C	66	GLN
1	C	81	ASN
1	C	85	ASN
2	D	277	GLN
2	D	376	ASN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 2 ligands modelled in this entry, 2 are monoatomic - leaving 0 for Mogul analysis.

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2		OWAB(Å ²)	Q<0.9
1	A	235/287 (81%)	0.21	15 (6%)	25 24	16, 25, 48, 64	0
1	C	235/287 (81%)	0.19	12 (5%)	33 32	16, 25, 49, 64	0
2	B	164/174 (94%)	0.65	17 (10%)	11 10	18, 29, 44, 62	0
2	D	165/174 (94%)	0.52	15 (9%)	15 13	18, 30, 45, 66	0
All	All	799/922 (86%)	0.36	59 (7%)	20 19	16, 27, 47, 66	0

All (59) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
2	B	394	ALA	7.1
2	B	305	THR	6.0
1	A	28	GLN	5.8
2	B	306	THR	5.4
1	A	207	LEU	5.3
2	B	307	ALA	5.0
1	C	209	ALA	4.9
2	B	395	ASN	4.8
1	A	208	GLY	4.7
2	D	395	ASN	4.3
2	D	394	ALA	4.1
1	A	64	ALA	3.9
1	C	208	GLY	3.9
1	C	198	SER	3.7
2	B	304	GLY	3.7
2	B	335	LEU	3.6
1	A	198	SER	3.6
1	C	28	GLN	3.6
1	A	121	THR	3.5
2	D	250	TYR	3.4
1	A	209	ALA	3.4

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	26	ALA	3.3
2	D	407	ILE	3.2
1	C	207	LEU	2.9
1	A	24	PRO	2.8
1	A	122	ILE	2.8
2	B	391	LEU	2.7
2	B	355	ASN	2.7
2	B	268	HIS	2.7
1	C	210	GLY	2.7
1	A	197	GLU	2.7
2	D	392	LYS	2.6
2	D	396	LEU	2.6
1	C	27	GLY	2.6
2	D	391	LEU	2.5
1	A	171	VAL	2.5
2	B	251	GLU	2.5
2	B	269	ASP	2.5
2	D	269	ASP	2.5
2	B	398	THR	2.4
1	A	63	GLU	2.4
1	A	27	GLY	2.3
1	C	63	GLU	2.3
2	B	396	LEU	2.3
1	C	71	TYR	2.3
2	D	415	LEU	2.3
2	D	401	ASN	2.3
2	D	399	ASN	2.2
2	D	306	THR	2.2
1	C	62	PRO	2.2
1	C	197	GLU	2.2
1	A	81	ASN	2.2
2	B	303	ILE	2.1
2	B	294	SER	2.1
1	C	143	TYR	2.0
2	B	276	LEU	2.0
2	D	335	LEU	2.0
2	D	369	ASP	2.0
2	D	398	THR	2.0

6.2 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
3	ZN	C	502	1/1	0.99	0.04	22,22,22,22	0
3	ZN	A	501	1/1	1.00	0.04	20,20,20,20	0

6.5 Other polymers [i](#)

There are no such residues in this entry.