



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 8, 2026 – 08:31 AM UTC

PDB ID : 1E1K / pdb_00001e1k
Title : ADRENODOXIN REDUCTASE in complex with NADP+ obtained by a soaking experiment
Authors : Ziegler, G.A.; Schulz, G.E.
Deposited on : 2000-05-09
Resolution : 1.95 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0
Mogul : 2022.3.0, CSD as543be (2022)
Xtriage (Phenix) : 2.0
EDS : 3.0
Buster-report : wwPDB partial adaption of 1.1.7 (2018)
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4 : 9.0.010 (Gargrove)
Density-Fitness : 1.0.12
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.95 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	3494 (1.96-1.96)
Clashscore	190562	3612 (1.96-1.96)
Ramachandran outliers	187476	3587 (1.96-1.96)
Sidechain outliers	187428	3587 (1.96-1.96)
RSRZ outliers	180081	3495 (1.96-1.96)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	460	

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
3	NAP	A	802	X	-	-	-

2 Entry composition [i](#)


There are 4 unique types of molecules in this entry. The entry contains 3959 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called ADRENODOXIN REDUCTASE.


Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	455	Total	C	N	O	S	0	1	0
			3512	2219	637	645	11			

- Molecule 2 is FLAVIN-ADENINE DINUCLEOTIDE (CCD ID: FAD) (formula: $C_{27}H_{33}N_9O_{15}P_2$).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
2	A	1	Total	C	N	O	P	0	0
			53	27	9	15	2		

- Molecule 3 is NADP NICOTINAMIDE-ADENINE-DINUCLEOTIDE PHOSPHATE (CCD ID: NAP) (formula: $C_{21}H_{28}N_7O_{17}P_3$).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
3	A	1	Total 48	C 21	N 7	O 17	P 3	0	0


- Molecule 4 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
4	A	346	Total O 346 346	0	0

3 Residue-property plots

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: ADRENODOXIN REDUCTASE



4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	60.90Å 62.40Å 78.20Å 90.00° 106.60° 90.00°	Depositor
Resolution (Å)	20.00 – 1.95 20.00 – 1.95	Depositor EDS
% Data completeness (in resolution range)	100.0 (20.00-1.95) 99.7 (20.00-1.95)	Depositor EDS
R_{merge}	0.08	Depositor
R_{sym}	0.07	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.69 (at 1.95Å)	Xtriage
Refinement program	REFMAC	Depositor
R, R_{free}	0.184 , 0.227 0.184 , 0.225	Depositor DCC
R_{free} test set	2108 reflections (5.13%)	wwPDB-VP
Wilson B-factor (Å ²)	18.5	Xtriage
Anisotropy	0.462	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.38 , 41.9	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.95	EDS
Total number of atoms	3959	wwPDB-VP
Average B, all atoms (Å ²)	26.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 6.65% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: FAD, NAP

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.80	0/3595	1.71	50/4885 (1.0%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	1

There are no bond length outliers.

All (50) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	31	ARG	CD-NE-CZ	15.04	145.46	124.40
1	A	372	PRO	CA-C-O	-10.40	113.61	121.31
1	A	106	ASP	CA-CB-CG	9.85	122.45	112.60
1	A	70	ARG	CD-NE-CZ	9.10	137.14	124.40
1	A	188	ARG	CD-NE-CZ	8.46	136.24	124.40
1	A	333	SER	N-CA-C	-8.44	99.64	110.53
1	A	283	VAL	CA-C-O	-8.03	111.70	120.85
1	A	331	TYR	N-CA-C	7.71	120.45	108.96
1	A	195	VAL	N-CA-C	7.70	119.24	107.99
1	A	414	LEU	O-C-N	7.07	130.21	122.15
1	A	278	ARG	NE-CZ-NH2	7.05	125.55	119.20
1	A	185	ARG	NE-CZ-NH2	-6.89	113.00	119.20
1	A	278	ARG	NH1-CZ-NH2	-6.76	110.52	119.30
1	A	337	ASP	CA-CB-CG	6.57	119.17	112.60
1	A	414	LEU	CA-C-O	-6.53	113.50	120.42
1	A	373	THR	CA-C-N	-6.33	114.94	122.16

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	373	THR	C-N-CA	-6.33	114.94	122.16
1	A	403	PRO	N-CA-C	6.24	125.31	112.47
1	A	60	ASN	CA-CB-CG	-5.92	106.68	112.60
1	A	233	ASP	CA-CB-CG	-5.91	106.69	112.60
1	A	140	ALA	O-C-N	-5.80	118.86	121.53
1	A	331	TYR	CA-C-O	5.75	127.61	121.45
1	A	249	LEU	CB-CA-C	-5.68	101.95	110.88
1	A	256	LYS	CA-CB-CG	5.68	125.45	114.10
1	A	108	GLN	CA-C-O	-5.63	114.33	120.81
1	A	321	PRO	CA-C-O	-5.62	114.93	121.34
1	A	233	ASP	CA-C-O	-5.61	114.93	120.82
1	A	40	GLN	OE1-CD-NE2	-5.61	116.99	122.60
1	A	403	PRO	CA-C-O	-5.59	110.42	120.60
1	A	334	ARG	N-CA-CB	-5.58	99.42	110.70
1	A	162	ARG	NE-CZ-NH2	-5.42	114.33	119.20
1	A	309	THR	N-CA-CB	5.38	117.87	109.97
1	A	83	GLY	N-CA-C	-5.33	108.35	114.69
1	A	157	ALA	CA-C-O	5.31	126.18	120.55
1	A	256	LYS	N-CA-C	-5.28	102.99	109.65
1	A	185	ARG	NH1-CZ-NH2	5.27	126.16	119.30
1	A	13	GLY	CA-C-O	-5.26	115.14	121.98
1	A	57	GLU	CA-CB-CG	5.23	124.56	114.10
1	A	310	ARG	CD-NE-CZ	-5.14	117.21	124.40
1	A	334	ARG	CA-CB-CG	5.13	124.36	114.10
1	A	370	ARG	NH1-CZ-NH2	-5.10	112.67	119.30
1	A	254	THR	CA-C-O	-5.09	113.52	118.97
1	A	99	VAL	CB-CA-C	-5.09	103.54	110.77
1	A	434	GLU	N-CA-CB	5.07	117.57	110.12
1	A	202	GLN	OE1-CD-NE2	-5.07	117.53	122.60
1	A	108	GLN	OE1-CD-NE2	5.06	127.66	122.60
1	A	172	GLU	CB-CA-C	-5.05	102.40	110.79
1	A	25	LEU	CA-C-O	-5.04	115.08	120.42
1	A	311	ALA	CA-C-O	-5.01	114.97	120.69
1	A	150	ILE	CA-C-O	-5.01	115.17	120.48

There are no chirality outliers.

All (1) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	330	GLY	Mainchain

5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3512	0	3544	45	0
2	A	53	0	31	0	0
3	A	48	0	24	3	0
4	A	346	0	0	8	0
All	All	3959	0	3599	45	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 6.

All (45) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:306:GLY:O	1:A:309:THR:HG23	1.85	0.77
1:A:124[A]:ARG:HD2	1:A:331:TYR:HE1	1.50	0.76
1:A:107:HIS:ND1	1:A:124[B]:ARG:NH2	2.36	0.73
1:A:160:VAL:HG11	1:A:326:LEU:HD13	1.71	0.73
1:A:160:VAL:HG11	1:A:326:LEU:CD1	2.19	0.72
1:A:215:GLN:HG2	4:A:2186:HOH:O	1.90	0.70
1:A:309:THR:C	1:A:310:ARG:HD2	2.19	0.67
1:A:29:HIS:HD2	1:A:31:ARG:H	1.41	0.65
1:A:104:ALA:HA	1:A:333:SER:HA	1.80	0.64
1:A:124[B]:ARG:HD3	1:A:331:TYR:HE1	1.64	0.63
1:A:209:GLU:OE2	3:A:802:NAP:N7N	2.33	0.61
1:A:106:ASP:O	1:A:331:TYR:HD1	1.84	0.60
1:A:124[A]:ARG:HD2	1:A:331:TYR:CE1	2.33	0.59
1:A:77:TYR:CE1	1:A:411:LYS:HE2	2.39	0.57
1:A:229:LEU:HD22	4:A:2192:HOH:O	2.05	0.56
1:A:433:GLU:HG2	1:A:457:LEU:HD13	1.89	0.55
1:A:124[B]:ARG:HD3	1:A:331:TYR:CE1	2.43	0.53
1:A:108:GLN:HE22	1:A:332:LYS:HE3	1.75	0.52
1:A:282:GLN:OE1	4:A:2243:HOH:O	2.19	0.51
1:A:10:CYS:SG	1:A:98:VAL:HG22	2.51	0.51
1:A:404:ARG:HA	4:A:2300:HOH:O	2.10	0.51
1:A:309:THR:HG21	4:A:2251:HOH:O	2.12	0.50
1:A:352:MET:HA	1:A:352:MET:HE2	1.95	0.49

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:29:HIS:CD2	1:A:31:ARG:H	2.27	0.48
1:A:309:THR:O	1:A:310:ARG:HD2	2.13	0.48
1:A:373:THR:O	3:A:802:NAP:O3D	2.32	0.47
1:A:396:ALA:HB1	1:A:398:HIS:CE1	2.50	0.47
1:A:375:VAL:HG12	3:A:802:NAP:H2N	1.95	0.47
1:A:215:GLN:NE2	4:A:2184:HOH:O	2.48	0.47
1:A:331:TYR:HD2	4:A:2290:HOH:O	1.98	0.46
1:A:231:LEU:HD23	1:A:250:LEU:HG	1.97	0.46
1:A:259:VAL:HG23	4:A:2214:HOH:O	2.16	0.45
1:A:55:HIS:N	1:A:56:PRO:CD	2.79	0.45
1:A:29:HIS:CD2	1:A:32:ALA:H	2.36	0.43
1:A:54:ASP:C	1:A:56:PRO:CD	2.91	0.42
1:A:211:ARG:HG2	1:A:215:GLN:NE2	2.34	0.42
1:A:288:ASP:CG	1:A:290:ARG:HG3	2.44	0.42
1:A:110:LEU:O	1:A:111:ASP:HB2	2.19	0.42
1:A:117:LEU:HA	1:A:118:PRO:HD3	1.86	0.41
1:A:19:PHE:HB3	1:A:65:PHE:CD1	2.56	0.41
1:A:229:LEU:O	1:A:229:LEU:HD23	2.20	0.41
1:A:224:ASP:HA	1:A:225:PRO:HD2	1.91	0.40
1:A:240:ARG:HB2	1:A:241:PRO:HD3	2.03	0.40
1:A:194:ILE:HB	1:A:274:LEU:HD12	2.04	0.40
1:A:223:LEU:HD21	1:A:274:LEU:HD22	2.03	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	454/460 (99%)	440 (97%)	13 (3%)	1 (0%)	43 36

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	404	ARG

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	375/379 (99%)	364 (97%)	11 (3%)	37 29

All (11) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	31	ARG
1	A	106	ASP
1	A	110	LEU
1	A	171	LEU
1	A	229	LEU
1	A	233	ASP
1	A	250	LEU
1	A	274	LEU
1	A	352	MET
1	A	425	SER
1	A	452	GLN

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (7) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	8	GLN
1	A	29	HIS
1	A	40	GLN
1	A	108	GLN
1	A	215	GLN
1	A	281	GLN
1	A	282	GLN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no oligosaccharides in this entry.

5.6 Ligand geometry ⓘ

2 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
3	NAP	A	802	-	50,52,52	2.33	12 (24%)	71,80,80	2.35	20 (28%)
2	FAD	A	801	-	58,58,58	1.70	11 (18%)	85,89,89	1.38	11 (12%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	NAP	A	802	-	1/1/12/12	4/35/67/67	0/5/5/5
2	FAD	A	801	-	-	2/34/50/50	0/6/6/6

All (23) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	A	802	NAP	P2B-O2B	-8.83	1.44	1.59

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	A	802	NAP	C3N-C7N	5.54	1.58	1.50
2	A	801	FAD	P-O3P	-5.38	1.53	1.59
3	A	802	NAP	PA-O3	5.27	1.65	1.59
3	A	802	NAP	P2B-O3X	-4.84	1.36	1.54
3	A	802	NAP	P2B-O2X	4.72	1.72	1.54
3	A	802	NAP	P2B-O1X	4.20	1.63	1.50
2	A	801	FAD	C1'-C2'	3.93	1.58	1.52
2	A	801	FAD	PA-O2A	-3.47	1.39	1.55
2	A	801	FAD	O5'-C5'	3.31	1.57	1.44
3	A	802	NAP	C5N-C4N	3.16	1.44	1.38
2	A	801	FAD	C5'-C4'	2.93	1.55	1.51
2	A	801	FAD	P-O2P	-2.85	1.42	1.55
2	A	801	FAD	C6-C7	-2.79	1.35	1.39
3	A	802	NAP	O4D-C1D	-2.66	1.37	1.40
2	A	801	FAD	P-O5'	-2.35	1.50	1.59
3	A	802	NAP	C2A-N3A	-2.23	1.30	1.33
2	A	801	FAD	C5X-N5	-2.20	1.35	1.39
3	A	802	NAP	PN-O3	2.17	1.61	1.59
3	A	802	NAP	O2B-C2B	-2.13	1.36	1.44
3	A	802	NAP	C6N-C5N	-2.10	1.34	1.38
2	A	801	FAD	O2-C2	-2.04	1.20	1.24
2	A	801	FAD	C4-N3	-2.04	1.35	1.38

All (31) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	A	802	NAP	C5N-C4N-C3N	-8.58	111.93	120.36
3	A	802	NAP	C6N-C5N-C4N	6.11	128.26	119.45
3	A	802	NAP	O4B-C1B-N9A	5.42	118.50	108.09
3	A	802	NAP	O2X-P2B-O2B	4.87	124.83	105.85
3	A	802	NAP	P2B-O2B-C2B	-4.69	110.90	123.43
3	A	802	NAP	O3X-P2B-O2B	4.64	123.93	105.85
3	A	802	NAP	O2X-P2B-O1X	-4.53	93.18	110.83
2	A	801	FAD	C5'-C4'-C3'	-4.25	104.21	112.22
3	A	802	NAP	PN-O5D-C5D	4.04	144.51	121.35
3	A	802	NAP	C5N-C6N-N1N	-3.76	115.25	120.38
3	A	802	NAP	C4D-O4D-C1D	-3.71	106.53	109.92
3	A	802	NAP	O3X-P2B-O2X	-3.67	94.05	107.80
3	A	802	NAP	C4N-C3N-C7N	-3.35	111.95	121.06
3	A	802	NAP	O2B-C2B-C1B	-2.93	99.73	110.05
3	A	802	NAP	C2N-C3N-C7N	2.68	127.19	119.46
2	A	801	FAD	O4-C4-C4X	-2.57	119.75	126.53

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	A	802	NAP	O4B-C4B-C5B	2.55	117.51	109.33
3	A	802	NAP	O7N-C7N-C3N	2.46	122.61	119.60
2	A	801	FAD	C9-C9A-N10	2.42	125.11	121.85
3	A	802	NAP	O5D-C5D-C4D	2.40	117.17	108.99
2	A	801	FAD	C2A-N1A-C6A	2.37	122.63	118.73
2	A	801	FAD	O3B-C3B-C2B	2.24	119.00	111.82
2	A	801	FAD	N6A-C6A-N1A	2.24	123.36	118.38
3	A	802	NAP	C2N-C3N-C4N	2.23	120.85	118.26
2	A	801	FAD	C8M-C8-C9	2.22	123.49	119.57
2	A	801	FAD	O4B-C1B-N9A	-2.18	103.89	108.09
2	A	801	FAD	C4'-C3'-C2'	-2.12	110.04	113.57
2	A	801	FAD	O3'-C3'-C4'	-2.08	104.20	108.93
3	A	802	NAP	C3N-C7N-N7N	-2.05	115.22	117.74
2	A	801	FAD	C4B-O4B-C1B	-2.04	104.96	109.47
3	A	802	NAP	O5B-C5B-C4B	2.00	115.81	108.99

All (1) chirality outliers are listed below:

Mol	Chain	Res	Type	Atom
3	A	802	NAP	C4B

All (6) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
3	A	802	NAP	C5B-O5B-PA-O2A
3	A	802	NAP	C5B-O5B-PA-O3
2	A	801	FAD	C2B-C1B-N9A-C8A
3	A	802	NAP	O4B-C4B-C5B-O5B
2	A	801	FAD	O3'-C3'-C4'-C5'
3	A	802	NAP	C4B-C5B-O5B-PA


There are no ring outliers.


1 monomer is involved in 3 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	A	802	NAP	3	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier.

Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.





5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2		OWAB(Å ²)	Q<0.9
1	A	455/460 (98%)	-0.10	17 (3%)	45 52	13, 23, 44, 69	1 (0%)

All (17) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	331	TYR	5.6
1	A	398	HIS	4.2
1	A	401	SER	4.0
1	A	400	PRO	4.0
1	A	399	LEU	3.0
1	A	403	PRO	3.0
1	A	440	ALA	2.7
1	A	442	GLY	2.5
1	A	346	LEU	2.5
1	A	234	ARG	2.5
1	A	339	SER	2.4
1	A	373	THR	2.4
1	A	236	LYS	2.2
1	A	231	LEU	2.2
1	A	237	GLU	2.1
1	A	235	ILE	2.1
1	A	233	ASP	2.1

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)


There are no oligosaccharides in this entry.


6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q<0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
3	NAP	A	802	48/48	0.88	0.16	25,42,64,66	0
2	FAD	A	801	53/53	0.98	0.04	13,17,22,25	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.





6.5 Other polymers [i](#)

There are no such residues in this entry.