



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 17, 2026 – 10:54 PM UTC

PDB ID : 1E1M / pdb_00001e1m
Title : ADRENODOXIN REDUCTASE in complex with NADPH obtained by a soaking experiment
Authors : Ziegler, G.A.; Schulz, G.E.
Deposited on : 2000-05-09
Resolution : 1.85 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Buster-report	:	wwPDB partial adaption of 1.1.7 (2018)
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.85 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	3428 (1.86-1.86)
Clashscore	190562	3579 (1.86-1.86)
Ramachandran outliers	187476	3553 (1.86-1.86)
Sidechain outliers	187428	3553 (1.86-1.86)
RSRZ outliers	180081	3429 (1.86-1.86)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	460	

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
3	NAP	A	802	X	-	-	-



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
3	A	1	Total 48	C 21	N 7	O 17	P 3	0	0


- Molecule 4 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
4	A	393	Total O 393 393	0	0

3 Residue-property plots

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: ADRENODOXIN REDUCTASE



4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	61.50Å 62.50Å 78.60Å 90.00° 106.70° 90.00°	Depositor
Resolution (Å)	18.00 – 1.85 18.00 – 1.85	Depositor EDS
% Data completeness (in resolution range)	96.3 (18.00-1.85) 96.3 (18.00-1.85)	Depositor EDS
R_{merge}	0.05	Depositor
R_{sym}	0.04	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.90 (at 1.85Å)	Xtriage
Refinement program	REFMAC	Depositor
R, R_{free}	0.188 , 0.228 0.191 , 0.226	Depositor DCC
R_{free} test set	2381 reflections (5.05%)	wwPDB-VP
Wilson B-factor (Å ²)	17.2	Xtriage
Anisotropy	0.316	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.40 , 49.6	EDS
L-test for twinning ²	$\langle L \rangle = 0.48$, $\langle L^2 \rangle = 0.31$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.94	EDS
Total number of atoms	3999	wwPDB-VP
Average B, all atoms (Å ²)	23.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 6.83% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: FAD, NAP

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.94	2/3584 (0.1%)	1.47	27/4871 (0.6%)

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	28	HIS	ND1-CE1	5.55	1.38	1.32
1	A	67	GLN	CD-NE2	-5.38	1.22	1.33

All (27) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	348	VAL	CA-C-N	-9.28	115.91	122.59
1	A	348	VAL	C-N-CA	-9.28	115.91	122.59
1	A	403	PRO	N-CA-C	8.26	129.48	112.47
1	A	415	ASP	CA-CB-CG	7.68	120.28	112.60
1	A	28	HIS	CB-CG-CD2	-7.06	122.02	131.20
1	A	195	VAL	N-CA-C	7.01	118.38	108.36
1	A	278	ARG	NE-CZ-NH2	6.95	125.45	119.20
1	A	402	GLY	CA-C-N	6.84	128.39	119.84
1	A	402	GLY	C-N-CA	6.84	128.39	119.84
1	A	188	ARG	CD-NE-CZ	6.70	133.78	124.40
1	A	207	ILE	N-CA-C	6.43	117.22	110.72
1	A	414	LEU	O-C-N	6.43	129.49	122.15
1	A	140	ALA	O-C-N	-6.36	118.60	121.53
1	A	414	LEU	CA-C-O	-6.23	113.82	120.42
1	A	259	VAL	N-CA-C	5.94	116.59	110.36
1	A	337	ASP	N-CA-CB	-5.90	102.57	110.77
1	A	188	ARG	NE-CZ-NH1	5.89	127.39	121.50
1	A	28	HIS	CB-CG-ND1	5.83	131.44	122.70
1	A	49	PHE	N-CA-CB	-5.39	102.49	110.53

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	70	ARG	CD-NE-CZ	5.38	131.93	124.40
1	A	374	GLY	N-CA-C	5.35	121.53	113.02
1	A	106	ASP	N-CA-C	5.34	117.45	108.96
1	A	419	VAL	CA-C-O	-5.30	114.82	120.59
1	A	195	VAL	N-CA-CB	-5.25	104.69	110.99
1	A	333	SER	N-CA-C	-5.18	103.56	110.55
1	A	39	LYS	N-CA-CB	-5.15	102.55	110.16
1	A	276	PHE	CA-CB-CG	-5.00	108.80	113.80

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3505	0	3535	41	0
2	A	53	0	31	1	0
3	A	48	0	24	5	0
4	A	393	0	0	11	0
All	All	3999	0	3590	44	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 6.

All (44) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:124:ARG:HD3	4:A:2154:HOH:O	1.79	0.81
1:A:160:VAL:HG11	1:A:326:LEU:CD1	2.17	0.73
1:A:306:GLY:O	1:A:309:THR:HG23	1.92	0.70
1:A:309:THR:C	1:A:310:ARG:HD2	2.17	0.69
1:A:29:HIS:HD2	1:A:31:ARG:H	1.41	0.68
1:A:375:VAL:HG12	3:A:802:NAP:H2N	1.76	0.68
1:A:160:VAL:HG11	1:A:326:LEU:HD13	1.80	0.63
1:A:104:ALA:O	1:A:124:ARG:NH2	2.32	0.63

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:215:GLN:HG2	4:A:2232:HOH:O	2.02	0.58
1:A:396:ALA:HB3	1:A:398:HIS:CE1	2.38	0.58
1:A:209:GLU:OE2	3:A:802:NAP:N7N	2.36	0.58
1:A:396:ALA:CB	1:A:398:HIS:CE1	2.90	0.54
1:A:160:VAL:HG11	1:A:326:LEU:HD12	1.89	0.54
1:A:309:THR:O	1:A:310:ARG:HD2	2.08	0.53
1:A:31:ARG:HH11	1:A:31:ARG:HB3	1.74	0.53
1:A:308:ALA:O	1:A:310:ARG:HD3	2.08	0.53
1:A:400:PRO:HD3	4:A:2343:HOH:O	2.10	0.51
1:A:166:THR:HG21	1:A:171:LEU:HD13	1.92	0.51
1:A:450:ASP:O	1:A:454:MET:HG3	2.12	0.50
1:A:433:GLU:HG2	1:A:457:LEU:HD13	1.95	0.49
1:A:124:ARG:CD	4:A:2154:HOH:O	2.49	0.49
1:A:104:ALA:HA	1:A:333:SER:HA	1.96	0.47
1:A:28:HIS:HE1	4:A:2023:HOH:O	1.97	0.46
1:A:55:HIS:N	1:A:56:PRO:CD	2.79	0.45
1:A:201:LEU:HD21	1:A:231:LEU:HD11	1.99	0.45
1:A:77:TYR:OH	1:A:411:LYS:HE2	2.17	0.45
3:A:802:NAP:O2N	4:A:2387:HOH:O	2.20	0.44
3:A:802:NAP:H4N	4:A:2184:HOH:O	2.17	0.44
1:A:330:GLY:C	1:A:331:TYR:CG	2.96	0.43
1:A:185:ARG:HD2	4:A:2205:HOH:O	2.18	0.43
1:A:229:LEU:O	1:A:229:LEU:HD23	2.19	0.43
1:A:396:ALA:HB3	1:A:398:HIS:ND1	2.34	0.42
1:A:404:ARG:HA	4:A:2346:HOH:O	2.19	0.42
1:A:251:ARG:HG2	1:A:255:GLU:OE2	2.20	0.42
1:A:310:ARG:HD2	1:A:310:ARG:N	2.33	0.42
1:A:411:LYS:HE3	4:A:2348:HOH:O	2.20	0.41
1:A:117:LEU:HA	1:A:118:PRO:HD3	1.90	0.41
1:A:216:LEU:HA	1:A:217:PRO:HD3	1.93	0.41
1:A:198:ARG:NE	3:A:802:NAP:O2X	2.40	0.41
1:A:54:ASP:C	1:A:56:PRO:CD	2.94	0.41
1:A:396:ALA:HB1	1:A:398:HIS:CE1	2.56	0.41
2:A:801:FAD:HM83	2:A:801:FAD:HM71	1.77	0.40
1:A:6:THR:HG23	1:A:6:THR:O	2.21	0.40
1:A:259:VAL:HG23	4:A:2265:HOH:O	2.22	0.40

There are no symmetry-related clashes.

5.3 Torsion angles

5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	453/460 (98%)	440 (97%)	11 (2%)	2 (0%)	30 17

All (2) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	404	ARG
1	A	403	PRO

5.3.2 Protein sidechains

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	374/379 (99%)	364 (97%)	10 (3%)	39 24

All (10) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	31	ARG
1	A	110	LEU
1	A	171	LEU
1	A	229	LEU
1	A	233	ASP
1	A	250	LEU
1	A	274	LEU
1	A	339	SER
1	A	352	MET

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	A	452	GLN

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (4) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	28	HIS
1	A	29	HIS
1	A	40	GLN
1	A	398	HIS

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

2 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
3	NAP	A	802	-	50,52,52	2.18	12 (24%)	71,80,80	2.28	22 (30%)
2	FAD	A	801	-	58,58,58	2.07	21 (36%)	85,89,89	1.44	13 (15%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	NAP	A	802	-	1/1/12/12	4/35/67/67	0/5/5/5
2	FAD	A	801	-	-	3/34/50/50	0/6/6/6

All (33) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	A	802	NAP	P2B-O2B	-8.63	1.44	1.59
2	A	801	FAD	P-O3P	-7.89	1.51	1.59
2	A	801	FAD	C6-C7	-4.72	1.33	1.39
3	A	802	NAP	PA-O3	4.69	1.64	1.59
3	A	802	NAP	C5N-C4N	4.55	1.46	1.38
3	A	802	NAP	P2B-O2X	4.31	1.70	1.54
2	A	801	FAD	C5'-C4'	4.09	1.57	1.51
2	A	801	FAD	PA-O2A	-3.60	1.38	1.55
3	A	802	NAP	C4N-C3N	3.57	1.44	1.39
2	A	801	FAD	C1'-C2'	3.52	1.57	1.52
3	A	802	NAP	C6N-C5N	-3.35	1.31	1.38
3	A	802	NAP	P2B-O3X	-3.34	1.42	1.54
2	A	801	FAD	O5'-C5'	3.00	1.55	1.44
2	A	801	FAD	C10-N10	-2.85	1.31	1.37
2	A	801	FAD	C2-N1	-2.81	1.30	1.36
2	A	801	FAD	PA-O3P	2.68	1.62	1.59
3	A	802	NAP	O4D-C1D	-2.54	1.37	1.40
3	A	802	NAP	O2B-C2B	-2.49	1.35	1.44
2	A	801	FAD	P-O2P	-2.47	1.43	1.55
2	A	801	FAD	C5X-N5	-2.46	1.34	1.39
2	A	801	FAD	O4B-C1B	2.40	1.47	1.42
3	A	802	NAP	C3N-C7N	2.36	1.54	1.50
2	A	801	FAD	C8-C7	2.31	1.46	1.40
2	A	801	FAD	C2A-N3A	2.27	1.38	1.33
2	A	801	FAD	O2-C2	-2.25	1.19	1.24
3	A	802	NAP	P2B-O1X	2.21	1.57	1.50
2	A	801	FAD	C4-N3	-2.13	1.34	1.38
2	A	801	FAD	C4A-N9A	2.11	1.42	1.37
2	A	801	FAD	C5B-C4B	2.08	1.57	1.51
2	A	801	FAD	C2-N3	2.06	1.43	1.39
3	A	802	NAP	O3D-C3D	-2.02	1.38	1.43
2	A	801	FAD	C2A-N1A	2.02	1.37	1.33

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	A	801	FAD	C5A-C6A	2.01	1.46	1.41

All (35) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	A	802	NAP	C5N-C4N-C3N	-8.31	112.19	120.36
3	A	802	NAP	O2X-P2B-O2B	5.85	128.66	105.85
3	A	802	NAP	C6N-C5N-C4N	5.23	126.98	119.45
3	A	802	NAP	O4B-C4B-C5B	4.50	123.74	109.33
3	A	802	NAP	O3X-P2B-O2X	-4.16	92.19	107.80
3	A	802	NAP	P2B-O2B-C2B	-3.98	112.81	123.43
3	A	802	NAP	O4B-C1B-N9A	3.77	115.33	108.09
3	A	802	NAP	O2X-P2B-O1X	-3.61	96.77	110.83
3	A	802	NAP	PN-O5D-C5D	3.54	141.65	121.35
3	A	802	NAP	C4N-C3N-C7N	-3.41	111.78	121.06
2	A	801	FAD	C7M-C7-C6	3.37	125.50	119.57
3	A	802	NAP	O2B-C2B-C3B	-3.26	99.98	111.68
3	A	802	NAP	C2N-C3N-C7N	3.12	128.47	119.46
3	A	802	NAP	O3D-C3D-C2D	3.04	121.56	111.82
3	A	802	NAP	C5B-C4B-C3B	3.00	126.00	115.21
2	A	801	FAD	N3A-C2A-N1A	-3.00	124.05	128.58
2	A	801	FAD	C4-C4X-N5	2.95	122.28	118.21
2	A	801	FAD	C5'-C4'-C3'	-2.86	106.82	112.22
2	A	801	FAD	O3'-C3'-C4'	-2.79	102.60	108.93
3	A	802	NAP	C5N-C6N-N1N	-2.68	116.73	120.38
3	A	802	NAP	O5B-C5B-C4B	2.65	118.03	108.99
2	A	801	FAD	O2-C2-N1	2.65	126.20	121.80
3	A	802	NAP	C2A-N1A-C6A	2.57	122.94	118.73
2	A	801	FAD	C7M-C7-C8	-2.54	115.58	120.76
3	A	802	NAP	O5D-C5D-C4D	2.40	117.17	108.99
2	A	801	FAD	C10-N1-C2	2.38	122.00	116.85
2	A	801	FAD	O3B-C3B-C2B	2.36	119.37	111.82
2	A	801	FAD	O2-C2-N3	-2.29	114.19	118.58
2	A	801	FAD	O4B-C1B-C2B	-2.23	101.85	106.62
2	A	801	FAD	C4X-C4-N3	2.16	118.76	113.25
3	A	802	NAP	C3N-C7N-N7N	-2.11	115.14	117.74
2	A	801	FAD	C2A-N1A-C6A	2.09	122.16	118.73
3	A	802	NAP	C2B-C1B-N9A	-2.07	110.35	113.75
3	A	802	NAP	C5D-C4D-C3D	-2.04	107.87	115.21
3	A	802	NAP	O2N-PN-O5D	2.01	116.67	107.57

All (1) chirality outliers are listed below:

Mol	Chain	Res	Type	Atom
3	A	802	NAP	C4B

All (7) torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
3	A	802	NAP	O4B-C4B-C5B-O5B
2	A	801	FAD	O3'-C3'-C4'-C5'
3	A	802	NAP	C3B-C4B-C5B-O5B
2	A	801	FAD	O3'-C3'-C4'-O4'
2	A	801	FAD	C2B-C1B-N9A-C8A
3	A	802	NAP	C3D-C4D-C5D-O5D
3	A	802	NAP	O4D-C4D-C5D-O5D


There are no ring outliers.

2 monomers are involved in 6 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	A	802	NAP	5	0
2	A	801	FAD	1	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.





5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	455/460 (98%)	0.14	31 (6%) 23 25	11, 20, 41, 64	0

All (31) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	398	HIS	4.9
1	A	106	ASP	3.7
1	A	234	ARG	3.5
1	A	396	ALA	3.4
1	A	442	GLY	3.3
1	A	401	SER	3.2
1	A	373	THR	3.1
1	A	231	LEU	3.1
1	A	255	GLU	2.9
1	A	403	PRO	2.9
1	A	400	PRO	2.8
1	A	374	GLY	2.8
1	A	331	TYR	2.6
1	A	440	ALA	2.6
1	A	236	LYS	2.5
1	A	402	GLY	2.5
1	A	342	PHE	2.4
1	A	415	ASP	2.4
1	A	412	ALA	2.4
1	A	416	SER	2.4
1	A	235	ILE	2.3
1	A	305	ILE	2.3
1	A	346	LEU	2.3
1	A	339	SER	2.3
1	A	397	GLY	2.2
1	A	232	GLN	2.2
1	A	229	LEU	2.2

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	124	ARG	2.2
1	A	441	SER	2.2
1	A	343	ASP	2.1
1	A	237	GLU	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)




In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
3	NAP	A	802	48/48	0.83	0.19	21,52,69,71	0
2	FAD	A	801	53/53	0.98	0.05	10,13,22,24	0




The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.

Electron density around NAP A 802:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)

**Electron density around FAD A 801:**

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



6.5 Other polymers [i](#)

There are no such residues in this entry.