



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 6, 2026 – 05:21 AM UTC

PDB ID : 5E1E / pdb_00005e1e
Title : Human JAK1 kinase in complex with compound 30 at 2.30 Angstroms resolution
Authors : Ferguson, A.D.
Deposited on : 2015-09-29
Resolution : 2.30 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Buster-report	:	wwPDB partial adaption of 1.1.7 (2018)
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.30 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	6319 (2.30-2.30)
Clashscore	190562	6919 (2.30-2.30)
Ramachandran outliers	187476	6854 (2.30-2.30)
Sidechain outliers	187428	6854 (2.30-2.30)
RSRZ outliers	180081	6325 (2.30-2.30)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	290	7% 87% 9% ..
1	B	290	5% 88% 8% .

2 Entry composition [i](#)


There are 4 unique types of molecules in this entry. The entry contains 4944 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Tyrosine-protein kinase JAK1.


Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
1	A	281	Total	C	N	O	P	S	0	4	0
			2333	1485	400	430	2	16			
1	B	282	Total	C	N	O	P	S	0	2	0
			2319	1475	398	429	2	15			

- Molecule 2 is DI(HYDROXYETHYL)ETHER (CCD ID: PEG) (formula: C₄H₁₀O₃).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
2	A	1	Total	C	O	0	0
			7	4	3		

- Molecule 3 is 6-chloro-2-(2-fluoro-4,5-dimethoxyphenyl)-N-(piperidin-4-ylmethyl)-3H-imidazo[4,5-b]pyridin-7-amine (CCD ID: 5JG) (formula: C₂₀H₂₃ClFN₅O₂).



Mol	Chain	Residues	Atoms						ZeroOcc	AltConf
3	A	1	Total	C	Cl	F	N	O	0	0
			29	20	1	1	5	2		
3	B	1	Total	C	Cl	F	N	O	0	0
			29	20	1	1	5	2		


- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	117	Total	O	0	0
			117	117		
4	B	110	Total	O	0	0
			110	110		


3 Residue-property plots

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Tyrosine-protein kinase JAK1



- Molecule 1: Tyrosine-protein kinase JAK1



4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	43.09Å 88.71Å 174.93Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	48.73 – 2.30 48.73 – 2.30	Depositor EDS
% Data completeness (in resolution range)	99.8 (48.73-2.30) 99.8 (48.73-2.30)	Depositor EDS
R_{merge}	0.11	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.75 (at 2.29Å)	Xtriage
Refinement program	BUSTER 2.11.6	Depositor
R, R_{free}	0.225 , 0.268 0.219 , 0.267	Depositor DCC
R_{free} test set	1312 reflections (4.27%)	wwPDB-VP
Wilson B-factor (Å ²)	35.3	Xtriage
Anisotropy	0.217	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.30 , 45.8	EDS
L-test for twinning ²	$\langle L \rangle = 0.54$, $\langle L^2 \rangle = 0.38$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.95	EDS
Total number of atoms	4944	wwPDB-VP
Average B, all atoms (Å ²)	43.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The analyses of the Patterson function reveals a significant off-origin peak that is 56.42 % of the origin peak, indicating pseudo-translational symmetry. The chance of finding a peak of this or larger height randomly in a structure without pseudo-translational symmetry is equal to 2.7430e-05. The detected translational NCS is most likely also responsible for the elevated intensity ratio.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: 5JG, PTR, PEG

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# $ Z > 5$	RMSZ	# $ Z > 5$
1	A	0.87	0/2348	1.27	5/3157 (0.2%)
1	B	0.86	0/2336	1.28	8/3141 (0.3%)
All	All	0.87	0/4684	1.28	13/6298 (0.2%)

There are no bond length outliers.

All (13) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	1127	LEU	CA-C-N	6.03	128.37	120.28
1	B	1127	LEU	C-N-CA	6.03	128.37	120.28
1	A	1127	LEU	CA-C-N	5.83	128.10	120.28
1	A	1127	LEU	C-N-CA	5.83	128.10	120.28
1	B	1019	ILE	N-CA-C	-5.54	102.38	109.30
1	B	1063	ASP	CA-C-N	5.29	127.23	120.56
1	B	1063	ASP	C-N-CA	5.29	127.23	120.56
1	A	1063	ASP	CA-C-N	5.25	127.18	120.56
1	A	1063	ASP	C-N-CA	5.25	127.18	120.56
1	A	895	ASP	CA-CB-CG	5.15	117.75	112.60
1	B	919	ILE	N-CA-CB	5.08	117.45	110.54
1	B	1044	PRO	CA-C-N	5.03	127.62	120.53
1	B	1044	PRO	C-N-CA	5.03	127.62	120.53

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen

atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2333	0	2325	11	0
1	B	2319	0	2316	9	0
2	A	7	0	10	0	0
3	A	29	0	23	4	0
3	B	29	0	23	3	0
4	A	117	0	0	0	0
4	B	110	0	0	0	0
All	All	4944	0	4697	23	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 2.

All (23) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
3:B:1201:5JG:N16	3:B:1201:5JG:H9	1.91	0.83
3:A:1202:5JG:H9	3:A:1202:5JG:N16	2.02	0.73
1:B:957:GLU:O	3:B:1201:5JG:H8	2.01	0.60
1:A:957:GLU:O	3:A:1202:5JG:H8	2.02	0.60
1:A:910:LEU:HD11	1:A:919:ILE:HA	1.85	0.59
1:B:909:SER:HB3	1:B:953:LYS:HG2	1.87	0.55
1:B:1029:GLU:HB3	1:B:1032:LYS:HG3	1.90	0.52
1:A:1029:GLU:HB3	1:A:1032:LYS:HG3	1.90	0.52
3:B:1201:5JG:N16	3:B:1201:5JG:C22	2.72	0.48
1:A:1000:VAL:HG13	1:A:1028:ILE:HD11	1.95	0.48
1:B:1000:VAL:HG13	1:B:1028:ILE:HD11	1.95	0.48
1:B:944:CYS:HB3	1:B:953:LYS:HB2	1.97	0.47
1:A:944:CYS:HB3	1:A:953:LYS:HB2	1.98	0.46
1:A:1044:PRO:HD2	1:A:1048:TYR:CZ	2.53	0.43
1:A:1010:LEU:HD21	3:A:1202:5JG:H24	1.99	0.43
1:B:910:LEU:HD11	1:B:919:ILE:HA	2.01	0.43
1:A:1069:VAL:O	1:A:1073:GLU:HG3	2.19	0.42
1:B:1069:VAL:O	1:B:1073:GLU:HG3	2.18	0.42
1:B:926:ILE:HG12	1:B:954:LEU:HB2	2.02	0.41
1:B:1044:PRO:HD2	1:B:1048:TYR:CZ	2.55	0.41
1:A:926:ILE:HG12	1:A:954:LEU:HB2	2.02	0.41
1:A:1048:TYR:CZ	1:A:1058:PHE:HZ	2.39	0.41
1:A:1021:ASP:HB2	3:A:1202:5JG:H16	2.02	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	277/290 (96%)	269 (97%)	7 (2%)	1 (0%)	30	38
1	B	276/290 (95%)	269 (98%)	7 (2%)	0	100	100
All	All	553/580 (95%)	538 (97%)	14 (2%)	1 (0%)	43	55

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	1021	ASP

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	256/257 (100%)	250 (98%)	6 (2%)	44	63
1	B	255/257 (99%)	249 (98%)	6 (2%)	43	62
All	All	511/514 (99%)	499 (98%)	12 (2%)	43	63

All (12) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	872	LYS
1	A	927	GLU

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	A	1029	GLU
1	A	1037	VAL
1	A	1102	THR
1	A	1129	ARG
1	B	871	GLU
1	B	919	ILE
1	B	933	TYR
1	B	1037	VAL
1	B	1102	THR
1	B	1144	ASN

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (6) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	950	ASN
1	A	1016	GLN
1	B	885	HIS
1	B	950	ASN
1	B	979	GLN
1	B	1055	GLN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

4 non-standard protein/DNA/RNA residues are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
1	PTR	B	1035	1	15,16,17	1.40	2 (13%)	17,22,24	1.20	1 (5%)
1	PTR	A	1034	1	15,16,17	1.47	3 (20%)	17,22,24	1.18	1 (5%)

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
1	PTR	B	1034	1	15,16,17	1.39	3 (20%)	17,22,24	1.06	1 (5%)
1	PTR	A	1035	1	15,16,17	1.43	2 (13%)	17,22,24	1.22	1 (5%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
1	PTR	B	1035	1	-	0/10/11/13	0/1/1/1
1	PTR	A	1034	1	-	0/10/11/13	0/1/1/1
1	PTR	B	1034	1	-	0/10/11/13	0/1/1/1
1	PTR	A	1035	1	-	0/10/11/13	0/1/1/1

All (10) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	1035	PTR	CE2-CZ	2.86	1.44	1.38
1	A	1034	PTR	CE1-CZ	2.74	1.43	1.38
1	A	1034	PTR	P-OH	-2.67	1.54	1.59
1	B	1034	PTR	CE1-CZ	2.56	1.43	1.38
1	B	1034	PTR	CB-CA	2.42	1.58	1.53
1	B	1034	PTR	P-OH	-2.42	1.55	1.59
1	B	1035	PTR	CE1-CZ	2.36	1.43	1.38
1	A	1034	PTR	CB-CA	2.25	1.58	1.53
1	A	1035	PTR	CE1-CD1	2.15	1.42	1.38
1	B	1035	PTR	CE2-CZ	2.14	1.42	1.38

All (4) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	1035	PTR	OH-CZ-CE2	2.67	127.23	119.22
1	B	1034	PTR	OH-CZ-CE1	2.67	127.23	119.22
1	A	1034	PTR	OH-CZ-CE1	2.57	126.92	119.22
1	B	1035	PTR	OH-CZ-CE2	2.51	126.73	119.22

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

3 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
3	5JG	A	1202	-	31,32,32	0.31	0	40,45,45	1.13	3 (7%)
3	5JG	B	1201	-	31,32,32	0.33	0	40,45,45	1.25	4 (10%)
2	PEG	A	1201	-	6,6,6	0.25	0	5,5,5	0.15	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	5JG	A	1202	-	-	7/13/21/21	0/4/4/4
3	5JG	B	1201	-	-	3/13/21/21	0/4/4/4
2	PEG	A	1201	-	-	2/4/4/4	-

There are no bond length outliers.

All (7) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	B	1201	5JG	C24-C23-C22	-4.25	102.83	111.47
3	A	1202	5JG	C23-C22-N21	4.18	119.61	112.83
3	B	1201	5JG	C23-C22-N21	3.62	118.70	112.83
3	A	1202	5JG	C24-C23-C22	-2.81	105.76	111.47
3	B	1201	5JG	C27-C28-C23	-2.70	107.96	112.14
3	A	1202	5JG	C27-C28-C23	-2.67	108.00	112.14
3	B	1201	5JG	C28-C23-C22	-2.13	107.14	111.47

There are no chirality outliers.

All (12) torsion outliers are listed below:



Mol	Chain	Res	Type	Atoms
3	A	1202	5JG	N21-C22-C23-C28
3	B	1201	5JG	C23-C22-N21-C17
3	A	1202	5JG	C8-C3-O2-C1
2	A	1201	PEG	C1-C2-O2-C3
2	A	1201	PEG	C4-C3-O2-C2
3	A	1202	5JG	C4-C3-O2-C1
3	A	1202	5JG	C23-C22-N21-C17
3	A	1202	5JG	N21-C22-C23-C24
3	A	1202	5JG	N13-C12-C5-C6
3	B	1201	5JG	N13-C12-C5-C6
3	A	1202	5JG	C15-C17-N21-C22
3	B	1201	5JG	C15-C17-N21-C22

There are no ring outliers.

2 monomers are involved in 7 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	A	1202	5JG	4	0
3	B	1201	5JG	3	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.



5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ > 2		OWAB(Å ²)	Q < 0.9
1	A	279/290 (96%)	0.57	21 (7%)	20 22	15, 40, 75, 97	4 (1%)
1	B	280/290 (96%)	0.48	14 (5%)	34 35	19, 40, 68, 101	2 (0%)
All	All	559/580 (96%)	0.52	35 (6%)	26 28	15, 40, 71, 101	6 (1%)

All (35) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	1099[A]	MET	7.1
1	A	884	GLY	4.9
1	A	1154	LYS	4.7
1	A	865	VAL	4.7
1	A	886	PHE	4.7
1	B	885	HIS	4.5
1	B	1154	LYS	4.4
1	B	865	VAL	3.9
1	A	917	ASN	3.8
1	B	919	ILE	3.6
1	A	933[A]	TYR	3.5
1	A	885	HIS	3.4
1	B	950	ASN	3.4
1	B	917	ASN	3.3
1	A	911	LYS	3.0
1	A	945	THR	2.9
1	B	946	GLU	2.8
1	A	1090	LYS	2.7
1	A	1091	MET	2.5
1	B	933	TYR	2.4
1	B	1031	ASP	2.4
1	A	950	ASN	2.3
1	B	886	PHE	2.3
1	B	884	GLY	2.3

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	1041	ARG	2.3
1	A	883	GLU	2.2
1	B	911	LYS	2.2
1	A	919	ILE	2.2
1	A	1031	ASP	2.1
1	A	869	HIS	2.1
1	A	918	HIS	2.1
1	A	1095	THR	2.1
1	B	871	GLU	2.1
1	A	1058	PHE	2.1
1	B	1127	LEU	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
1	PTR	A	1034	16/17	0.89	0.11	52,60,77,79	0
1	PTR	B	1034	16/17	0.90	0.10	50,58,72,73	0
1	PTR	B	1035	16/17	0.90	0.10	47,50,66,67	0
1	PTR	A	1035	16/17	0.93	0.10	48,54,66,67	0

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)




In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.


Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
2	PEG	A	1201	7/7	0.77	0.21	64,65,66,66	0
3	5JG	A	1202	29/29	0.92	0.09	24,31,36,38	0
3	5JG	B	1201	29/29	0.94	0.08	22,28,35,42	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.

Electron density around 5JG A 1202:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)





6.5 Other polymers [i](#)

There are no such residues in this entry.