



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 8, 2026 – 04:48 PM UTC

PDB ID : 8E15 / pdb_00008e15
Title : A computationally stabilized hMPV F protein
Authors : Huang, J.; Gonzalez, K.; Mousa, J.; Strauch, E.
Deposited on : 2022-08-09
Resolution : 2.41 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0
Mogul : 2022.3.0, CSD as543be (2022)
Xtriage (Phenix) : 2.0
EDS : 3.0
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4 : 9.0.010 (Gargrove)
Density-Fitness : 1.0.12
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.41 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	6062 (2.44-2.40)
Clashscore	190562	6562 (2.44-2.40)
Ramachandran outliers	187476	6481 (2.44-2.40)
Sidechain outliers	187428	6482 (2.44-2.40)
RSRZ outliers	180081	6066 (2.44-2.40)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	F	102	6% 56% 17% 27%
2	G	422	5% 76% 12% 13%
3	A	3	33% 67%

2 Entry composition

There are 5 unique types of molecules in this entry. The entry contains 3438 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called F2 protein.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	F	74	Total	C	N	O	S	0	0	0
			580	361	94	123	2			

There are 3 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
F	90	ASN	ALA	engineered mutation	UNP Q8B9P0
F	100	ARG	GLN	engineered mutation	UNP Q8B9P0
F	101	ARG	SER	engineered mutation	UNP Q8B9P0

- Molecule 2 is a protein called F1 protein with Fibrin peptide.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	G	369	Total	C	N	O	S	0	0	0
			2780	1739	482	540	19			

There are 14 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
G	114	GLU	THR	engineered mutation	UNP Q8B9P0
G	130	ASP	LEU	engineered mutation	UNP Q8B9P0
G	159	LEU	ALA	engineered mutation	UNP Q8B9P0
G	185	PRO	ALA	engineered mutation	UNP Q8B9P0
G	203	ILE	VAL	engineered mutation	UNP Q8B9P0
G	430	GLN	VAL	engineered mutation	UNP Q8B9P0
G	449	ASP	VAL	engineered mutation	UNP Q8B9P0
G	491	SER	-	linker	UNP Q8B9P0
G	519	HIS	-	expression tag	UNP Q76VI8
G	520	HIS	-	expression tag	UNP Q76VI8
G	521	HIS	-	expression tag	UNP Q76VI8
G	522	HIS	-	expression tag	UNP Q76VI8
G	523	HIS	-	expression tag	UNP Q76VI8

Continued on next page...

Continued from previous page...


Chain	Residue	Modelled	Actual	Comment	Reference
G	524	HIS	-	expression tag	UNP Q76VI8

- Molecule 3 is an oligosaccharide called beta-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose.



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
3	A	3	Total	C	N	O	0	0	0
			39	22	2	15			

- Molecule 4 is 2-acetamido-2-deoxy-beta-D-glucopyranose (CCD ID: NAG) (formula: $C_8H_{15}NO_6$).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
4	F	1	Total	C	N	O	0	0
			14	8	1	5		
4	G	1	Total	C	N	O	0	0
			14	8	1	5		


- Molecule 5 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	F	3	Total 3	O 3	0	0
5	G	8	Total 8	O 8	0	0


3 Residue-property plots

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: F2 protein



- Molecule 2: F1 protein with Fibrin peptide



- Molecule 3: beta-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose



4 Data and refinement statistics

Property	Value	Source
Space group	I 21 3	Depositor
Cell constants a, b, c, α , β , γ	178.19Å 178.19Å 178.19Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	47.62 – 2.41 47.62 – 2.41	Depositor EDS
% Data completeness (in resolution range)	99.9 (47.62-2.41) 99.9 (47.62-2.41)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.22 (at 2.42Å)	Xtriage
Refinement program	PHENIX 1.20.1_4487	Depositor
R, R_{free}	0.204 , 0.249 0.204 , 0.249	Depositor DCC
R_{free} test set	1872 reflections (5.15%)	wwPDB-VP
Wilson B-factor (Å ²)	64.9	Xtriage
Anisotropy	0.000	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.35 , 43.9	EDS
L-test for twinning ²	$\langle L \rangle = 0.51$, $\langle L^2 \rangle = 0.34$	Xtriage
Estimated twinning fraction	0.018 for -l,-k,-h	Xtriage
F_o, F_c correlation	0.96	EDS
Total number of atoms	3438	wwPDB-VP
Average B, all atoms (Å ²)	70.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 2.43% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: BMA, NAG

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	F	0.43	0/586	0.60	0/795
2	G	0.39	0/2822	0.58	0/3828
All	All	0.40	0/3408	0.58	0/4623

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	F	580	0	570	19	0
2	G	2780	0	2753	38	0
3	A	39	0	34	0	0
4	F	14	0	13	0	0
4	G	14	0	13	0	0
5	F	3	0	0	0	0
5	G	8	0	0	1	0
All	All	3438	0	3383	45	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 7.

All (45) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:G:233:ASN:OD1	5:G:701:HOH:O	1.91	0.88
2:G:411:THR:HG22	2:G:413:GLN:H	1.43	0.81
1:F:55:VAL:O	1:F:75:LYS:NZ	2.25	0.69
1:F:56:GLU:HA	1:F:75:LYS:NZ	2.08	0.69
1:F:70:GLU:HB3	2:G:193:PHE:CD2	2.29	0.68
2:G:168:PHE:HE2	2:G:193:PHE:HD1	1.45	0.64
1:F:51:GLU:HB3	2:G:166:LYS:HD3	1.78	0.64
2:G:392:ILE:HG22	2:G:418:VAL:HG12	1.80	0.62
1:F:46:ASN:ND2	2:G:158:LEU:HD23	2.16	0.60
1:F:30:THR:HG23	2:G:381:LEU:HD11	1.82	0.59
2:G:396:ARG:HG2	2:G:396:ARG:HH11	1.69	0.56
2:G:323:GLU:H	2:G:323:GLU:CD	2.14	0.55
2:G:222:MET:HE1	2:G:273:LEU:HD11	1.88	0.55
1:F:56:GLU:HA	1:F:75:LYS:HZ2	1.72	0.54
2:G:130:ASP:OD1	2:G:131:GLU:N	2.42	0.53
1:F:46:ASN:HD22	2:G:158:LEU:HB3	1.74	0.52
2:G:294:GLU:HB2	2:G:299:TYR:CE1	2.44	0.52
1:F:70:GLU:HB3	2:G:193:PHE:HD2	1.75	0.52
2:G:173:LEU:O	2:G:177:ILE:HG13	2.10	0.51
2:G:419:THR:HG22	2:G:424:VAL:HA	1.93	0.49
2:G:139:ASN:HA	2:G:142:LYS:HD2	1.94	0.49
1:F:53:GLY:HA2	2:G:166:LYS:HG3	1.93	0.49
2:G:113:ALA:HB2	2:G:339:ALA:HB2	1.96	0.48
2:G:411:THR:HG22	2:G:413:GLN:N	2.19	0.48
2:G:217:ILE:HD11	2:G:255:GLY:HA3	1.97	0.47
1:F:37:SER:HB3	2:G:283:CYS:SG	2.55	0.47
1:F:62:ASP:CG	2:G:182:CYS:H	2.24	0.46
2:G:396:ARG:HG2	2:G:396:ARG:NH1	2.31	0.46
1:F:36:LEU:HB3	2:G:280:ASP:HA	1.97	0.46
1:F:46:ASN:ND2	2:G:158:LEU:HB3	2.31	0.45
1:F:89:LEU:HA	1:F:89:LEU:HD12	1.66	0.45
2:G:446:PHE:O	2:G:448:PRO:HD3	2.18	0.44
2:G:453:GLU:O	2:G:455:GLN:HG2	2.18	0.44
1:F:53:GLY:CA	2:G:166:LYS:HG3	2.48	0.44
1:F:81:LEU:HD12	1:F:81:LEU:HA	1.76	0.43
2:G:296:LYS:O	2:G:296:LYS:HG2	2.19	0.43
2:G:302:LEU:C	2:G:302:LEU:HD12	2.43	0.42
1:F:50:LEU:C	1:F:50:LEU:HD23	2.44	0.42
2:G:359:TYR:CG	2:G:360:PRO:HD2	2.55	0.42
2:G:245:LEU:HD23	2:G:245:LEU:HA	1.88	0.42

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:G:326:CYS:HA	2:G:334:PHE:O	2.20	0.42
1:F:60:CYS:HB2	1:F:68:LYS:HG3	2.02	0.41
2:G:373:VAL:HG23	2:G:420:ILE:HD12	2.03	0.41
2:G:132:SER:HA	2:G:135:THR:OG1	2.22	0.40
2:G:168:PHE:HE2	2:G:193:PHE:CD1	2.32	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	F	72/102 (71%)	68 (94%)	4 (6%)	0	100	100
2	G	367/422 (87%)	359 (98%)	8 (2%)	0	100	100
All	All	439/524 (84%)	427 (97%)	12 (3%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	F	67/94 (71%)	66 (98%)	1 (2%)	57	75
2	G	304/354 (86%)	301 (99%)	3 (1%)	68	82
All	All	371/448 (83%)	367 (99%)	4 (1%)	65	81

All (4) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	F	73	LEU
2	G	355	SER
2	G	427	LEU
2	G	431	GLU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (4) such sidechains are listed below:

Mol	Chain	Res	Type
2	G	178	ASN
2	G	313	ASN
2	G	402	GLN
2	G	455	GLN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

3 monosaccharides are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
3	NAG	A	1	3,2	14,14,15	1.15	2 (14%)	17,19,21	0.62	0
3	NAG	A	2	3	14,14,15	0.65	0	17,19,21	0.47	0
3	BMA	A	3	3	11,11,12	2.14	4 (36%)	15,15,17	1.20	2 (13%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	NAG	A	1	3,2	-	0/6/23/26	0/1/1/1
3	NAG	A	2	3	-	0/6/23/26	0/1/1/1
3	BMA	A	3	3	-	1/2/19/22	0/1/1/1

All (6) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	A	3	BMA	C4-C5	4.04	1.61	1.53
3	A	3	BMA	C4-C3	3.79	1.62	1.52
3	A	1	NAG	O5-C1	3.49	1.49	1.43
3	A	3	BMA	C1-C2	2.32	1.57	1.52
3	A	1	NAG	C1-C2	2.28	1.55	1.52
3	A	3	BMA	O5-C1	2.03	1.47	1.43

All (2) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	A	3	BMA	C1-O5-C5	2.87	116.03	112.19
3	A	3	BMA	C3-C4-C5	2.02	113.89	110.23

There are no chirality outliers.


All (1) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
3	A	3	BMA	O5-C5-C6-O6

There are no ring outliers.

No monomer is involved in short contacts.

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for oligosaccharide.



5.6 Ligand geometry [i](#)

2 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
4	NAG	G	601	2	14,14,15	0.61	1 (7%)	17,19,21	0.43	0
4	NAG	F	601	1	14,14,15	1.71	2 (14%)	17,19,21	1.25	1 (5%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	NAG	G	601	2	-	1/6/23/26	0/1/1/1
4	NAG	F	601	1	-	1/6/23/26	0/1/1/1

All (3) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
4	F	601	NAG	C1-C2	5.59	1.60	1.52
4	F	601	NAG	O5-C1	2.40	1.47	1.43
4	G	601	NAG	O5-C1	2.03	1.47	1.43

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
4	F	601	NAG	C2-N2-C7	3.84	128.04	122.90

There are no chirality outliers.

All (2) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
4	F	601	NAG	C1-C2-N2-C7
4	G	601	NAG	C1-C2-N2-C7

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	F	74/102 (72%)	0.56	6 (8%) 18 14	52, 73, 104, 112	0
2	G	369/422 (87%)	0.33	19 (5%) 33 30	51, 65, 96, 116	0
All	All	443/524 (84%)	0.37	25 (5%) 30 26	51, 65, 100, 116	0

All (25) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	F	91	ARG	3.9
1	F	59	THR	3.5
2	G	471	GLN	3.4
2	G	193	PHE	3.2
2	G	131	GLU	3.1
2	G	184	ILE	2.9
1	F	60	CYS	2.9
2	G	216	ALA	2.8
1	F	87	ASP	2.8
2	G	175	ARG	2.8
2	G	187	LEU	2.8
2	G	345	GLU	2.7
2	G	185	PRO	2.7
2	G	356	THR	2.6
2	G	170	SER	2.5
2	G	295	LYS	2.4
1	F	88	GLN	2.4
2	G	296	LYS	2.3
2	G	323	GLU	2.3
1	F	63	GLY	2.3
2	G	154	GLY	2.2
2	G	372	MET	2.2
2	G	355	SER	2.2
2	G	468	GLU	2.0

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
2	G	181	LYS	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)


There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
3	NAG	A	1	14/15	-	-	96,100,109,113	0
3	NAG	A	2	14/15	-	-	101,113,116,116	0
3	BMA	A	3	11/12	-	-	91,107,112,112	0

The following is a graphical depiction of the model fit to experimental electron density for oligosaccharide. Each fit is shown from different orientation to approximate a three-dimensional view.



6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
4	NAG	F	601	14/15	0.19	0.25	110,127,129,130	0
4	NAG	G	601	14/15	0.84	0.17	96,104,108,108	0

6.5 Other polymers [i](#)

There are no such residues in this entry.