



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 6, 2026 – 04:08 PM UTC

PDB ID : 6E2J / pdb_00006e2j
Title : Crystal structure of the heterocomplex between human keratin 1 coil 1B containing S233L mutation and wild-type human keratin 10 coil 1B
Authors : Eldirany, S.A.; Lomakin, I.B.; Bunick, C.G.
Deposited on : 2018-07-11
Resolution : 2.39 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0
Mogul : 2022.3.0, CSD as543be (2022)
Xtriage (Phenix) : 2.0
EDS : 3.0
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4 : 9.0.010 (Gargrove)
Density-Fitness : 1.0.12
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.39 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	7164 (2.40-2.36)
Clashscore	190562	7722 (2.40-2.36)
Ramachandran outliers	187476	7626 (2.40-2.36)
Sidechain outliers	187428	7627 (2.40-2.36)
RSRZ outliers	180081	7170 (2.40-2.36)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	107	25% 88% 11%
2	B	104	32% 84% 15%

2 Entry composition

There are 4 unique types of molecules in this entry. The entry contains 3560 atoms, of which 1729 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Keratin, type II cytoskeletal 1.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
1	A	107	Total	C	H	N	O	S	0	0	0
			1766	554	872	152	183	5			

There are 2 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	225	MET	-	initiating methionine	UNP P04264
A	233	LEU	SER	engineered mutation	UNP P04264

- Molecule 2 is a protein called Keratin, type I cytoskeletal 10.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
2	B	104	Total	C	H	N	O	S	0	1	0
			1714	531	857	148	176	2			

There are 2 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
B	193	GLY	-	expression tag	UNP P13645
B	194	SER	-	expression tag	UNP P13645

- Molecule 3 is SULFATE ION (CCD ID: SO4) (formula: O₄S).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	A	1	Total	O	S	0	0
			5	4	1		
3	A	1	Total	O	S	0	0
			5	4	1		
3	A	1	Total	O	S	0	0
			5	4	1		
3	B	1	Total	O	S	0	0
			5	4	1		



- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	32	Total	O	0	0
			32	32		
4	B	28	Total	O	0	0
			28	28		



3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Keratin, type II cytoskeletal 1



- Molecule 2: Keratin, type I cytoskeletal 10



4 Data and refinement statistics

Property	Value	Source
Space group	P 64 2 2	Depositor
Cell constants a, b, c, α , β , γ	93.30Å 93.30Å 124.74Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	46.65 – 2.39 46.65 – 2.39	Depositor EDS
% Data completeness (in resolution range)	99.4 (46.65-2.39) 99.5 (46.65-2.39)	Depositor EDS
R_{merge}	0.14	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.92 (at 2.39Å)	Xtriage
Refinement program	PHENIX	Depositor
R, R_{free}	0.271 , 0.294 0.285 , 0.306	Depositor DCC
R_{free} test set	700 reflections (5.24%)	wwPDB-VP
Wilson B-factor (Å ²)	66.9	Xtriage
Anisotropy	0.347	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.39 , 74.0	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.33$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.94	EDS
Total number of atoms	3560	wwPDB-VP
Average B, all atoms (Å ²)	106.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 6.19% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: SO4

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.20	0/903	0.36	0/1212
2	B	0.21	0/866	0.34	0/1165
All	All	0.20	0/1769	0.35	0/2377

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	894	872	872	9	1
2	B	857	857	857	16	0
3	A	15	0	0	1	0
3	B	5	0	0	0	0
4	A	32	0	0	0	0
4	B	28	0	0	1	0
All	All	1831	1729	1729	18	1

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 5.

All (18) close contacts within the same asymmetric unit are listed below, sorted by their clash

magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:283:GLU:OE1	2:B:256:ARG:NH2	2.13	0.81
2:B:232:ASP:O	2:B:235:ARG:HG2	1.98	0.63
1:A:322:GLU:OE1	2:B:287:HIS:NE2	2.26	0.61
1:A:277:ARG:NE	2:B:241:GLU:OE1	2.30	0.58
1:A:240:ARG:NH2	3:A:403:SO4:O3	2.40	0.53
1:A:323:LEU:HD23	2:B:291:MET:HE1	1.91	0.53
2:B:282:TYR:C	2:B:282:TYR:CD1	2.87	0.53
2:B:195:ASP:OD1	2:B:195:ASP:N	2.38	0.52
1:A:323:LEU:HD23	2:B:291:MET:CE	2.40	0.52
2:B:245[B]:ARG:NH1	2:B:249:GLU:OE1	2.47	0.47
2:B:245[B]:ARG:NH2	4:B:401:HOH:O	2.30	0.47
2:B:210:ILE:O	2:B:214:THR:HG23	2.14	0.47
1:A:277:ARG:HD2	2:B:244:LEU:HD13	1.97	0.47
2:B:235:ARG:O	2:B:239:GLU:HG3	2.16	0.45
1:A:259:MET:HE3	2:B:228:ARG:HB2	2.00	0.44
2:B:232:ASP:HA	2:B:235:ARG:HG2	2.00	0.43
2:B:283:LEU:HD23	2:B:283:LEU:C	2.45	0.42
1:A:270:TYR:CZ	1:A:274:ILE:HD11	2.56	0.41

All (1) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:226:ASN:OD1	1:A:331:SER:OG[1_465]	2.05	0.15

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	105/107 (98%)	105 (100%)	0	0	100	100
2	B	103/104 (99%)	102 (99%)	1 (1%)	0	100	100

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
All	All	208/211 (99%)	207 (100%)	1 (0%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	101/101 (100%)	97 (96%)	4 (4%)	28	44
2	B	95/94 (101%)	93 (98%)	2 (2%)	47	67
All	All	196/195 (100%)	190 (97%)	6 (3%)	35	54

All (6) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	233	LEU
1	A	240	ARG
1	A	289	LYS
1	A	315	LEU
2	B	195	ASP
2	B	242	VAL

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (3) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	249	GLN
2	B	209	GLN
2	B	223	GLN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

4 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
3	SO4	A	401	-	4,4,4	0.24	0	6,6,6	0.09	0
3	SO4	A	403	-	4,4,4	0.31	0	6,6,6	0.16	0
3	SO4	A	402	-	4,4,4	0.31	0	6,6,6	0.10	0
3	SO4	B	301	-	4,4,4	0.24	0	6,6,6	0.08	0

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

1 monomer is involved in 1 short contact:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	A	403	SO4	1	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues ⓘ

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	107/107 (100%)	1.38	27 (25%) 1 1	66, 92, 195, 238	0
2	B	104/104 (100%)	1.63	33 (31%) 1 1	61, 96, 186, 300	1 (0%)
All	All	211/211 (100%)	1.50	60 (28%) 1 1	61, 94, 193, 300	1 (0%)

All (60) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
2	B	193	GLY	6.0
2	B	244	LEU	5.9
2	B	236	LEU	5.6
2	B	243	ALA	4.9
2	B	248	VAL	4.9
1	A	330	ILE	4.8
1	A	225	MET	4.4
2	B	196	TYR	4.4
2	B	242	VAL	4.2
1	A	231	PHE	3.8
2	B	194	SER	3.8
1	A	321	ALA	3.7
2	B	245[A]	ARG	3.6
1	A	323	LEU	3.5
1	A	266	TYR	3.4
1	A	230	TYR	3.4
1	A	285	VAL	3.4
1	A	227	LEU	3.2
2	B	235	ARG	3.1
2	B	200	TYR	3.1
2	B	294	LEU	3.0
2	B	199	TYR	3.0
1	A	318	LEU	3.0
2	B	267	ALA	2.9

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	328	THR	2.9
2	B	211	LEU	2.8
1	A	241	ARG	2.8
2	B	202	THR	2.7
2	B	238	TYR	2.7
1	A	319	TYR	2.7
1	A	326	MET	2.6
2	B	247	SER	2.6
1	A	307	ASN	2.5
1	A	317	ALA	2.5
2	B	246	GLN	2.5
2	B	195	ASP	2.5
2	B	239	GLU	2.4
2	B	250	ALA	2.4
2	B	296	ASN	2.4
2	B	203	ILE	2.4
2	B	276	LEU	2.4
1	A	315	LEU	2.4
1	A	267	ARG	2.3
1	A	226	ASN	2.3
1	A	259	MET	2.3
1	A	234	PHE	2.2
1	A	233	LEU	2.2
2	B	251	ASP	2.2
2	B	197	SER	2.2
2	B	252	ILE	2.2
2	B	208	ASN	2.2
2	B	240	ASN	2.1
2	B	241	GLU	2.1
1	A	331	SER	2.1
1	A	273	GLU	2.1
2	B	280	LEU	2.1
2	B	224	ILE	2.1
1	A	324	SER	2.0
1	A	322	GLU	2.0
1	A	327	GLN	2.0

6.2 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
3	SO4	A	403	5/5	0.38	0.18	193,194,195,196	0
3	SO4	A	402	5/5	0.48	0.18	190,190,192,192	0
3	SO4	B	301	5/5	0.80	0.11	130,130,132,132	0
3	SO4	A	401	5/5	0.88	0.20	137,138,138,140	0

6.5 Other polymers [i](#)

There are no such residues in this entry.