



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 8, 2026 – 02:51 PM UTC

PDB ID : 3E30 / pdb_00003e30
Title : Protein farnesyltransferase complexed with FPP and ethylene diamine inhibitor 4
Authors : Hast, M.A.; Beese, L.S.
Deposited on : 2008-08-06
Resolution : 2.45 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Buster-report	:	wwPDB partial adaption of 1.1.7 (2018)
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.45 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
Clashscore	190562	1229 (2.46-2.46)
Ramachandran outliers	187476	1218 (2.46-2.46)
Sidechain outliers	187428	1218 (2.46-2.46)
RSRZ outliers	180081	1190 (2.46-2.46)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	377	2% 75% 10% 14%
2	B	437	2% 84% 10% 7%

2 Entry composition [i](#)

There are 6 unique types of molecules in this entry. The entry contains 6450 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Protein farnesyltransferase/geranylgeranyltransferase type-1 subunit alpha.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	323	Total	C	N	O	S	0	0	0
			2745	1746	481	513	5			


- Molecule 2 is a protein called Protein farnesyltransferase subunit beta.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	B	407	Total	C	N	O	S	0	0	0
			3203	2050	550	580	23			

- Molecule 3 is ZINC ION (CCD ID: ZN) (formula: Zn).


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	B	1	Total	Zn	0	0
			1	1		

- Molecule 4 is FARNESYL DIPHOSPHATE (CCD ID: FPP) (formula: C₁₅H₂₈O₇P₂).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
4	B	1	Total	C	O	P	0	0
			24	15	7	2		

- Molecule 5 is tert-butyl 4-((2-((4-cyanophenyl)[(1-methyl-1H-imidazol-5-yl)methyl]amino)ethyl)[(1-methyl-1H-imidazol-4-yl)sulfonyl]amino}methyl)piperidine-1-carboxylate (CCD ID: ED4) (formula: C₂₉H₄₀N₈O₄S).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
5	B	1	Total	C	N	O	S	0	0
			42	29	8	4	1		

- Molecule 6 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
6	A	217	Total O 217 217	0	0
6	B	218	Total O 218 218	0	0

4 Data and refinement statistics

Property	Value	Source
Space group	P 61	Depositor
Cell constants a, b, c, α , β , γ	173.31Å 173.31Å 70.15Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	40.72 – 2.45 40.72 – 2.45	Depositor EDS
% Data completeness (in resolution range)	97.6 (40.72-2.45) 97.5 (40.72-2.45)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.34 (at 2.45Å)	Xtriage
Refinement program	REFMAC 5.4.0077	Depositor
R, R_{free}	0.184 , 0.222 0.185 , (Not available)	Depositor DCC
R_{free} test set	No test flags present.	wwPDB-VP
Wilson B-factor (Å ²)	35.0	Xtriage
Anisotropy	0.096	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.35 , 44.1	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.33$	Xtriage
Estimated twinning fraction	0.031 for h,-h-k,-l	Xtriage
F_o, F_c correlation	0.95	EDS
Total number of atoms	6450	wwPDB-VP
Average B, all atoms (Å ²)	35.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.46% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: ED4, ZN, FPP

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.46	0/2812	0.77	7/3817 (0.2%)
2	B	0.47	0/3292	0.76	0/4473
All	All	0.47	0/6104	0.76	7/8290 (0.1%)

There are no bond length outliers.

All (7) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	185	GLU	N-CA-C	5.64	117.23	111.14
1	A	270	VAL	CA-C-N	5.37	125.69	119.47
1	A	270	VAL	C-N-CA	5.37	125.69	119.47
1	A	78	VAL	CA-C-N	5.31	125.25	119.78
1	A	78	VAL	C-N-CA	5.31	125.25	119.78
1	A	85	SER	CA-C-N	5.19	125.47	119.92
1	A	85	SER	C-N-CA	5.19	125.47	119.92

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2745	0	2666	20	0
2	B	3203	0	3130	29	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
3	B	1	0	0	0	0
4	B	24	0	25	2	0
5	B	42	0	40	3	0
6	A	217	0	0	3	0
6	B	218	0	0	1	0
All	All	6450	0	5861	52	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 4.

All (52) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:152:MET:O	1:A:156:ILE:HG23	1.67	0.92
1:A:346:LYS:HA	1:A:346:LYS:HE2	1.58	0.84
6:A:1103:HOH:O	2:B:98:ALA:HB3	1.77	0.81
1:A:346:LYS:HE3	1:A:357:ARG:HH22	1.47	0.79
1:A:152:MET:O	1:A:156:ILE:CG2	2.35	0.74
2:B:87:ARG:NH2	2:B:125:ASP:OD2	2.34	0.59
2:B:308:LEU:HD13	2:B:330:PHE:HB3	1.85	0.58
2:B:28:PRO:HD2	2:B:29:GLU:OE1	2.05	0.57
1:A:303:GLN:HB2	1:A:304:PRO:HD3	1.88	0.56
2:B:75:ARG:NH2	2:B:393:VAL:O	2.39	0.56
2:B:325:MET:HA	2:B:325:MET:HE2	1.87	0.56
2:B:325:MET:SD	2:B:381:MET:HG3	2.46	0.54
1:A:329:ASN:O	1:A:330:LYS:C	2.52	0.52
2:B:149:HIS:ND1	2:B:152:PRO:HD2	2.27	0.50
6:A:1103:HOH:O	5:B:1003:ED4:CAE	2.59	0.50
1:A:96:PHE:HA	1:A:126:LEU:HD13	1.94	0.49
2:B:218:THR:HB	2:B:219:PRO:HD2	1.95	0.49
2:B:239:ILE:HB	2:B:252:THR:HA	1.95	0.49
1:A:294:ASN:O	1:A:298:GLN:HG3	2.13	0.48
1:A:302:LEU:HB3	1:A:306:HIS:HB2	1.95	0.48
1:A:77:PRO:HG3	1:A:102:TYR:CZ	2.48	0.48
2:B:202:ARG:HD2	4:B:1002:FPP:H152	1.96	0.47
2:B:376:PHE:CD2	2:B:376:PHE:C	2.93	0.47
2:B:149:HIS:HE1	2:B:193:MET:HE2	1.79	0.47
2:B:121:ILE:O	2:B:124:THR:HG22	2.15	0.46
2:B:192:LEU:HD23	2:B:199:VAL:CG2	2.46	0.46
1:A:87:VAL:O	1:A:88:VAL:C	2.59	0.46
2:B:282:MET:SD	2:B:289:GLN:HB3	2.56	0.45

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:326:GLN:HA	1:A:330:LYS:HE3	1.98	0.45
1:A:346:LYS:HE3	1:A:357:ARG:NH2	2.23	0.45
2:B:201:VAL:HG12	2:B:251:TYR:CD2	2.52	0.45
1:A:346:LYS:HA	1:A:346:LYS:CE	2.40	0.44
2:B:192:LEU:HD23	2:B:199:VAL:HG23	2.01	0.43
2:B:201:VAL:HG11	2:B:251:TYR:HB3	2.00	0.43
1:A:311:LEU:C	1:A:311:LEU:HD23	2.43	0.43
2:B:193:MET:SD	2:B:203:SER:HB3	2.58	0.43
2:B:149:HIS:CE1	2:B:193:MET:HE2	2.53	0.43
1:A:71:GLU:OE1	1:A:71:GLU:N	2.48	0.43
2:B:55:ILE:HD13	2:B:354:PRO:HG2	2.02	0.42
2:B:253:PHE:HA	2:B:307:LEU:HD21	2.02	0.42
1:A:189:ILE:HD11	1:A:205:HIS:HD2	1.85	0.42
6:A:1103:HOH:O	5:B:1003:ED4:HAEA	2.19	0.41
2:B:324:SER:O	2:B:325:MET:HE2	2.20	0.41
1:A:156:ILE:HG22	1:A:172:ARG:HH12	1.86	0.41
2:B:301:SER:O	2:B:305:ALA:HB3	2.20	0.41
2:B:423:PHE:N	2:B:423:PHE:CD2	2.88	0.41
1:A:287:ARG:HA	1:A:287:ARG:HD2	1.64	0.40
1:A:308:SER:O	1:A:312:ILE:HG12	2.21	0.40
2:B:51:VAL:O	2:B:55:ILE:HG12	2.22	0.40
2:B:174:ARG:HD3	6:B:1120:HOH:O	2.21	0.40
2:B:423:PHE:N	2:B:423:PHE:HD2	2.18	0.40
4:B:1002:FPP:H102	5:B:1003:ED4:NBB	2.36	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	321/377 (85%)	304 (95%)	16 (5%)	1 (0%)	36 45

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
2	B	405/437 (93%)	396 (98%)	9 (2%)	0	100	100
All	All	726/814 (89%)	700 (96%)	25 (3%)	1 (0%)	48	61

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	326	GLN

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	301/338 (89%)	291 (97%)	10 (3%)	33	48
2	B	344/371 (93%)	341 (99%)	3 (1%)	70	81
All	All	645/709 (91%)	632 (98%)	13 (2%)	48	63

All (13) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	58	LEU
1	A	109	ARG
1	A	143	SER
1	A	156	ILE
1	A	173	ARG
1	A	303	GLN
1	A	324	GLU
1	A	334	LEU
1	A	348	LYS
1	A	366	LYS
2	B	53	GLU
2	B	68	VAL
2	B	351	LEU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (11) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	80	GLN
1	A	89	GLN
1	A	135	HIS
1	A	145	GLN
1	A	303	GLN
1	A	326	GLN
1	A	364	GLN
2	B	128	GLN
2	B	179	GLN
2	B	332	GLN
2	B	383	HIS

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 3 ligands modelled in this entry, 1 is monoatomic - leaving 2 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# $ Z > 2$	Counts	RMSZ	# $ Z > 2$
5	ED4	B	1003	3	43,45,45	2.65	8 (18%)	54,65,65	2.82	13 (24%)
4	FPP	B	1002	-	22,23,23	2.10	8 (36%)	27,31,31	1.24	4 (14%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
5	ED4	B	1003	3	-	12/38/50/50	0/4/4/4
4	FPP	B	1002	-	-	1/25/25/25	-

All (16) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
5	B	1003	ED4	CAY-NBK	-9.28	1.27	1.45
5	B	1003	ED4	CBE-CAJ	-6.83	1.30	1.44
5	B	1003	ED4	OBC-CBD	6.63	1.45	1.33
5	B	1003	ED4	OAI-SBP	6.31	1.49	1.43
5	B	1003	ED4	OAH-SBP	5.52	1.48	1.43
5	B	1003	ED4	CBH-SBP	-4.34	1.65	1.76
4	B	1002	FPP	C11-C12	-3.51	1.39	1.50
4	B	1002	FPP	PB-O1B	3.34	1.60	1.50
4	B	1002	FPP	C6-C7	-3.29	1.40	1.50
4	B	1002	FPP	C1-C2	-3.16	1.40	1.49
4	B	1002	FPP	C7-C8	3.02	1.40	1.33
4	B	1002	FPP	C2-C3	2.89	1.39	1.33
5	B	1003	ED4	OBC-CBO	-2.82	1.43	1.48
4	B	1002	FPP	PA-O1A	2.73	1.60	1.50
4	B	1002	FPP	C12-C13	2.50	1.39	1.32
5	B	1003	ED4	CBG-NBN	-2.28	1.34	1.38

All (17) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
5	B	1003	ED4	CAR-NBM-CAQ	8.83	110.20	106.71
5	B	1003	ED4	NBM-CAQ-NBB	-7.81	109.67	112.66
5	B	1003	ED4	OBC-CBD-NBJ	7.70	122.46	110.78
5	B	1003	ED4	OAH-SBP-OAI	-6.42	110.48	119.69
5	B	1003	ED4	CAX-NBL-CAZ	-5.80	107.35	116.80
5	B	1003	ED4	OBC-CBD-OAG	-5.78	117.13	126.43
5	B	1003	ED4	NBN-CAP-NBA	-5.66	109.57	112.94
5	B	1003	ED4	CAU-NBJ-CAV	4.27	121.39	112.68
5	B	1003	ED4	CAO-NBA-CAP	3.45	109.75	105.24
5	B	1003	ED4	CBG-CAO-NBA	-3.27	102.85	109.79
5	B	1003	ED4	CAX-NBL-SBP	-2.90	111.93	117.88
5	B	1003	ED4	CAZ-NBL-SBP	-2.89	111.02	117.32

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
4	B	1002	FPP	O2A-PA-O3A	2.74	114.67	107.27
4	B	1002	FPP	C10-C8-C9	2.58	119.71	115.23
5	B	1003	ED4	OAG-CBD-NBJ	-2.11	120.40	124.30
4	B	1002	FPP	O2B-PB-O1B	-2.02	102.96	110.83
4	B	1002	FPP	O3B-PB-O2B	2.02	115.36	107.80

There are no chirality outliers.

All (13) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
5	B	1003	ED4	CAX-NBL-SBP-CBH
5	B	1003	ED4	NBL-CAZ-CBI-CAS
5	B	1003	ED4	NBJ-CBD-OBC-CBO
5	B	1003	ED4	OAG-CBD-OBC-CBO
5	B	1003	ED4	NBL-CAZ-CBI-CAT
5	B	1003	ED4	CAX-NBL-SBP-OAI
5	B	1003	ED4	CAX-NBL-SBP-OAH
5	B	1003	ED4	CAZ-NBL-SBP-OAI
5	B	1003	ED4	CAD-CBO-OBC-CBD
5	B	1003	ED4	CAE-CBO-OBC-CBD
5	B	1003	ED4	CAZ-NBL-SBP-OAH
5	B	1003	ED4	CAZ-NBL-SBP-CBH
4	B	1002	FPP	PB-O3A-PA-O2A


There are no ring outliers.

2 monomers are involved in 4 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
5	B	1003	ED4	3	0
4	B	1002	FPP	2	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient

equivalents in the CSD to analyse the geometry.



5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	323/377 (85%)	0.14	9 (2%) 55 56	17, 35, 62, 72	0
2	B	407/437 (93%)	-0.08	9 (2%) 62 64	19, 32, 50, 67	0
All	All	730/814 (89%)	0.02	18 (2%) 58 61	17, 33, 56, 72	0

All (18) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
2	B	19	TRP	4.4
1	A	377	VAL	4.2
2	B	17	PRO	3.6
1	A	375	ALA	3.5
1	A	55	PHE	3.4
1	A	374	PRO	3.4
2	B	18	VAL	3.3
2	B	423	PHE	3.1
1	A	376	SER	3.0
1	A	286	ASP	2.8
1	A	372	ASP	2.6
2	B	325	MET	2.4
1	A	373	ILE	2.3
2	B	20	SER	2.3
2	B	64	PHE	2.2
1	A	371	SER	2.1
2	B	266	ARG	2.1
2	B	417	GLN	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)


There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.




Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
5	ED4	B	1003	42/42	0.93	0.10	21,32,48,49	0
4	FPP	B	1002	24/24	0.97	0.08	20,24,27,29	0
3	ZN	B	1001	1/1	1.00	0.01	20,20,20,20	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.



Electron density around FPP B 1002:

$2mF_o - DF_c$ (at 0.7 rmsd) in gray
 $mF_o - DF_c$ (at 3 rmsd) in purple (negative)
and green (positive)



6.5 Other polymers [i](#)

There are no such residues in this entry.