



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 24, 2026 – 12:45 AM UTC

PDB ID : 3E33 / pdb_00003e33
Title : Protein farnesyltransferase complexed with FPP and ethylenediamine scaffold inhibitor 7
Authors : Hast, M.A.; Beese, L.S.
Deposited on : 2008-08-06
Resolution : 1.90 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0
Mogul : 2022.3.0, CSD as543be (2022)
Xtriage (Phenix) : 2.0
EDS : 3.0
Buster-report : wwPDB partial adaption of 1.1.7 (2018)
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4 : 9.0.010 (Gargrove)
Density-Fitness : 1.0.12
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.90 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
Clashscore	190562	8410 (1.90-1.90)
Ramachandran outliers	187476	8333 (1.90-1.90)
Sidechain outliers	187428	8333 (1.90-1.90)
RSRZ outliers	180081	7790 (1.90-1.90)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	377	
2	B	437	

2 Entry composition [i](#)

There are 6 unique types of molecules in this entry. The entry contains 6239 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Protein farnesyltransferase/geranylgeranyltransferase type-1 subunit alpha.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	323	Total	C	N	O	S	0	0	0
			2745	1746	481	513	5			


- Molecule 2 is a protein called Protein farnesyltransferase subunit beta.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	B	407	Total	C	N	O	S	0	0	0
			3203	2050	550	580	23			

- Molecule 3 is ZINC ION (CCD ID: ZN) (formula: Zn).


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	B	1	Total	Zn	0	0
			1	1		

- Molecule 4 is FARNESYL DIPHOSPHATE (CCD ID: FPP) (formula: C₁₅H₂₈O₇P₂).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
4	B	1	Total	C	O	P	0	0
			24	15	7	2		

- Molecule 5 is N-benzyl-N-(2-{(4-cyanophenyl)[(1-methyl-1H-imidazol-5-yl)methyl]amino}ethyl)-2-methylbenzenesulfonamide (CCD ID: ED7) (formula: $C_{28}H_{29}N_5O_2S$).




Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
5	B	1	Total	C	N	O	S	0	0
			36	28	5	2	1		

- Molecule 6 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
6	A	117	Total O 117 117	0	0
6	B	113	Total O 113 113	0	0

- Molecule 1: Protein farnesyltransferase/geranylgeranyltransferase type-1 subunit alpha



4 Data and refinement statistics

Property	Value	Source
Space group	P 61	Depositor
Cell constants a, b, c, α , β , γ	171.04Å 171.04Å 69.39Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	33.98 – 1.90 33.98 – 1.90	Depositor EDS
% Data completeness (in resolution range)	99.2 (33.98-1.90) 99.2 (33.98-1.90)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.29 (at 1.89Å)	Xtriage
Refinement program	REFMAC 5.4.0077	Depositor
R, R_{free}	0.201 , 0.222 0.202 , (Not available)	Depositor DCC
R_{free} test set	No test flags present.	wwPDB-VP
Wilson B-factor (Å ²)	27.9	Xtriage
Anisotropy	0.115	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.38 , 35.6	EDS
L-test for twinning ²	$\langle L \rangle = 0.50$, $\langle L^2 \rangle = 0.33$	Xtriage
Estimated twinning fraction	0.022 for h,-h-k,-l	Xtriage
F_o, F_c correlation	0.95	EDS
Total number of atoms	6239	wwPDB-VP
Average B, all atoms (Å ²)	30.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.29% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: ED7, ZN, FPP

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.56	0/2812	0.73	3/3817 (0.1%)
2	B	0.56	0/3292	0.74	0/4473
All	All	0.56	0/6104	0.74	3/8290 (0.0%)

There are no bond length outliers.

All (3) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	85	SER	CA-C-N	5.60	125.55	119.78
1	A	85	SER	C-N-CA	5.60	125.55	119.78
1	A	217	ASP	N-CA-C	5.40	117.16	111.28

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2745	0	2666	22	0
2	B	3203	0	3130	23	0
3	B	1	0	0	0	0
4	B	24	0	25	2	0
5	B	36	0	29	1	0
6	A	117	0	0	1	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
6	B	113	0	0	1	0
All	All	6239	0	5850	44	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 4.

All (44) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:152:MET:O	1:A:156:ILE:HG23	1.71	0.90
1:A:302:LEU:C	1:A:304:PRO:CD	2.50	0.83
1:A:156:ILE:HG22	1:A:172:ARG:HH12	1.46	0.79
1:A:302:LEU:C	1:A:304:PRO:HD3	2.15	0.72
1:A:302:LEU:C	1:A:304:PRO:HD2	2.16	0.70
1:A:302:LEU:O	1:A:304:PRO:HD2	1.93	0.68
1:A:303:GLN:N	1:A:304:PRO:HD3	2.13	0.64
2:B:308:LEU:HD13	2:B:330:PHE:HB3	1.82	0.60
2:B:325:MET:SD	2:B:381:MET:HG3	2.42	0.59
1:A:152:MET:O	1:A:156:ILE:CG2	2.50	0.57
1:A:77:PRO:HG3	1:A:102:TYR:CZ	2.40	0.55
2:B:193:MET:SD	2:B:203:SER:HB3	2.46	0.55
4:B:1002:FPP:H102	5:B:1003:ED7:HAN	1.89	0.54
2:B:75:ARG:NH2	2:B:393:VAL:O	2.41	0.54
1:A:303:GLN:N	1:A:304:PRO:CD	2.69	0.53
1:A:156:ILE:HG22	1:A:172:ARG:NH1	2.19	0.53
2:B:239:ILE:HB	2:B:252:THR:HA	1.91	0.52
2:B:30:HIS:HE1	6:B:2824:HOH:O	1.92	0.51
1:A:311:LEU:C	1:A:311:LEU:HD23	2.36	0.50
2:B:202:ARG:HD2	4:B:1002:FPP:H142	1.92	0.50
2:B:51:VAL:O	2:B:55:ILE:HG12	2.13	0.49
2:B:422:GLY:O	2:B:423:PHE:HB2	2.12	0.49
2:B:192:LEU:HD23	2:B:199:VAL:CG2	2.42	0.48
2:B:185:LYS:HD3	2:B:191:PHE:CZ	2.48	0.48
2:B:28:PRO:HD2	2:B:29:GLU:OE1	2.14	0.48
1:A:361:ARG:HB3	2:B:322:ALA:HB1	1.96	0.47
1:A:208:TRP:NE1	1:A:212:GLU:HG3	2.30	0.46
2:B:220:ASP:HA	2:B:223:GLU:HG2	1.97	0.46
1:A:156:ILE:HG21	6:A:2672:HOH:O	2.17	0.45
1:A:112:ARG:HA	1:A:140:LEU:CD2	2.46	0.45
1:A:294:ASN:O	1:A:298:GLN:HG3	2.15	0.45
2:B:325:MET:HA	2:B:325:MET:HE2	1.99	0.44

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:B:192:LEU:HD23	2:B:199:VAL:HG23	2.00	0.43
2:B:408:VAL:O	2:B:412:THR:HG23	2.19	0.43
1:A:198:LYS:HD2	2:B:294:LYS:HE3	2.00	0.43
2:B:301:SER:O	2:B:305:ALA:HB3	2.17	0.43
2:B:333:GLN:HG3	2:B:387:MET:SD	2.59	0.43
1:A:156:ILE:CG2	1:A:172:ARG:HH12	2.24	0.42
2:B:253:PHE:HA	2:B:307:LEU:HD21	2.01	0.42
2:B:220:ASP:HA	2:B:223:GLU:CG	2.50	0.42
1:A:346:LYS:HE3	1:A:357:ARG:NH2	2.35	0.41
1:A:373:ILE:HA	1:A:374:PRO:HD3	1.97	0.41
1:A:329:ASN:O	1:A:330:LYS:C	2.64	0.40
2:B:276:TRP:O	2:B:280:ARG:HG2	2.22	0.40

There are no symmetry-related clashes.

5.3 Torsion angles ⓘ

5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	321/377 (85%)	307 (96%)	13 (4%)	1 (0%)	36	29
2	B	405/437 (93%)	398 (98%)	7 (2%)	0	100	100
All	All	726/814 (89%)	705 (97%)	20 (3%)	1 (0%)	48	40

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	304	PRO

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	301/338 (89%)	292 (97%)	9 (3%)	36	30
2	B	344/371 (93%)	339 (98%)	5 (2%)	57	56
All	All	645/709 (91%)	631 (98%)	14 (2%)	45	42

All (14) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	55	PHE
1	A	85	SER
1	A	156	ILE
1	A	257	GLU
1	A	269	LEU
1	A	303	GLN
1	A	324	GLU
1	A	334	LEU
1	A	348	LYS
2	B	29	GLU
2	B	68	VAL
2	B	87	ARG
2	B	265	GLU
2	B	351	LEU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (10) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	89	GLN
1	A	108	GLN
1	A	135	HIS
1	A	145	GLN
1	A	162	GLN
1	A	241	HIS
2	B	56	GLN
2	B	128	GLN

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
2	B	194	HIS
2	B	410	GLN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no oligosaccharides in this entry.

5.6 Ligand geometry ⓘ

Of 3 ligands modelled in this entry, 1 is monoatomic - leaving 2 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
4	FPP	B	1002	-	22,23,23	2.12	5 (22%)	27,31,31	1.18	3 (11%)
5	ED7	B	1003	3	39,39,39	2.01	8 (20%)	49,54,54	2.34	9 (18%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	FPP	B	1002	-	-	3/25/25/25	-

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
5	ED7	B	1003	3	-	5/31/31/31	0/4/4/4

All (13) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
5	B	1003	ED7	OAE-SBJ	7.45	1.51	1.43
5	B	1003	ED7	OAD-SBJ	6.61	1.50	1.43
4	B	1002	FPP	PB-O2B	5.39	1.74	1.54
4	B	1002	FPP	PB-O3B	5.16	1.74	1.54
4	B	1002	FPP	PA-O2A	5.11	1.79	1.55
5	B	1003	ED7	CBF-SBJ	-3.05	1.74	1.78
5	B	1003	ED7	CAY-CBE	2.90	1.53	1.50
5	B	1003	ED7	SBJ-NBH	2.78	1.67	1.63
5	B	1003	ED7	CBE-NBI	-2.74	1.33	1.38
4	B	1002	FPP	PB-O1B	2.28	1.57	1.50
4	B	1002	FPP	PA-O1A	2.15	1.58	1.50
5	B	1003	ED7	CAB-NBI	2.11	1.50	1.46
5	B	1003	ED7	CAT-CBE	-2.02	1.34	1.36

All (12) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
5	B	1003	ED7	NBI-CAU-NAZ	-10.43	106.73	112.94
5	B	1003	ED7	OAE-SBJ-OAD	-7.05	108.59	119.59
5	B	1003	ED7	OAD-SBJ-NBH	5.17	111.55	106.69
5	B	1003	ED7	CAX-NBH-SBJ	-4.12	107.75	117.47
5	B	1003	ED7	CAA-CBA-CBF	-3.57	120.79	124.17
5	B	1003	ED7	CBF-SBJ-NBH	2.94	112.15	106.69
4	B	1002	FPP	C10-C8-C9	2.80	120.09	115.23
5	B	1003	ED7	CAT-NAZ-CAU	2.70	108.77	105.24
5	B	1003	ED7	CAL-CBA-CBF	2.69	119.11	116.27
5	B	1003	ED7	CAB-NBI-CBE	-2.39	124.15	126.69
4	B	1002	FPP	C6-C7-C8	-2.25	122.48	127.62
4	B	1002	FPP	C15-C13-C14	2.16	119.55	114.59

There are no chirality outliers.

All (8) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
5	B	1003	ED7	CAX-NBH-SBJ-CBF
5	B	1003	ED7	CAW-NBH-SBJ-OAD

Continued on next page...

Continued from previous page...


Mol	Chain	Res	Type	Atoms
5	B	1003	ED7	CAW-NBH-SBJ-CBF
5	B	1003	ED7	CAX-NBH-SBJ-OAE
4	B	1002	FPP	C4-C3-C5-C6
4	B	1002	FPP	C2-C3-C5-C6
5	B	1003	ED7	CBC-CAX-NBH-SBJ
4	B	1002	FPP	PB-O3A-PA-O2A


There are no ring outliers.

2 monomers are involved in 2 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
4	B	1002	FPP	2	0
5	B	1003	ED7	1	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.





5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	323/377 (85%)	0.90	49 (15%) 5 5	17, 31, 54, 62	0
2	B	407/437 (93%)	0.60	33 (8%) 18 19	18, 27, 45, 60	0
All	All	730/814 (89%)	0.73	82 (11%) 10 10	17, 28, 50, 62	0

All (82) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
2	B	19	TRP	7.2
1	A	55	PHE	7.2
2	B	423	PHE	6.5
2	B	17	PRO	6.2
2	B	380	ALA	5.9
1	A	375	ALA	5.7
1	A	377	VAL	5.5
1	A	374	PRO	5.4
1	A	373	ILE	5.3
2	B	64	PHE	5.2
2	B	18	VAL	5.0
1	A	334	LEU	4.7
1	A	58	LEU	4.5
2	B	121	ILE	4.5
1	A	304	PRO	4.3
2	B	70	ARG	4.0
1	A	56	LEU	3.9
1	A	286	ASP	3.7
1	A	371	SER	3.6
2	B	325	MET	3.6
2	B	382	LEU	3.6
1	A	327	CYS	3.6
1	A	364	GLN	3.6
1	A	376	SER	3.4

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
2	B	175	GLU	3.3
1	A	59	ASP	3.3
1	A	305	SER	3.3
2	B	422	GLY	3.2
1	A	300	LEU	3.1
2	B	20	SER	3.1
1	A	326	GLN	3.1
2	B	266	ARG	3.1
1	A	306	HIS	3.0
1	A	372	ASP	3.0
2	B	317	ALA	3.0
1	A	302	LEU	2.9
1	A	265	GLU	2.9
1	A	328	ASP	2.9
2	B	319	GLY	2.9
2	B	21	GLU	2.9
1	A	332	ASP	2.8
2	B	420	VAL	2.8
2	B	381	MET	2.7
1	A	330	LYS	2.7
2	B	62	TYR	2.7
1	A	337	ALA	2.6
1	A	314	PHE	2.6
2	B	386	VAL	2.6
1	A	217	ASP	2.6
2	B	120	GLN	2.6
1	A	333	ILE	2.6
2	B	80	HIS	2.5
1	A	296	LEU	2.5
2	B	416	LEU	2.5
1	A	331	GLU	2.5
2	B	419	PRO	2.5
2	B	118	ILE	2.5
1	A	287	ARG	2.4
2	B	379	GLY	2.4
2	B	22	PRO	2.4
2	B	71	LEU	2.4
1	A	310	TYR	2.4
1	A	75	ILE	2.4
2	B	76	GLU	2.4
1	A	292	TYR	2.3
1	A	368	SER	2.3

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	356	TRP	2.2
1	A	339	GLU	2.2
1	A	336	LYS	2.2
1	A	299	LEU	2.2
1	A	338	LEU	2.2
1	A	324	GLU	2.2
2	B	23	LEU	2.2
1	A	366	LYS	2.1
2	B	223	GLU	2.1
2	B	347	ALA	2.1
1	A	329	ASN	2.1
1	A	312	ILE	2.1
1	A	370	GLU	2.1
1	A	303	GLN	2.0
1	A	317	ASP	2.0
1	A	316	VAL	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)




In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
5	ED7	B	1003	36/36	0.91	0.11	19,31,39,41	0
4	FPP	B	1002	24/24	0.94	0.14	22,30,35,38	0
3	ZN	B	1001	1/1	1.00	0.02	20,20,20,20	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.




Electron density around ED7 B 1003:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



Electron density around FPP B 1002:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



6.5 Other polymers [i](#)

There are no such residues in this entry.