



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 5, 2026 – 09:51 PM UTC

PDB ID : 3E3F / pdb_00003e3f
Title : H. influenzae beta-carbonic anhydrase, variant V47A with 100 mM bicarbonate
Authors : Rowlett, R.S.; Mysliwiec, M.
Deposited on : 2008-08-07
Resolution : 2.30 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0
Mogul : 2022.3.0, CSD as543be (2022)
Xtriage (Phenix) : 2.0
EDS : 3.0
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4 : 9.0.010 (Gargrove)
Density-Fitness : 1.0.12
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.30 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	6319 (2.30-2.30)
Clashscore	190562	6919 (2.30-2.30)
Ramachandran outliers	187476	6854 (2.30-2.30)
Sidechain outliers	187428	6854 (2.30-2.30)
RSRZ outliers	180081	6325 (2.30-2.30)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	229	
1	B	229	

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
3	BCT	A	233	-	-	X	-

2 Entry composition

There are 4 unique types of molecules in this entry. The entry contains 3000 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Carbonic anhydrase 2.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	182	Total	C	N	O	S	0	0	0
			1443	918	259	259	7			
1	B	183	Total	C	N	O	S	0	0	0
			1451	922	260	262	7			

There are 2 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	47	ALA	VAL	engineered mutation	UNP P45148
B	47	ALA	VAL	engineered mutation	UNP P45148

- Molecule 2 is ZINC ION (CCD ID: ZN) (formula: Zn).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	A	1	Total	Zn	0	0
			1	1		
2	B	1	Total	Zn	0	0
			1	1		

- Molecule 3 is BICARBONATE ION (CCD ID: BCT) (formula: CHO₃).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	A	1	Total	C	O	0	0
			4	1	3		
3	A	1	Total	C	O	0	0
			4	1	3		
3	A	1	Total	C	O	0	0
			4	1	3		
3	A	1	Total	C	O	0	0
			4	1	3		
3	B	1	Total	C	O	0	0
			4	1	3		


- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	47	Total	O	0	0
			47	47		
4	B	37	Total	O	0	0
			37	37		


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: Carbonic anhydrase 2



• Molecule 1: Carbonic anhydrase 2



4 Data and refinement statistics

Property	Value	Source
Space group	P 41 21 2	Depositor
Cell constants a, b, c, α , β , γ	83.90Å 83.90Å 184.87Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	28.93 – 2.30 28.93 – 2.30	Depositor EDS
% Data completeness (in resolution range)	99.1 (28.93-2.30) 99.0 (28.93-2.30)	Depositor EDS
R_{merge}	0.07	Depositor
R_{sym}	0.07	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.50 (at 2.31Å)	Xtriage
Refinement program	REFMAC	Depositor
R, R_{free}	0.193 , 0.230 0.241 , 0.261	Depositor DCC
R_{free} test set	1523 reflections (5.05%)	wwPDB-VP
Wilson B-factor (Å ²)	34.1	Xtriage
Anisotropy	1.243	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.37 , 64.1	EDS
L-test for twinning ²	$\langle L \rangle = 0.47$, $\langle L^2 \rangle = 0.30$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.91	EDS
Total number of atoms	3000	wwPDB-VP
Average B, all atoms (Å ²)	48.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.98% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: BCT, ZN

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	1.06	1/1475 (0.1%)	1.01	2/2002 (0.1%)
1	B	1.00	3/1483 (0.2%)	1.03	0/2013
All	All	1.03	4/2958 (0.1%)	1.02	2/4015 (0.0%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	1

All (4) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	B	166	SER	C-O	-5.89	1.17	1.24
1	A	123	ILE	CA-CB	5.69	1.61	1.54
1	B	193	VAL	CA-CB	-5.19	1.48	1.54
1	B	116	ILE	C-O	-5.09	1.18	1.24

All (2) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	44	ASP	CB-CA-C	-5.31	102.72	110.95
1	A	61	PHE	N-CA-C	-5.02	100.99	109.07

There are no chirality outliers.

All (1) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	45	SER	Peptide

5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1443	0	1439	27	0
1	B	1451	0	1444	22	0
2	A	1	0	0	0	0
2	B	1	0	0	1	0
3	A	16	0	0	5	0
3	B	4	0	0	0	0
4	A	47	0	0	0	0
4	B	37	0	0	0	0
All	All	3000	0	2883	49	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 8.

All (49) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:47:ALA:CB	1:B:52:LEU:HD11	2.11	0.80
1:A:198:ARG:NH1	3:A:233:BCT:O2	2.15	0.79
1:B:86:ASP:O	1:B:89:LYS:NZ	2.19	0.71
1:A:165:LYS:NZ	3:A:233:BCT:O2	2.29	0.64
1:B:47:ALA:HB3	1:B:52:LEU:HD11	1.78	0.64
1:B:101:CYS:SG	2:B:230:ZN:ZN	1.88	0.61
1:A:45:SER:C	1:A:46:ARG:O	2.44	0.61
1:B:116:ILE:HG13	1:B:120:LEU:HD22	1.84	0.59
1:A:46:ARG:CZ	1:A:98:HIS:HD1	2.17	0.57
1:B:110:ASP:O	1:B:111:LYS:O	2.21	0.57
1:B:53:THR:O	1:B:55:LEU:HD13	2.06	0.56
1:B:47:ALA:HB1	1:B:52:LEU:HD21	1.86	0.55
1:A:50:GLU:OE2	3:A:231:BCT:O3	2.25	0.54
1:A:130:HIS:CD2	1:A:149:ILE:HG21	2.44	0.53
1:A:198:ARG:NH1	3:A:233:BCT:C	2.71	0.53

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:150:ASN:O	1:B:154:GLN:HG2	2.09	0.51
1:B:50:GLU:HG3	1:B:60:LEU:HD12	1.93	0.50
1:B:50:GLU:HB3	1:B:55:LEU:O	2.12	0.50
1:B:47:ALA:HB1	1:B:52:LEU:HD11	1.90	0.49
1:B:110:ASP:O	1:B:111:LYS:C	2.57	0.48
1:A:194:MET:HE1	1:A:196:THR:HG23	1.95	0.48
1:B:37:TYR:HB2	1:B:60:LEU:HD23	1.96	0.48
1:A:191:GLN:HE21	1:A:191:GLN:N	2.11	0.48
1:A:194:MET:CE	1:A:196:THR:HG23	2.45	0.46
1:B:189:VAL:HG12	1:B:191:GLN:NE2	2.30	0.46
1:A:56:GLU:H	1:A:56:GLU:CD	2.23	0.46
1:A:199:GLU:OE1	1:A:199:GLU:N	2.37	0.46
1:A:109:ALA:HB1	1:A:111:LYS:HG2	1.99	0.45
1:A:82:GLN:HA	1:A:163:ILE:HG21	1.99	0.45
1:A:133:LEU:HD12	1:A:133:LEU:O	2.17	0.45
1:A:133:LEU:HD12	1:A:133:LEU:C	2.42	0.45
1:A:98:HIS:O	1:A:147:THR:HG21	2.17	0.44
1:A:198:ARG:HH21	1:A:202:GLU:HG2	1.82	0.44
1:B:56:GLU:H	1:B:56:GLU:CD	2.26	0.44
1:B:76:ASN:OD1	1:B:76:ASN:C	2.61	0.44
1:B:54:ASN:CG	1:B:54:ASN:O	2.60	0.44
1:B:121:LEU:HD23	1:B:124:ARG:HD3	1.99	0.44
1:A:69:GLN:HE22	1:A:122:HIS:HB2	1.83	0.43
1:A:46:ARG:NH1	1:A:98:HIS:HD1	2.16	0.43
1:A:46:ARG:N	1:A:46:ARG:CD	2.81	0.42
1:A:192:GLY:HA2	1:A:211:ARG:HH12	1.84	0.42
1:A:198:ARG:NH1	3:A:233:BCT:O3	2.52	0.42
1:A:123:ILE:HD12	1:A:150:ASN:OD1	2.19	0.42
1:B:123:ILE:CD1	1:B:150:ASN:OD1	2.67	0.42
1:A:44:ASP:O	1:A:45:SER:C	2.62	0.42
1:B:93:ILE:HG21	1:B:158:LEU:HD21	2.03	0.41
1:A:81:VAL:O	1:A:85:VAL:HG23	2.21	0.41
1:B:44:ASP:O	1:B:45:SER:C	2.63	0.41
1:A:190:ASP:O	1:A:191:GLN:C	2.65	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	180/229 (79%)	169 (94%)	7 (4%)	4 (2%)	5	4
1	B	181/229 (79%)	170 (94%)	9 (5%)	2 (1%)	11	13
All	All	361/458 (79%)	339 (94%)	16 (4%)	6 (2%)	7	7

All (6) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	46	ARG
1	B	111	LYS
1	A	191	GLN
1	B	110	ASP
1	A	47	ALA
1	A	189	VAL

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	156/200 (78%)	144 (92%)	12 (8%)	12	16
1	B	157/200 (78%)	142 (90%)	15 (10%)	8	10
All	All	313/400 (78%)	286 (91%)	27 (9%)	10	13

All (27) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	34	THR
1	A	46	ARG
1	A	56	GLU
1	A	90	ILE
1	A	133	LEU
1	A	137	LEU
1	A	189	VAL
1	A	191	GLN
1	A	193	VAL
1	A	194	MET
1	A	198	ARG
1	A	212	LEU
1	B	45	SER
1	B	55	LEU
1	B	56	GLU
1	B	59	GLU
1	B	78	LEU
1	B	90	ILE
1	B	91	GLU
1	B	120	LEU
1	B	137	LEU
1	B	140	GLU
1	B	184	ASN
1	B	194	MET
1	B	198	ARG
1	B	212	LEU
1	B	215	LEU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (7) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	69	GLN
1	A	105	HIS
1	A	132	HIS
1	A	191	GLN
1	B	100	ASN
1	B	105	HIS
1	B	132	HIS

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 7 ligands modelled in this entry, 2 are monoatomic - leaving 5 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
3	BCT	A	232	-	3,3,3	0.51	0	2,3,3	0.20	0
3	BCT	A	231	-	3,3,3	0.61	0	2,3,3	1.42	0
3	BCT	A	233	-	3,3,3	0.74	0	2,3,3	0.78	0
3	BCT	A	234	-	3,3,3	0.55	0	2,3,3	0.58	0
3	BCT	B	231	-	3,3,3	0.60	0	2,3,3	0.36	0

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

2 monomers are involved in 5 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	A	231	BCT	1	0
3	A	233	BCT	4	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	182/229 (79%)	1.36	33 (18%) 3 4	38, 46, 59, 73	0
1	B	183/229 (79%)	1.50	46 (25%) 1 2	40, 48, 65, 67	0
All	All	365/458 (79%)	1.43	79 (21%) 2 3	38, 47, 64, 73	0

All (79) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	189	VAL	7.7
1	B	187	PHE	4.3
1	A	57	PRO	4.2
1	A	191	GLN	4.2
1	A	215	LEU	3.9
1	A	46	ARG	3.8
1	A	181	TYR	3.6
1	B	52	LEU	3.5
1	B	47	ALA	3.5
1	B	127	TRP	3.4
1	A	55	LEU	3.3
1	A	56	GLU	3.3
1	B	138	SER	3.2
1	B	37	TYR	3.2
1	B	57	PRO	3.2
1	A	34	THR	3.2
1	B	49	ALA	3.2
1	B	84	ALA	3.1
1	B	192	GLY	3.1
1	B	214	ILE	3.1
1	A	44	ASP	3.1
1	A	190	ASP	3.0
1	B	34	THR	2.9
1	A	36	HIS	2.8

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	B	48	PRO	2.8
1	A	192	GLY	2.8
1	A	78	LEU	2.8
1	B	55	LEU	2.7
1	B	113	LEU	2.7
1	B	53	THR	2.7
1	A	110	ASP	2.7
1	B	184	ASN	2.7
1	B	71	ILE	2.7
1	B	36	HIS	2.6
1	B	128	PHE	2.6
1	A	133	LEU	2.6
1	A	58	GLY	2.6
1	A	128	PHE	2.6
1	B	80	VAL	2.6
1	A	101	CYS	2.5
1	B	112	ASP	2.5
1	A	127	TRP	2.5
1	B	190	ASP	2.4
1	B	136	LYS	2.4
1	B	121	LEU	2.4
1	B	188	LEU	2.4
1	A	100	ASN	2.4
1	A	40	ILE	2.4
1	A	212	LEU	2.4
1	B	109	ALA	2.3
1	B	46	ARG	2.3
1	B	56	GLU	2.3
1	B	51	LYS	2.3
1	B	193	VAL	2.3
1	B	61	PHE	2.3
1	A	139	PRO	2.2
1	B	98	HIS	2.2
1	B	175	SER	2.2
1	B	119	TRP	2.2
1	A	193	VAL	2.2
1	B	54	ASN	2.2
1	B	101	CYS	2.2
1	A	169	GLU	2.2
1	A	121	LEU	2.2
1	A	74	ASP	2.2
1	A	112	ASP	2.2

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	188	LEU	2.1
1	B	111	LYS	2.1
1	A	54	ASN	2.1
1	B	45	SER	2.1
1	B	75	PHE	2.1
1	B	107	ALA	2.1
1	B	203	ILE	2.1
1	B	83	TYR	2.1
1	A	66	VAL	2.1
1	B	133	LEU	2.0
1	A	47	ALA	2.0
1	B	67	ALA	2.0
1	B	90	ILE	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
3	BCT	A	232	4/4	0.74	0.39	86,86,86,86	0
3	BCT	A	234	4/4	0.80	0.14	26,27,27,27	4
3	BCT	B	231	4/4	0.81	0.24	56,56,56,56	0
3	BCT	A	231	4/4	0.91	0.29	34,36,37,39	0
3	BCT	A	233	4/4	0.92	0.10	22,23,23,24	4
2	ZN	B	230	1/1	0.94	0.22	39,39,39,39	0
2	ZN	A	230	1/1	0.94	0.20	43,43,43,43	0

6.5 Other polymers [i](#)

There are no such residues in this entry.