



wwPDB X-ray Structure Validation Summary Report ⓘ

Mar 13, 2026 – 08:03 PM UTC

PDB ID : 3E3M / pdb_00003e3m
Title : Crystal structure of a LacI family transcriptional regulator from *Silicibacter pomeroyi*
Authors : Bonanno, J.B.; Freeman, J.; Bain, K.T.; Iizuka, M.; Ozyurt, S.; Smith, D.; Wasserman, S.; Sauder, J.M.; Burley, S.K.; Almo, S.C.; New York SGX Research Center for Structural Genomics (NYSGXRC)
Deposited on : 2008-08-07
Resolution : 1.60 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.60 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	4673 (1.60-1.60)
Clashscore	190562	4931 (1.60-1.60)
Ramachandran outliers	187476	4831 (1.60-1.60)
Sidechain outliers	187428	4830 (1.60-1.60)
RSRZ outliers	180081	4672 (1.60-1.60)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	355	
1	B	355	
1	C	355	
1	D	355	

2 Entry composition

There are 2 unique types of molecules in this entry. The entry contains 9477 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Transcriptional regulator, LacI family.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	270	Total	C	N	O	S	0	2	0
			2105	1336	369	393	7			
1	B	270	Total	C	N	O	S	0	8	0
			2136	1359	375	394	8			
1	C	270	Total	C	N	O	S	0	1	0
			2100	1333	366	394	7			
1	D	270	Total	C	N	O	S	0	0	0
			2100	1332	369	392	7			

There are 44 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	-1	MET	-	expression tag	UNP Q5LKY7
A	0	SER	-	expression tag	UNP Q5LKY7
A	1	LEU	-	expression tag	UNP Q5LKY7
A	346	GLU	-	expression tag	UNP Q5LKY7
A	347	GLY	-	expression tag	UNP Q5LKY7
A	348	HIS	-	expression tag	UNP Q5LKY7
A	349	HIS	-	expression tag	UNP Q5LKY7
A	350	HIS	-	expression tag	UNP Q5LKY7
A	351	HIS	-	expression tag	UNP Q5LKY7
A	352	HIS	-	expression tag	UNP Q5LKY7
A	353	HIS	-	expression tag	UNP Q5LKY7
B	-1	MET	-	expression tag	UNP Q5LKY7
B	0	SER	-	expression tag	UNP Q5LKY7
B	1	LEU	-	expression tag	UNP Q5LKY7
B	346	GLU	-	expression tag	UNP Q5LKY7
B	347	GLY	-	expression tag	UNP Q5LKY7
B	348	HIS	-	expression tag	UNP Q5LKY7
B	349	HIS	-	expression tag	UNP Q5LKY7
B	350	HIS	-	expression tag	UNP Q5LKY7
B	351	HIS	-	expression tag	UNP Q5LKY7
B	352	HIS	-	expression tag	UNP Q5LKY7

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
B	353	HIS	-	expression tag	UNP Q5LKY7
C	-1	MET	-	expression tag	UNP Q5LKY7
C	0	SER	-	expression tag	UNP Q5LKY7
C	1	LEU	-	expression tag	UNP Q5LKY7
C	346	GLU	-	expression tag	UNP Q5LKY7
C	347	GLY	-	expression tag	UNP Q5LKY7
C	348	HIS	-	expression tag	UNP Q5LKY7
C	349	HIS	-	expression tag	UNP Q5LKY7
C	350	HIS	-	expression tag	UNP Q5LKY7
C	351	HIS	-	expression tag	UNP Q5LKY7
C	352	HIS	-	expression tag	UNP Q5LKY7
C	353	HIS	-	expression tag	UNP Q5LKY7
D	-1	MET	-	expression tag	UNP Q5LKY7
D	0	SER	-	expression tag	UNP Q5LKY7
D	1	LEU	-	expression tag	UNP Q5LKY7
D	346	GLU	-	expression tag	UNP Q5LKY7
D	347	GLY	-	expression tag	UNP Q5LKY7
D	348	HIS	-	expression tag	UNP Q5LKY7
D	349	HIS	-	expression tag	UNP Q5LKY7
D	350	HIS	-	expression tag	UNP Q5LKY7
D	351	HIS	-	expression tag	UNP Q5LKY7
D	352	HIS	-	expression tag	UNP Q5LKY7
D	353	HIS	-	expression tag	UNP Q5LKY7


- Molecule 2 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
2	A	266	Total O 266 266	0	0
2	B	272	Total O 272 272	0	0
2	C	229	Total O 229 229	0	0
2	D	269	Total O 269 269	0	0


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Transcriptional regulator, LacI family



- Molecule 1: Transcriptional regulator, LacI family




- Molecule 1: Transcriptional regulator, LacI family



- Molecule 1: Transcriptional regulator, LacI family





4 Data and refinement statistics

Property	Value	Source
Space group	P 1	Depositor
Cell constants a, b, c, α , β , γ	60.40Å 60.44Å 90.77Å 82.91° 76.96° 87.22°	Depositor
Resolution (Å)	15.00 – 1.60 15.00 – 1.60	Depositor EDS
% Data completeness (in resolution range)	95.6 (15.00-1.60) 95.4 (15.00-1.60)	Depositor EDS
R_{merge}	0.09	Depositor
R_{sym}	0.09	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.00 (at 1.60Å)	Xtriage
Refinement program	REFMAC	Depositor
R, R_{free}	0.184 , 0.215 0.188 , 0.217	Depositor DCC
R_{free} test set	7881 reflections (5.04%)	wwPDB-VP
Wilson B-factor (Å ²)	14.9	Xtriage
Anisotropy	0.117	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.41 , 48.3	EDS
L-test for twinning ²	$\langle L \rangle = 0.47$, $\langle L^2 \rangle = 0.29$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.95	EDS
Total number of atoms	9477	wwPDB-VP
Average B, all atoms (Å ²)	22.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 5.34% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.62	0/2152	0.78	0/2919
1	B	0.59	0/2201	0.79	0/2984
1	C	0.57	0/2144	0.78	0/2909
1	D	0.61	0/2141	0.78	0/2904
All	All	0.59	0/8638	0.78	0/11716

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2105	0	2118	4	0
1	B	2136	0	2177	9	0
1	C	2100	0	2106	10	0
1	D	2100	0	2111	8	0
2	A	266	0	0	3	0
2	B	272	0	0	1	0
2	C	229	0	0	1	0
2	D	269	0	0	1	0
All	All	9477	0	8512	29	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 2.

The worst 5 of 29 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:74:LEU:HD21	1:B:120[A]:MET:HE1	1.59	0.82
1:C:156:LYS:NZ	1:C:332:THR:OG1	2.14	0.73
1:C:240:GLU:OE2	1:C:266:ARG:NE	2.33	0.60
1:D:122:ARG:HD2	2:D:387:HOH:O	2.07	0.55
1:A:103:LEU:HB2	1:B:103:LEU:HB2	1.90	0.54

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	268/355 (76%)	262 (98%)	6 (2%)	0	100	100
1	B	274/355 (77%)	270 (98%)	4 (2%)	0	100	100
1	C	267/355 (75%)	262 (98%)	5 (2%)	0	100	100
1	D	266/355 (75%)	262 (98%)	4 (2%)	0	100	100
All	All	1075/1420 (76%)	1056 (98%)	19 (2%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	226/301 (75%)	222 (98%)	4 (2%)	51	28
1	B	232/301 (77%)	227 (98%)	5 (2%)	45	22
1	C	225/301 (75%)	224 (100%)	1 (0%)	84	75
1	D	225/301 (75%)	222 (99%)	3 (1%)	61	40
All	All	908/1204 (75%)	895 (99%)	13 (1%)	61	38

5 of 13 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	B	306	ILE
1	B	333	LEU
1	D	145	ARG
1	D	75	LEU
1	D	118	GLU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 9 such sidechains are listed below:

Mol	Chain	Res	Type
1	D	244	GLN
1	D	277	GLN
1	A	221	GLN
1	C	115	GLN
1	C	159	HIS

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

There are no ligands in this entry.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	270/355 (76%)	-0.22	4 (1%) 72 76	10, 18, 28, 37	2 (0%)
1	B	270/355 (76%)	-0.15	1 (0%) 88 91	11, 19, 28, 32	8 (2%)
1	C	270/355 (76%)	0.08	3 (1%) 78 82	13, 22, 30, 39	1 (0%)
1	D	270/355 (76%)	-0.11	1 (0%) 88 91	13, 20, 30, 39	0
All	All	1080/1420 (76%)	-0.10	9 (0%) 82 86	10, 20, 29, 39	11 (1%)

The worst 5 of 9 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	145	ARG	2.9
1	D	147	SER	2.5
1	C	198	TRP	2.4
1	A	96	GLN	2.3
1	A	186	ARG	2.2

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

There are no ligands in this entry.

6.5 Other polymers [i](#)

There are no such residues in this entry.