



wwPDB X-ray Structure Validation Summary Report ⓘ

Mar 12, 2026 – 05:47 AM UTC

PDB ID : 3E3N / pdb_00003e3n
Title : The Glycogen phosphorylase b R state- AMP complex
Authors : Leonidas, D.D.; Zographos, S.E.; Oikonomakos, N.G.
Deposited on : 2008-08-07
Resolution : 2.70 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0
Mogul : 2022.3.0, CSD as543be (2022)
Xtriage (Phenix) : 2.0
EDS : 3.0
Buster-report : wwPDB partial adaption of 1.1.7 (2018)
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4 : 9.0.010 (Gargrove)
Density-Fitness : 1.0.12
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.70 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	3538 (2.70-2.70)
Clashscore	190562	3843 (2.70-2.70)
Ramachandran outliers	187476	3778 (2.70-2.70)
Sidechain outliers	187428	3778 (2.70-2.70)
RSRZ outliers	180081	3538 (2.70-2.70)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	842	2% 71% 21% . .
1	B	842	2% 70% 21% 5% .
1	C	842	74% 17% 5% 5%
1	D	842	74% 19% . .
1	E	842	% 72% 19% . 5%

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
1	F	842	% 72% 22%
1	G	842	2% 71% 20% 6%
1	H	842	3% 73% 19% 5%

2 Entry composition


There are 4 unique types of molecules in this entry. The entry contains 53512 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Glycogen phosphorylase, muscle form.


Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
1	A	814	Total	C	N	O	P	S	0	0	0
			6635	4225	1175	1204	1	30			
1	B	815	Total	C	N	O	P	S	0	0	0
			6642	4230	1176	1205	1	30			
1	C	802	Total	C	N	O	P	S	0	0	0
			6536	4165	1155	1186	1	29			
1	D	815	Total	C	N	O	P	S	0	0	0
			6643	4229	1176	1207	1	30			
1	E	803	Total	C	N	O	P	S	0	0	0
			6545	4170	1156	1189	1	29			
1	F	813	Total	C	N	O	P	S	0	0	0
			6626	4220	1173	1202	1	30			
1	G	814	Total	C	N	O	P	S	0	0	0
			6635	4225	1175	1204	1	30			
1	H	814	Total	C	N	O	P	S	0	0	0
			6636	4224	1175	1206	1	30			

- Molecule 2 is ADENOSINE MONOPHOSPHATE (CCD ID: AMP) (formula: C₁₀H₁₄N₅O₇P).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
2	A	1	Total	C	N	O	P	0	0
			23	10	5	7	1		
2	B	1	Total	C	N	O	P	0	0
			23	10	5	7	1		
2	C	1	Total	C	N	O	P	0	0
			23	10	5	7	1		
2	D	1	Total	C	N	O	P	0	0
			23	10	5	7	1		
2	E	1	Total	C	N	O	P	0	0
			23	10	5	7	1		
2	F	1	Total	C	N	O	P	0	0
			23	10	5	7	1		
2	G	1	Total	C	N	O	P	0	0
			23	10	5	7	1		
2	H	1	Total	C	N	O	P	0	0
			23	10	5	7	1		

- Molecule 3 is SULFATE ION (CCD ID: SO4) (formula: O₄S).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	A	1	Total	O	S	0	0
			5	4	1		
3	A	1	Total	O	S	0	0
			5	4	1		
3	B	1	Total	O	S	0	0
			5	4	1		
3	B	1	Total	O	S	0	0
			5	4	1		
3	C	1	Total	O	S	0	0
			5	4	1		
3	C	1	Total	O	S	0	0
			5	4	1		
3	D	1	Total	O	S	0	0
			5	4	1		
3	D	1	Total	O	S	0	0
			5	4	1		
3	E	1	Total	O	S	0	0
			5	4	1		
3	E	1	Total	O	S	0	0
			5	4	1		
3	F	1	Total	O	S	0	0
			5	4	1		
3	F	1	Total	O	S	0	0
			5	4	1		
3	G	1	Total	O	S	0	0
			5	4	1		
3	G	1	Total	O	S	0	0
			5	4	1		


Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	H	1	Total	O	S	0	0
			5	4	1		
3	H	1	Total	O	S	0	0
			5	4	1		


- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	56	Total	O	0	0
			56	56		
4	B	32	Total	O	0	0
			32	32		
4	C	36	Total	O	0	0
			36	36		
4	D	62	Total	O	0	0
			62	62		
4	E	27	Total	O	0	0
			27	27		
4	F	38	Total	O	0	0
			38	38		
4	G	69	Total	O	0	0
			69	69		
4	H	30	Total	O	0	0
			30	30		




• Molecule 1: Glycogen phosphorylase, muscle form


Chain C: 74% 17% 5% 5%




• Molecule 1: Glycogen phosphorylase, muscle form

Chain D: 74% 19% 7%







- Molecule 1: Glycogen phosphorylase, muscle form




- Molecule 1: Glycogen phosphorylase, muscle form







- Molecule 1: Glycogen phosphorylase, muscle form



4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	119.02Å 188.08Å 175.91Å 90.00° 108.92° 90.00°	Depositor
Resolution (Å)	29.79 – 2.70 29.79 – 2.70	Depositor EDS
% Data completeness (in resolution range)	100.0 (29.79-2.70) 94.3 (29.79-2.70)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	0.09	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.71 (at 2.68Å)	Xtriage
Refinement program	REFMAC 5.2.0005	Depositor
R, R_{free}	0.192 , 0.256 (Not available) , 0.265	Depositor DCC
R_{free} test set	9450 reflections (5.00%)	wwPDB-VP
Wilson B-factor (Å ²)	41.2	Xtriage
Anisotropy	0.029	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.30 , 44.0	EDS
L-test for twinning ²	$\langle L \rangle = 0.50$, $\langle L^2 \rangle = 0.34$	Xtriage
Estimated twinning fraction	0.000 for h,-k,-h-l	Xtriage
F_o, F_c correlation	0.94	EDS
Total number of atoms	53512	wwPDB-VP
Average B, all atoms (Å ²)	24.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The analyses of the Patterson function reveals a significant off-origin peak that is 74.27 % of the origin peak, indicating pseudo-translational symmetry. The chance of finding a peak of this or larger height randomly in a structure without pseudo-translational symmetry is equal to 1.5777e-06. The detected translational NCS is most likely also responsible for the elevated intensity ratio.*

¹ Intensities estimated from amplitudes.

² Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: LLP, SO4, AMP

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# $ Z > 5$	RMSZ	# $ Z > 5$
1	A	0.59	0/6756	0.91	3/9137 (0.0%)
1	B	0.56	0/6764	0.91	6/9149 (0.1%)
1	C	0.56	0/6654	0.89	3/8999 (0.0%)
1	D	0.59	1/6764 (0.0%)	0.91	4/9148 (0.0%)
1	E	0.56	1/6663 (0.0%)	0.88	7/9011 (0.1%)
1	F	0.58	0/6747	0.90	4/9125 (0.0%)
1	G	0.60	2/6756 (0.0%)	0.92	7/9137 (0.1%)
1	H	0.56	1/6756 (0.0%)	0.90	10/9136 (0.1%)
All	All	0.58	5/53860 (0.0%)	0.90	44/72842 (0.1%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	1
1	H	0	1
All	All	0	2

All (5) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	G	657	ILE	CA-CB	8.07	1.58	1.54
1	H	82	ILE	CA-CB	5.67	1.61	1.54
1	G	82	ILE	CA-CB	5.13	1.60	1.54
1	E	230	VAL	CA-CB	5.02	1.58	1.54
1	D	82	ILE	CA-CB	5.00	1.60	1.54

The worst 5 of 44 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	675	GLY	N-CA-C	-7.83	101.69	112.18
1	B	280	TYR	N-CA-C	7.49	124.13	108.39
1	C	380	LEU	CA-C-N	7.37	126.90	119.24
1	C	380	LEU	C-N-CA	7.37	126.90	119.24
1	H	8	GLU	N-CA-C	-6.80	103.28	112.26

There are no chirality outliers.

All (2) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	454	GLY	Peptide
1	H	7	GLN	Peptide

5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	6635	0	6598	115	0
1	B	6642	0	6605	128	0
1	C	6536	0	6500	94	0
1	D	6643	0	6602	90	0
1	E	6545	0	6506	104	0
1	F	6626	0	6590	97	0
1	G	6635	0	6598	126	0
1	H	6636	0	6595	111	0
2	A	23	0	12	1	0
2	B	23	0	12	2	0
2	C	23	0	12	0	0
2	D	23	0	12	0	0
2	E	23	0	12	0	0
2	F	23	0	12	0	0
2	G	23	0	12	1	0
2	H	23	0	12	1	0
3	A	10	0	0	0	0
3	B	10	0	0	0	0
3	C	10	0	0	0	0
3	D	10	0	0	0	0
3	E	10	0	0	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
3	F	10	0	0	1	0
3	G	10	0	0	1	0
3	H	10	0	0	0	0
4	A	56	0	0	1	0
4	B	32	0	0	1	0
4	C	36	0	0	3	0
4	D	62	0	0	1	0
4	E	27	0	0	2	0
4	F	38	0	0	0	0
4	G	69	0	0	1	0
4	H	30	0	0	3	0
All	All	53512	0	52690	842	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 8.

The worst 5 of 842 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:C:274:ASN:HA	1:C:277:ARG:HG3	1.34	1.09
1:H:615:MET:HE1	1:H:761:ILE:HG12	1.30	1.06
1:G:615:MET:HE1	1:G:761:ILE:HG12	1.43	1.01
1:G:274:ASN:HA	1:G:277:ARG:HG3	1.42	1.01
1:D:336:GLN:HE21	1:D:825:TRP:HE1	0.99	0.98

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	807/842 (96%)	771 (96%)	27 (3%)	9 (1%)	11 29

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	B	808/842 (96%)	757 (94%)	39 (5%)	12 (2%)	8	22
1	C	793/842 (94%)	745 (94%)	40 (5%)	8 (1%)	12	32
1	D	808/842 (96%)	763 (94%)	38 (5%)	7 (1%)	14	35
1	E	794/842 (94%)	752 (95%)	36 (4%)	6 (1%)	16	37
1	F	806/842 (96%)	759 (94%)	42 (5%)	5 (1%)	21	44
1	G	807/842 (96%)	763 (94%)	34 (4%)	10 (1%)	10	27
1	H	807/842 (96%)	756 (94%)	44 (6%)	7 (1%)	14	35
All	All	6430/6736 (96%)	6066 (94%)	300 (5%)	64 (1%)	12	32

5 of 64 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	16	ARG
1	A	166	PHE
1	A	835	PRO
1	B	16	ARG
1	B	273	GLU

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	703/730 (96%)	613 (87%)	90 (13%)	4	11
1	B	704/730 (96%)	616 (88%)	88 (12%)	4	11
1	C	692/730 (95%)	603 (87%)	89 (13%)	4	10
1	D	704/730 (96%)	607 (86%)	97 (14%)	3	9
1	E	693/730 (95%)	611 (88%)	82 (12%)	5	13
1	F	702/730 (96%)	605 (86%)	97 (14%)	3	9
1	G	703/730 (96%)	610 (87%)	93 (13%)	4	10
1	H	703/730 (96%)	620 (88%)	83 (12%)	5	13
All	All	5604/5840 (96%)	4885 (87%)	719 (13%)	4	11

5 of 719 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	F	177	GLU
1	G	274	ASN
1	F	279	LEU
1	F	165	ILE
1	F	649	ARG

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 203 such sidechains are listed below:

Mol	Chain	Res	Type
1	E	541	ASN
1	F	579	ASN
1	H	744	GLN
1	E	727	ASN
1	F	235	ASN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

8 non-standard protein/DNA/RNA residues are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
1	LLP	D	680	1	23,24,25	1.74	4 (17%)	25,32,34	1.46	8 (32%)
1	LLP	B	680	1	23,24,25	1.76	5 (21%)	25,32,34	1.39	5 (20%)
1	LLP	E	680	1	23,24,25	1.68	3 (13%)	25,32,34	1.58	6 (24%)
1	LLP	G	680	1	23,24,25	1.59	3 (13%)	25,32,34	1.53	5 (20%)
1	LLP	C	680	1	23,24,25	1.68	3 (13%)	25,32,34	1.40	4 (16%)
1	LLP	F	680	1	23,24,25	1.61	2 (8%)	25,32,34	1.43	5 (20%)

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
1	LLP	H	680	1	23,24,25	1.73	5 (21%)	25,32,34	1.28	5 (20%)
1	LLP	A	680	1	23,24,25	1.86	3 (13%)	25,32,34	1.42	5 (20%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
1	LLP	D	680	1	-	4/16/17/19	0/1/1/1
1	LLP	B	680	1	-	3/16/17/19	0/1/1/1
1	LLP	E	680	1	-	1/16/17/19	0/1/1/1
1	LLP	G	680	1	-	0/16/17/19	0/1/1/1
1	LLP	C	680	1	-	3/16/17/19	0/1/1/1
1	LLP	F	680	1	-	2/16/17/19	0/1/1/1
1	LLP	H	680	1	-	3/16/17/19	0/1/1/1
1	LLP	A	680	1	-	2/16/17/19	0/1/1/1

The worst 5 of 28 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	680	LLP	O3-C3	-6.30	1.22	1.36
1	C	680	LLP	O3-C3	-5.82	1.23	1.36
1	B	680	LLP	O3-C3	-5.82	1.23	1.36
1	H	680	LLP	O3-C3	-5.78	1.23	1.36
1	E	680	LLP	O3-C3	-5.66	1.23	1.36

The worst 5 of 43 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	E	680	LLP	OP4-C5'-C5	3.69	116.27	109.36
1	B	680	LLP	OP4-C5'-C5	3.40	115.74	109.36
1	G	680	LLP	CE-NZ-C4'	-3.16	108.61	118.72
1	C	680	LLP	C5-C6-N1	-3.02	118.93	123.83
1	G	680	LLP	C5-C6-N1	-2.98	118.98	123.83

There are no chirality outliers.

5 of 18 torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
1	A	680	LLP	C5'-OP4-P-OP2
1	A	680	LLP	C5'-OP4-P-OP3
1	B	680	LLP	C5'-OP4-P-OP1
1	B	680	LLP	C5'-OP4-P-OP2
1	B	680	LLP	C5'-OP4-P-OP3

There are no ring outliers.

8 monomers are involved in 26 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
1	D	680	LLP	1	0
1	B	680	LLP	2	0
1	E	680	LLP	4	0
1	G	680	LLP	6	0
1	C	680	LLP	4	0
1	F	680	LLP	3	0
1	H	680	LLP	3	0
1	A	680	LLP	3	0

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

24 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
3	SO4	D	845	-	4,4,4	0.21	0	6,6,6	0.19	0
3	SO4	D	844	-	4,4,4	0.25	0	6,6,6	0.18	0
2	AMP	H	843	-	25,25,25	1.48	5 (20%)	37,38,38	1.82	9 (24%)
3	SO4	F	845	-	4,4,4	0.24	0	6,6,6	0.21	0
3	SO4	H	844	-	4,4,4	0.22	0	6,6,6	0.17	0

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
2	AMP	A	843	-	25,25,25	1.47	4 (16%)	37,38,38	1.97	8 (21%)
3	SO4	F	844	-	4,4,4	0.26	0	6,6,6	0.23	0
2	AMP	D	843	-	25,25,25	1.53	4 (16%)	37,38,38	1.97	9 (24%)
3	SO4	A	844	-	4,4,4	0.23	0	6,6,6	0.20	0
3	SO4	B	844	-	4,4,4	0.25	0	6,6,6	0.13	0
3	SO4	C	844	-	4,4,4	0.26	0	6,6,6	0.24	0
3	SO4	E	845	-	4,4,4	0.25	0	6,6,6	0.08	0
3	SO4	E	844	-	4,4,4	0.22	0	6,6,6	0.24	0
2	AMP	B	843	-	25,25,25	1.45	3 (12%)	37,38,38	1.78	7 (18%)
2	AMP	G	843	-	25,25,25	1.40	4 (16%)	37,38,38	1.96	9 (24%)
3	SO4	A	845	-	4,4,4	0.24	0	6,6,6	0.17	0
3	SO4	B	845	-	4,4,4	0.23	0	6,6,6	0.18	0
3	SO4	C	845	-	4,4,4	0.24	0	6,6,6	0.19	0
3	SO4	H	899	-	4,4,4	0.25	0	6,6,6	0.13	0
2	AMP	C	843	-	25,25,25	1.45	4 (16%)	37,38,38	1.83	10 (27%)
2	AMP	E	843	-	25,25,25	1.49	4 (16%)	37,38,38	1.85	9 (24%)
3	SO4	G	845	-	4,4,4	0.24	0	6,6,6	0.10	0
2	AMP	F	843	-	25,25,25	1.41	3 (12%)	37,38,38	2.08	10 (27%)
3	SO4	G	844	-	4,4,4	0.25	0	6,6,6	0.10	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	AMP	H	843	-	-	0/10/26/26	0/3/3/3
2	AMP	D	843	-	-	0/10/26/26	0/3/3/3
2	AMP	A	843	-	-	0/10/26/26	0/3/3/3
2	AMP	G	843	-	-	0/10/26/26	0/3/3/3
2	AMP	F	843	-	-	0/10/26/26	0/3/3/3
2	AMP	C	843	-	-	0/10/26/26	0/3/3/3
2	AMP	E	843	-	-	0/10/26/26	0/3/3/3
2	AMP	B	843	-	-	0/10/26/26	0/3/3/3

The worst 5 of 31 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	B	843	AMP	C5-C4	5.00	1.48	1.39
2	H	843	AMP	C5-C4	5.00	1.48	1.39
2	D	843	AMP	C5-C4	4.98	1.48	1.39

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	E	843	AMP	C5-C4	4.97	1.48	1.39
2	C	843	AMP	C5-C4	4.95	1.47	1.39

The worst 5 of 71 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	F	843	AMP	C5-C4-N3	-6.22	118.16	126.72
2	A	843	AMP	C5-C4-N3	-6.21	118.17	126.72
2	D	843	AMP	C5-C4-N3	-5.96	118.51	126.72
2	G	843	AMP	C5-C4-N3	-5.47	119.19	126.72
2	E	843	AMP	C5-C4-N3	-5.27	119.46	126.72

There are no chirality outliers.



There are no torsion outliers.



There are no ring outliers.

6 monomers are involved in 7 short contacts:


Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	H	843	AMP	1	0
3	F	845	SO4	1	0
2	A	843	AMP	1	0
2	B	843	AMP	2	0
2	G	843	AMP	1	0
3	G	844	SO4	1	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.







Ligand AMP G 843



Ligand AMP C 843





5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	813/842 (96%)	-0.29	14 (1%) 69 67	7, 21, 42, 58	0
1	B	814/842 (96%)	-0.12	20 (2%) 58 55	4, 25, 50, 75	0
1	C	801/842 (95%)	-0.12	4 (0%) 87 86	3, 21, 41, 54	0
1	D	814/842 (96%)	-0.26	3 (0%) 88 87	6, 22, 38, 54	0
1	E	802/842 (95%)	-0.10	7 (0%) 81 80	5, 23, 43, 54	0
1	F	812/842 (96%)	-0.06	11 (1%) 73 72	3, 20, 38, 57	0
1	G	813/842 (96%)	-0.22	14 (1%) 69 67	7, 21, 41, 57	0
1	H	813/842 (96%)	0.05	23 (2%) 55 51	8, 28, 54, 69	0
All	All	6482/6736 (96%)	-0.14	96 (1%) 72 70	3, 23, 44, 75	0

The worst 5 of 96 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	F	164	GLY	5.0
1	D	288	GLY	4.9
1	A	288	GLY	4.7
1	H	380	LEU	4.7
1	H	165	ILE	4.1

6.2 Non-standard residues in protein, DNA, RNA chains ⓘ

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
1	LLP	F	680	24/25	0.95	0.07	15,17,22,23	0

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
1	LLP	C	680	24/25	0.96	0.06	13,16,23,23	0
1	LLP	D	680	24/25	0.96	0.07	16,17,21,24	0
1	LLP	E	680	24/25	0.96	0.07	16,19,22,23	0
1	LLP	B	680	24/25	0.96	0.07	19,27,31,32	0
1	LLP	G	680	24/25	0.96	0.07	17,18,21,23	0
1	LLP	H	680	24/25	0.96	0.07	22,25,27,27	0
1	LLP	A	680	24/25	0.97	0.06	17,19,22,24	0

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
3	SO4	C	844	5/5	0.73	0.14	88,88,88,88	0
3	SO4	E	845	5/5	0.77	0.11	95,95,96,96	0
3	SO4	H	899	5/5	0.79	0.13	80,80,80,81	0
3	SO4	B	845	5/5	0.81	0.11	90,90,90,91	0
3	SO4	F	845	5/5	0.83	0.13	86,86,87,87	0
3	SO4	G	845	5/5	0.84	0.13	82,83,83,84	0
3	SO4	A	845	5/5	0.84	0.12	97,98,98,98	0
3	SO4	F	844	5/5	0.86	0.12	58,59,59,60	0
3	SO4	D	845	5/5	0.86	0.11	83,84,84,85	0
3	SO4	E	844	5/5	0.86	0.09	71,72,73,73	0
3	SO4	C	845	5/5	0.86	0.12	58,59,60,60	0
3	SO4	D	844	5/5	0.91	0.07	66,66,66,67	0
3	SO4	B	844	5/5	0.91	0.10	67,67,68,68	0
2	AMP	D	843	23/23	0.95	0.07	27,29,31,31	0
2	AMP	E	843	23/23	0.95	0.07	33,37,39,39	0
3	SO4	A	844	5/5	0.95	0.07	47,47,48,48	0
3	SO4	H	844	5/5	0.95	0.09	46,46,46,47	0
2	AMP	C	843	23/23	0.95	0.07	28,32,34,34	0
3	SO4	G	844	5/5	0.96	0.07	40,41,42,42	0
2	AMP	B	843	23/23	0.96	0.06	26,35,36,36	0
2	AMP	A	843	23/23	0.96	0.06	25,27,29,29	0

Continued on next page...




Continued from previous page...

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
2	AMP	G	843	23/23	0.96	0.06	23,28,30,30	0
2	AMP	F	843	23/23	0.97	0.06	20,26,27,27	0
2	AMP	H	843	23/23	0.97	0.07	28,34,34,36	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.




Electron density around AMP D 843:

2mF_o-DF_c (at 0.7 rmsd) in gray
mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around AMP E 843:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around AMP C 843:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around AMP B 843:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around AMP A 843:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around AMP G 843:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around AMP F 843:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



Electron density around AMP H 843:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



6.5 Other polymers [i](#)

There are no such residues in this entry.