



# Full wwPDB X-ray Structure Validation Report ⓘ

Mar 5, 2026 – 08:43 PM UTC

PDB ID : 4E4P / pdb\_00004e4p  
Title : Second native structure of Xylanase A1 from Paenibacillus sp. JDR-2  
Authors : Pozharski, E.; St John, F.J.  
Deposited on : 2012-03-13  
Resolution : 1.92 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0  
Xtrriage (Phenix) : 2.0  
EDS : 3.0  
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)  
CCP4 : 9.0.010 (Gargrove)  
Density-Fitness : 1.0.12  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.49


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 1.92 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	180053	1188 (1.92-1.92)
Clashscore	190562	1209 (1.92-1.92)
Ramachandran outliers	187476	1195 (1.92-1.92)
Sidechain outliers	187428	1195 (1.92-1.92)
RSRZ outliers	180081	1188 (1.92-1.92)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	341	 6% 70% 14% • 14%
1	B	341	 6% 70% 14% • 14%

## 2 Entry composition [i](#)

There are 4 unique types of molecules in this entry. The entry contains 4878 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Endo-1,4-beta-xylanase A.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	293	2322	1468	400	445	9	0	0	0
1	B	293	2322	1468	400	445	9	0	0	0

There are 14 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	1	GLY	-	expression tag	UNP C6CRV0
A	2	SER	-	expression tag	UNP C6CRV0
A	3	HIS	-	expression tag	UNP C6CRV0
A	4	MET	-	expression tag	UNP C6CRV0
A	339	ALA	-	expression tag	UNP C6CRV0
A	340	GLU	-	expression tag	UNP C6CRV0
A	341	GLN	-	expression tag	UNP C6CRV0
B	1	GLY	-	expression tag	UNP C6CRV0
B	2	SER	-	expression tag	UNP C6CRV0
B	3	HIS	-	expression tag	UNP C6CRV0
B	4	MET	-	expression tag	UNP C6CRV0
B	339	ALA	-	expression tag	UNP C6CRV0
B	340	GLU	-	expression tag	UNP C6CRV0
B	341	GLN	-	expression tag	UNP C6CRV0

- Molecule 2 is MAGNESIUM ION (CCD ID: MG) (formula: Mg).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	A	2	Total	Mg	0	0
			2	2		
2	B	2	Total	Mg	0	0
			2	2		

- Molecule 3 is CHLORIDE ION (CCD ID: CL) (formula: Cl).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	A	1	Total Cl 2 2	0	1
3	B	1	Total Cl 2 2	0	1


- Molecule 4 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
4	A	116	Total O 117 117	0	1
4	B	109	Total O 109 109	0	0


### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Endo-1,4-beta-xylanase A



- Molecule 1: Endo-1,4-beta-xylanase A



## 4 Data and refinement statistics i

Property	Value	Source
Space group	C 1 2 1	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	179.52Å 58.53Å 65.02Å 90.00° 108.81° 90.00°	Depositor
Resolution (Å)	50.00 – 1.92 50.00 – 1.92	Depositor EDS
% Data completeness (in resolution range)	98.7 (50.00-1.92) 98.7 (50.00-1.92)	Depositor EDS
$R_{merge}$	0.11	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	1.17 (at 1.92Å)	Xtrriage
Refinement program	REFMAC 5.5.0102	Depositor
R, $R_{free}$	0.253 , 0.295 0.269 , 0.303	Depositor DCC
$R_{free}$ test set	2440 reflections (4.98%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	30.1	Xtrriage
Anisotropy	0.436	Xtrriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.32 , 35.4	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.42$ , $\langle L^2 \rangle = 0.25$	Xtrriage
Estimated twinning fraction	0.046 for -h-2*1,-k,l	Xtrriage
$F_o, F_c$ correlation	0.96	EDS
Total number of atoms	4878	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	44.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The analyses of the Patterson function reveals a significant off-origin peak that is 99.77 % of the origin peak, indicating pseudo-translational symmetry. The chance of finding a peak of this or larger height randomly in a structure without pseudo-translational symmetry is equal to 2.5868e-14. The detected translational NCS is most likely also responsible for the elevated intensity ratio.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: CL, MG

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	A	1.34	7/2374 (0.3%)	1.16	6/3226 (0.2%)
1	B	1.33	3/2374 (0.1%)	1.17	7/3226 (0.2%)
All	All	1.33	10/4748 (0.2%)	1.17	13/6452 (0.2%)

All (10) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	B	193	ASP	N-CA	7.95	1.55	1.45
1	A	299	ILE	C-O	-7.92	1.15	1.24
1	A	174	ALA	C-O	5.83	1.30	1.24
1	A	193	ASP	N-CA	5.65	1.52	1.45
1	A	230	GLY	C-O	5.24	1.30	1.23
1	A	191	TYR	CA-C	5.17	1.58	1.52
1	A	212	ILE	C-O	5.14	1.29	1.24
1	B	81	HIS	CA-C	5.11	1.58	1.52
1	B	212	ILE	CA-CB	-5.03	1.48	1.54
1	A	81	HIS	CA-C	5.02	1.58	1.52

All (13) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	213	ASN	N-CA-C	6.45	118.31	111.28
1	B	52	LEU	CA-C-N	-6.05	116.75	122.37
1	B	52	LEU	C-N-CA	-6.05	116.75	122.37
1	B	51	ALA	N-CA-C	-6.04	105.74	113.23
1	B	39	ASP	CB-CA-C	-5.90	99.16	109.07
1	B	46	ALA	N-CA-C	5.60	119.21	112.38
1	A	104	VAL	N-CA-C	-5.50	104.68	109.19
1	A	143	ASN	CB-CA-C	5.45	115.07	111.20
1	A	39	ASP	CB-CA-C	-5.38	99.50	109.29

*Continued on next page...*

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	25	ASP	CA-C-N	-5.18	115.67	124.31
1	A	25	ASP	C-N-CA	-5.18	115.67	124.31
1	B	25	ASP	CA-C-N	-5.04	116.35	125.66
1	B	25	ASP	C-N-CA	-5.04	116.35	125.66

There are no chirality outliers.

There are no planarity outliers.

## 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2322	0	2245	27	2
1	B	2322	0	2245	29	2
2	A	2	0	0	0	0
2	B	2	0	0	0	0
3	A	2	0	0	0	0
3	B	2	0	0	0	0
4	A	117	0	0	0	0
4	B	109	0	0	1	0
All	All	4878	0	4490	56	2

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 6.

All (56) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:196:GLU:HG3	1:A:231:MET:HE2	1.61	0.80
1:A:82:VAL:HG21	1:A:137:ASN:HB2	1.62	0.80
1:B:82:VAL:HG21	1:B:137:ASN:HB2	1.64	0.79
1:B:293:LYS:NZ	1:B:293:LYS:HB2	2.02	0.75
1:A:293:LYS:HB2	1:A:293:LYS:NZ	2.04	0.73
1:B:293:LYS:NZ	1:B:293:LYS:CB	2.55	0.69
1:B:34:LEU:HD12	1:B:76:MET:HE1	1.76	0.68
1:A:25:ASP:O	1:A:26:LEU:HB2	1.96	0.66

Continued on next page...

*Continued from previous page...*

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:32:GLU:OE2	1:A:35:LYS:HE3	1.95	0.66
1:B:293:LYS:HB2	1:B:293:LYS:HZ2	1.61	0.64
1:A:19:ASN:HB2	1:A:38:HIS:CD2	2.33	0.63
1:A:284:LEU:HD22	1:A:288:LEU:HG	1.81	0.63
1:B:25:ASP:O	1:B:26:LEU:HB2	1.99	0.63
1:B:19:ASN:HB2	1:B:38:HIS:CD2	2.36	0.60
1:B:293:LYS:CB	1:B:293:LYS:HZ3	2.14	0.60
1:A:293:LYS:NZ	1:A:293:LYS:CB	2.66	0.58
1:A:7:LEU:H	1:A:7:LEU:HD22	1.69	0.58
1:A:34:LEU:HD12	1:A:76:MET:HE1	1.86	0.57
1:A:293:LYS:HB2	1:A:293:LYS:HZ2	1.69	0.57
1:A:43:ALA:CB	1:A:67:MET:HE1	2.36	0.56
1:B:284:LEU:HD22	1:B:288:LEU:HG	1.86	0.56
1:A:293:LYS:HB2	1:A:293:LYS:HZ3	1.71	0.54
1:B:196:GLU:HG3	1:B:231:MET:HE2	1.90	0.53
1:B:32:GLU:OE2	1:B:35:LYS:HD3	2.10	0.52
1:B:173:LEU:O	1:B:177:GLU:HG3	2.09	0.51
1:A:258:VAL:O	1:A:299:ILE:HA	2.10	0.51
1:A:43:ALA:HB2	1:A:67:MET:HE1	1.94	0.51
1:B:7:LEU:HD22	1:B:7:LEU:H	1.75	0.50
1:B:43:ALA:HB2	1:B:67:MET:HE1	1.94	0.50
1:B:43:ALA:CB	1:B:67:MET:HE1	2.42	0.50
1:B:240:ASN:OD1	1:B:241:PRO:HD2	2.12	0.48
1:A:147:PRO:HB2	1:A:203:THR:HG22	1.95	0.48
1:A:68:ILE:CD1	1:A:129:LYS:HB3	2.43	0.48
1:B:155:ARG:HD2	1:B:157:THR:HG22	1.96	0.47
1:B:25:ASP:O	1:B:26:LEU:CB	2.62	0.47
1:B:212:ILE:CG2	1:B:225:LEU:HD23	2.44	0.47
1:B:233:GLY:HA2	1:B:235:TYR:CZ	2.51	0.46
1:B:68:ILE:HG12	1:B:78:MET:HE3	1.98	0.45
1:B:169:GLU:CD	1:B:215:ARG:HH22	2.24	0.45
1:B:223:LYS:NZ	4:B:514:HOH:O	2.50	0.44
1:A:240:ASN:OD1	1:A:241:PRO:HD2	2.17	0.44
1:A:26:LEU:HA	1:A:31:LEU:HG	1.99	0.44
1:B:23:ALA:O	1:B:25:ASP:O	2.34	0.44
1:A:293:LYS:CB	1:A:293:LYS:HZ3	2.28	0.44
1:A:155:ARG:HD2	1:A:157:THR:HG22	2.00	0.43
1:A:23:ALA:O	1:A:25:ASP:O	2.36	0.43
1:A:45:ASN:HA	1:A:48:LYS:HD2	2.00	0.43
1:B:142:ASP:O	1:B:143:ASN:C	2.61	0.43
1:B:26:LEU:HA	1:B:31:LEU:HG	2.00	0.43

*Continued on next page...*

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:293:LYS:HZ3	1:B:293:LYS:HB3	1.84	0.42
1:A:277:LEU:HB3	1:A:279:VAL:HG12	2.02	0.41
1:A:233:GLY:HA2	1:A:235:TYR:CZ	2.55	0.41
1:B:24:GLU:C	1:B:25:ASP:O	2.63	0.41
1:A:68:ILE:HG12	1:A:78:MET:HE3	2.03	0.41
1:A:112:LEU:O	1:A:116:ARG:HG3	2.21	0.41
1:B:216:TYR:CG	1:B:225:LEU:HB2	2.56	0.41

All (2) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:11:TYR:OH	1:B:283:TYR:OH[4_546]	1.98	0.22
1:A:283:TYR:OH	1:B:11:TYR:OH[4_546]	2.11	0.09

### 5.3 Torsion angles [i](#)

#### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	289/341 (85%)	281 (97%)	8 (3%)	0	100	100
1	B	289/341 (85%)	282 (98%)	7 (2%)	0	100	100
All	All	578/682 (85%)	563 (97%)	15 (3%)	0	100	100

There are no Ramachandran outliers to report.

#### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	247/288 (86%)	236 (96%)	11 (4%)	24	9
1	B	247/288 (86%)	235 (95%)	12 (5%)	22	8
All	All	494/576 (86%)	471 (95%)	23 (5%)	23	9

All (23) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	27	GLU
1	A	29	THR
1	A	31	LEU
1	A	33	LEU
1	A	34	LEU
1	A	68	ILE
1	A	250	LYS
1	A	277	LEU
1	A	284	LEU
1	A	290	LYS
1	A	293	LYS
1	B	27	GLU
1	B	30	ARG
1	B	31	LEU
1	B	33	LEU
1	B	34	LEU
1	B	68	ILE
1	B	250	LYS
1	B	277	LEU
1	B	284	LEU
1	B	287	GLN
1	B	290	LYS
1	B	293	LYS

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (9) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	37	HIS
1	A	94	ASN
1	A	102	ASN
1	A	170	GLN

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type
1	A	298	HIS
1	B	37	HIS
1	B	102	ASN
1	B	220	HIS
1	B	295	HIS

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

### 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

### 5.6 Ligand geometry [i](#)

Of 8 ligands modelled in this entry, 8 are monoatomic - leaving 0 for Mogul analysis.

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

### 5.7 Other polymers [i](#)

There are no such residues in this entry.

### 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data

### 6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	293/341 (85%)	0.70	20 (6%) 23 28	25, 41, 82, 107	0
1	B	293/341 (85%)	0.65	19 (6%) 25 29	26, 42, 79, 107	0
All	All	586/682 (85%)	0.68	39 (6%) 24 28	25, 41, 80, 107	0

All (39) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	277	LEU	6.0
1	B	277	LEU	5.3
1	B	99	ASP	4.4
1	A	29	THR	3.8
1	B	5	ALA	3.8
1	A	99	ASP	3.6
1	A	5	ALA	3.4
1	A	98	ASP	3.3
1	B	283	TYR	3.0
1	A	33	LEU	2.9
1	A	283	TYR	2.9
1	B	29	THR	2.8
1	B	98	ASP	2.8
1	A	284	LEU	2.7
1	A	14	ASP	2.6
1	B	27	GLU	2.6
1	A	27	GLU	2.6
1	B	33	LEU	2.6
1	B	148	ALA	2.5
1	A	112	LEU	2.4
1	B	26	LEU	2.4
1	A	69	ASP	2.4
1	B	284	LEU	2.4
1	B	69	ASP	2.3

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	RSRZ
1	A	28	GLY	2.3
1	B	180	ASP	2.3
1	B	14	ASP	2.3
1	B	113	ASP	2.3
1	A	32	GLU	2.3
1	A	19	ASN	2.3
1	A	65	ASP	2.3
1	A	36	MET	2.2
1	A	39	ASP	2.2
1	B	65	ASP	2.2
1	B	19	ASN	2.2
1	B	36	MET	2.2
1	A	180	ASP	2.2
1	A	148	ALA	2.2
1	B	28	GLY	2.0

## 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

## 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å <sup>2</sup> )	Q<0.9
2	MG	A	401	1/1	0.76	0.13	38,38,38,38	0
2	MG	A	403	1/1	0.86	0.10	43,43,43,43	0
2	MG	B	403	1/1	0.86	0.10	41,41,41,41	0
2	MG	B	401	1/1	0.95	0.08	41,41,41,41	0
3	CL	A	402[A]	1/1	0.96	0.09	33,33,33,33	1
3	CL	A	402[B]	1/1	0.96	0.09	30,30,30,30	1
3	CL	B	402[A]	1/1	0.98	0.11	33,33,33,33	1
3	CL	B	402[B]	1/1	0.98	0.11	35,35,35,35	1

## 6.5 Other polymers [i](#)

There are no such residues in this entry.