



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 5, 2026 – 08:48 PM UTC

PDB ID : 5E4N / pdb_00005e4n
Title : 3-Deoxy-D-arabino-heptulosonate 7-phosphate synthase from Mycobacterium tuberculosis with D-tyrosine bound in the tyrosine and phenylalanine binding sites
Authors : Reichau, S.; Jiao, W.; Parker, E.J.
Deposited on : 2015-10-06
Resolution : 2.05 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0
Mogul : 2022.3.0, CSD as543be (2022)
Xtriage (Phenix) : 2.0
EDS : 3.0
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4 : 9.0.010 (Gargrove)
Density-Fitness : 1.0.12
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.05 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	2260 (2.04-2.04)
Clashscore	190562	2333 (2.04-2.04)
Ramachandran outliers	187476	2318 (2.04-2.04)
Sidechain outliers	187428	2318 (2.04-2.04)
RSRZ outliers	180081	2260 (2.04-2.04)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	464	10% 88% 11% .
1	B	464	8% 89% 9% .

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
2	SO4	B	503	-	-	X	-

2 Entry composition [i](#)

There are 6 unique types of molecules in this entry. The entry contains 7744 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called 3-deoxy-D-arabinoheptulosonate-7-phosphate synthase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	459	Total	C	N	O	S	0	10	0
			3604	2255	660	669	20			
1	B	454	Total	C	N	O	S	0	7	0
			3536	2211	647	658	20			

There are 4 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	-1	GLY	-	expression tag	UNP A0A0E8NFD1
A	0	ALA	-	expression tag	UNP A0A0E8NFD1
B	-1	GLY	-	expression tag	UNP A0A0E8NFD1
B	0	ALA	-	expression tag	UNP A0A0E8NFD1

- Molecule 2 is SULFATE ION (CCD ID: SO4) (formula: O₄S).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
2	A	1	Total O S 5 4 1	0	0
2	B	1	Total O S 5 4 1	0	0
2	B	1	Total O S 5 4 1	0	0
2	B	1	Total O S 5 4 1	0	0


- Molecule 3 is CHLORIDE ION (CCD ID: CL) (formula: Cl).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	A	2	Total Cl 2 2	0	0
3	B	2	Total Cl 2 2	0	0

- Molecule 4 is MANGANESE (II) ION (CCD ID: MN) (formula: Mn).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
4	A	1	Total Mn 1 1	0	0
4	B	1	Total Mn 1 1	0	0

- Molecule 5 is D-TYROSINE (CCD ID: DTY) (formula: C₉H₁₁NO₃).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
5	A	1	Total	C	N	O	0	0
			13	9	1	3		
5	B	1	Total	C	N	O	0	0
			13	9	1	3		
5	B	1	Total	C	N	O	0	0
			13	9	1	3		


- Molecule 6 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
6	A	249	Total	O	0	0
			249	249		
6	B	290	Total	O	0	0
			290	290		


3 Residue-property plots

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: 3-deoxy-D-arabinoheptulosonate-7-phosphate synthase



- Molecule 1: 3-deoxy-D-arabinoheptulosonate-7-phosphate synthase



4 Data and refinement statistics

Property	Value	Source
Space group	P 32 2 1	Depositor
Cell constants a, b, c, α , β , γ	205.74Å 205.74Å 66.64Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	47.37 – 2.05 47.37 – 2.05	Depositor EDS
% Data completeness (in resolution range)	100.0 (47.37-2.05) 100.0 (47.37-2.05)	Depositor EDS
R_{merge}	0.17	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	3.39 (at 2.05Å)	Xtriage
Refinement program	REFMAC 5.7.0029	Depositor
R, R_{free}	0.199 , 0.229 0.179 , 0.201	Depositor DCC
R_{free} test set	5048 reflections (4.99%)	wwPDB-VP
Wilson B-factor (Å ²)	25.5	Xtriage
Anisotropy	0.115	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.36 , 40.8	EDS
L-test for twinning ²	$\langle L \rangle = 0.45$, $\langle L^2 \rangle = 0.27$	Xtriage
Estimated twinning fraction	0.046 for -h,-k,l	Xtriage
F_o, F_c correlation	0.95	EDS
Total number of atoms	7744	wwPDB-VP
Average B, all atoms (Å ²)	32.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 2.72% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality ⓘ

5.1 Standard geometry ⓘ

Bond lengths and bond angles in the following residue types are not validated in this section: SO4, DTY, CL, MN

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.74	1/3690 (0.0%)	0.85	0/5017
1	B	0.79	0/3617	0.90	6/4916 (0.1%)
All	All	0.76	1/7307 (0.0%)	0.88	6/9933 (0.1%)

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	16	PRO	CA-C	5.37	1.54	1.51

All (6) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	338	MET	N-CA-C	6.86	121.81	113.17
1	B	15	LEU	CA-C-N	-5.50	114.71	120.38
1	B	15	LEU	C-N-CA	-5.50	114.71	120.38
1	B	348	PRO	CA-C-N	-5.24	114.27	119.56
1	B	348	PRO	C-N-CA	-5.24	114.27	119.56
1	B	348	PRO	N-CA-C	5.04	116.85	110.70

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3604	0	3572	45	0
1	B	3536	0	3491	42	0
2	A	5	0	0	0	0
2	B	15	0	0	3	0
3	A	2	0	0	0	0
3	B	2	0	0	0	0
4	A	1	0	0	0	0
4	B	1	0	0	0	0
5	A	13	0	9	0	0
5	B	26	0	19	5	0
6	A	249	0	0	2	0
6	B	290	0	0	4	0
All	All	7744	0	7091	80	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 6.

All (80) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:372:THR:HG22	1:B:382:ARG:HE	1.33	0.93
1:B:48[A]:MET:HE3	1:B:170:VAL:HG23	1.52	0.92
1:B:17:PRO:HA	2:B:503:SO4:O1	1.75	0.86
1:B:23:ARG:NH2	2:B:503:SO4:O2	2.14	0.81
1:B:48[A]:MET:HE2	1:B:167:SER:HA	1.63	0.79
1:A:12:LEU:HD11	5:B:507:DTY:O	1.84	0.78
1:A:48[A]:MET:HE3	1:A:170:VAL:HG23	1.66	0.76
1:A:38:PRO:HG2	1:A:40:TRP:CZ2	2.21	0.76
1:A:10:ASP:HB3	1:B:171:ARG:HH22	1.56	0.71
1:A:391:VAL:HG12	1:A:459[A]:MET:HE1	1.72	0.69
1:B:49[A]:ARG:NH1	6:B:601:HOH:O	2.25	0.69
1:B:337:ARG:HH21	1:B:369:HIS:HB3	1.60	0.65
1:B:311[A]:MET:HE3	1:B:316:ALA:N	2.11	0.65
1:B:100:ARG:HD3	6:B:609:HOH:O	1.96	0.64
1:A:48[A]:MET:HE2	1:A:167:SER:HA	1.81	0.63
1:A:48[A]:MET:CE	1:A:170:VAL:HG23	2.29	0.62
1:B:48[A]:MET:HE3	1:B:170:VAL:CG2	2.27	0.62
1:B:48[A]:MET:CE	1:B:167:SER:HA	2.30	0.61
1:A:138:ASP:HA	6:A:651:HOH:O	2.01	0.60
1:A:10:ASP:CB	1:B:171:ARG:HH22	2.14	0.59
1:B:337:ARG:O	1:B:337:ARG:HG3	2.01	0.59
1:B:372:THR:CG2	1:B:382:ARG:HE	2.11	0.59

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:12:LEU:HD13	1:B:92:MET:HA	1.83	0.59
1:A:268:GLU:HB3	1:A:269:PRO:CD	2.33	0.58
1:A:48[A]:MET:CE	1:A:167:SER:HA	2.34	0.57
1:A:12:LEU:CD1	5:B:507:DTY:H	2.17	0.56
1:A:68[B]:GLN:HG2	1:A:302:PRO:HG3	1.89	0.55
1:A:395:PHE:CD1	1:A:459[A]:MET:CE	2.91	0.54
1:B:381:THR:HA	1:B:442:PRO:HG3	1.89	0.54
1:B:197:VAL:HA	1:B:200:TRP:CE3	2.42	0.54
1:B:145:ARG:NH1	6:B:605:HOH:O	2.40	0.54
1:B:18:LEU:N	2:B:503:SO4:O1	2.39	0.53
1:A:381:THR:HA	1:A:442:PRO:HG3	1.90	0.53
1:A:133:LYS:NZ	1:A:440:CYS:SG	2.76	0.53
1:B:83:GLN:HA	1:B:124:VAL:O	2.10	0.52
1:A:64:ILE:O	1:A:68[B]:GLN:HG3	2.10	0.51
1:A:338:MET:O	1:A:342:LYS:HB2	2.10	0.51
1:B:372:THR:HG21	6:B:812:HOH:O	2.10	0.51
1:A:9:ILE:HG12	1:B:3:TRP:HD1	1.77	0.50
1:A:83:GLN:HA	1:A:124:VAL:O	2.11	0.50
1:B:311[A]:MET:HE3	1:B:316:ALA:CA	2.41	0.50
1:B:281[A]:ILE:HD11	1:B:296:ALA:HB2	1.93	0.49
1:A:41:PRO:HG2	1:A:44:GLN:OE1	2.13	0.49
1:A:48[A]:MET:CE	1:A:170:VAL:CG2	2.90	0.49
1:A:96[A]:GLU:HB3	1:A:97:PRO:HD3	1.94	0.49
1:A:48[A]:MET:HE3	1:A:170:VAL:CG2	2.39	0.48
1:A:10:ASP:HB3	1:B:171:ARG:NH2	2.25	0.48
1:A:77:GLY:HA2	1:A:404:HIS:CD2	2.49	0.48
1:B:399[A]:ARG:HH11	1:B:399[A]:ARG:HG2	1.79	0.48
1:A:337:ARG:O	1:A:337:ARG:HG3	2.15	0.47
1:B:53:GLU:OE2	5:B:508:DTY:N	2.38	0.47
1:A:6:ASP:OD1	1:B:4:THR:HG22	2.15	0.47
1:B:382:ARG:HD2	1:B:442:PRO:HG2	1.96	0.47
1:A:38:PRO:HB2	1:A:40:TRP:CE2	2.51	0.45
1:B:311[A]:MET:CE	1:B:316:ALA:N	2.79	0.45
1:A:9:ILE:HG21	1:A:51:VAL:HG21	1.98	0.45
1:A:129:GLY:HA2	1:A:131:TYR:CE2	2.52	0.45
1:B:338:MET:O	1:B:342:LYS:HB2	2.17	0.45
1:A:382:ARG:HD2	1:A:442:PRO:HG2	1.98	0.44
1:A:268:GLU:HB3	1:A:269:PRO:HD2	2.00	0.44
1:B:52:LEU:O	1:B:55:VAL:HG22	2.18	0.43
1:B:48[A]:MET:CE	1:B:170:VAL:CG2	2.96	0.43
1:A:39:THR:O	1:A:39:THR:OG1	2.33	0.43

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:281[A]:ILE:HD11	1:A:296:ALA:HB2	2.00	0.43
6:A:845:HOH:O	1:B:191:LEU:HD13	2.18	0.43
1:A:263:ASP:OD1	1:A:264:GLY:N	2.52	0.42
1:B:38:PRO:HG2	1:B:40:TRP:CZ2	2.54	0.42
1:A:413:THR:OG1	1:A:415:GLU:HG2	2.20	0.42
1:B:372:THR:HG22	1:B:382:ARG:NE	2.16	0.42
1:B:23:ARG:HH22	5:B:508:DTY:HB2	1.83	0.42
1:A:9:ILE:CG2	1:A:51:VAL:HG21	2.50	0.42
1:A:12:LEU:HD11	5:B:507:DTY:H	1.86	0.41
1:B:311[A]:MET:CE	1:B:316:ALA:CA	2.98	0.41
1:A:10:ASP:CB	1:B:171:ARG:NH2	2.81	0.41
1:A:262:SER:OG	1:A:269:PRO:HA	2.20	0.41
1:B:44:GLN:O	1:B:48[A]:MET:HG2	2.21	0.41
1:A:259:LEU:HD13	1:A:271:LEU:HD13	2.03	0.41
1:A:96[B]:GLU:HB3	1:A:97:PRO:HD3	2.03	0.41
1:B:441:ASP:HB3	1:B:442:PRO:HD2	2.03	0.40
1:A:395:PHE:CD1	1:A:459[A]:MET:HE2	2.57	0.40

There are no symmetry-related clashes.

5.3 Torsion angles ⓘ

5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	465/464 (100%)	450 (97%)	15 (3%)	0	100	100
1	B	455/464 (98%)	442 (97%)	12 (3%)	1 (0%)	43	37
All	All	920/928 (99%)	892 (97%)	27 (3%)	1 (0%)	48	43

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	B	440	CYS

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	379/376 (101%)	373 (98%)	6 (2%)	55	56
1	B	370/376 (98%)	368 (100%)	2 (0%)	81	85
All	All	749/752 (100%)	741 (99%)	8 (1%)	65	68

All (8) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	4	THR
1	A	40	TRP
1	A	368	MET
1	A	374	GLU
1	A	399	ARG
1	A	430	THR
1	B	40	TRP
1	B	450	LEU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (1) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	392	GLN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 13 ligands modelled in this entry, 6 are monoatomic - leaving 7 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
2	SO4	B	502	-	4,4,4	0.45	0	6,6,6	0.27	0
2	SO4	A	501	-	4,4,4	0.10	0	6,6,6	0.48	0
2	SO4	B	501	-	4,4,4	0.26	0	6,6,6	0.31	0
5	DTY	B	508	-	12,13,13	0.91	0	13,17,17	1.02	1 (7%)
2	SO4	B	503	-	4,4,4	0.28	0	6,6,6	0.82	0
5	DTY	B	507	-	12,13,13	0.75	0	13,17,17	1.12	0
5	DTY	A	505	-	12,13,13	1.34	2 (16%)	13,17,17	0.95	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
5	DTY	B	507	-	-	2/8/8/8	0/1/1/1
5	DTY	A	505	-	-	3/8/8/8	0/1/1/1
5	DTY	B	508	-	-	5/8/8/8	0/1/1/1

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
5	A	505	DTY	OXT-C	-3.41	1.19	1.30
5	A	505	DTY	O-C	2.23	1.28	1.22

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
5	B	508	DTY	CB-CA-N	2.32	120.29	111.40

There are no chirality outliers.

All (10) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
5	A	505	DTY	N-CA-CB-CG
5	A	505	DTY	C-CA-CB-CG
5	B	507	DTY	N-CA-CB-CG
5	B	508	DTY	O-C-CA-N
5	B	508	DTY	N-CA-CB-CG
5	B	508	DTY	OXT-C-CA-N
5	B	507	DTY	C-CA-CB-CG
5	B	508	DTY	C-CA-CB-CG
5	B	508	DTY	OXT-C-CA-CB
5	A	505	DTY	CA-CB-CG-CD1

There are no ring outliers.

3 monomers are involved in 8 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
5	B	508	DTY	2	0
2	B	503	SO4	3	0
5	B	507	DTY	3	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2		OWAB(Å ²)	Q<0.9
1	A	459/464 (98%)	0.18	47 (10%)	12 12	10, 30, 71, 103	10 (2%)
1	B	454/464 (97%)	-0.01	37 (8%)	18 18	10, 24, 63, 100	7 (1%)
All	All	913/928 (98%)	0.09	84 (9%)	14 14	10, 27, 64, 103	17 (1%)

All (84) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	267	GLY	8.0
1	B	15	LEU	7.7
1	A	3	TRP	7.0
1	A	264	GLY	5.7
1	A	235	ASP	5.6
1	B	236	ARG	5.5
1	A	239	GLN	5.3
1	A	238	LEU	5.1
1	B	240	THR	5.1
1	A	13	PRO	5.1
1	B	195	HIS	5.0
1	B	239	GLN	5.0
1	B	238	LEU	4.8
1	A	195	HIS	4.8
1	B	1	MET	4.7
1	A	269	PRO	4.5
1	B	9	ILE	4.5
1	A	240	THR	4.4
1	B	237	ASN	4.4
1	A	234	ALA	4.3
1	B	373	HIS	4.2
1	A	12	LEU	4.2
1	B	377	THR	4.1
1	A	268	GLU	4.1

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	15	LEU	4.0
1	B	234	ALA	3.9
1	B	200	TRP	3.9
1	A	1	MET	3.9
1	B	379	PHE	3.8
1	A	14	SER	3.7
1	A	462	ASP	3.7
1	A	237	ASN	3.5
1	B	197	VAL	3.4
1	B	16	PRO	3.4
1	B	376	SER	3.4
1	B	263	ASP	3.3
1	A	338	MET	3.3
1	B	372	THR	3.3
1	A	197	VAL	3.3
1	A	379	PHE	3.3
1	B	269	PRO	3.2
1	A	11	GLN	3.2
1	A	200	TRP	3.1
1	A	137	ALA	3.0
1	B	341[A]	HIS	2.9
1	A	337	ARG	2.9
1	B	268	GLU	2.9
1	B	433	ALA	2.9
1	A	9	ILE	2.9
1	A	136	SER	2.8
1	A	263	ASP	2.8
1	A	2	ASN	2.7
1	A	373	HIS	2.7
1	B	235	ASP	2.7
1	A	377	THR	2.7
1	A	236	ARG	2.7
1	A	40	TRP	2.7
1	A	10	ASP	2.6
1	A	378	GLY	2.6
1	A	202	ARG	2.6
1	A	262	SER	2.4
1	A	376	SER	2.4
1	A	241	ALA	2.4
1	A	374	GLU	2.4
1	A	207	THR	2.4
1	B	337	ARG	2.4

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	341[A]	HIS	2.4
1	B	338	MET	2.3
1	B	378	GLY	2.3
1	B	207	THR	2.3
1	A	192	ALA	2.2
1	B	462	ASP	2.2
1	B	198	HIS	2.2
1	B	201	ASN	2.2
1	B	8	PRO	2.1
1	A	135	ARG	2.1
1	B	206	ARG	2.1
1	A	196	LEU	2.1
1	B	136	SER	2.1
1	A	399	ARG	2.0
1	B	213	ARG	2.0
1	A	383	HIS	2.0
1	A	375	SER	2.0
1	B	190	GLY	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
3	CL	A	504	1/1	0.86	0.14	62,62,62,62	0
5	DTY	B	507	13/13	0.88	0.12	28,31,38,44	0
2	SO4	B	502	5/5	0.92	0.09	68,68,72,75	0
5	DTY	B	508	13/13	0.92	0.11	25,33,36,37	0
3	CL	A	502	1/1	0.93	0.10	61,61,61,61	0

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
3	CL	B	506	1/1	0.93	0.09	61,61,61,61	0
2	SO4	B	503	5/5	0.94	0.15	34,34,37,38	5
5	DTY	A	505	13/13	0.96	0.07	20,23,24,24	0
4	MN	B	505	1/1	0.99	0.08	32,32,32,32	1
2	SO4	B	501	5/5	0.99	0.05	26,27,29,30	0
3	CL	B	504	1/1	0.99	0.04	39,39,39,39	0
2	SO4	A	501	5/5	0.99	0.05	28,30,30,33	0
4	MN	A	503	1/1	1.00	0.04	29,29,29,29	1

6.5 Other polymers [i](#)

There are no such residues in this entry.