



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 5, 2026 – 04:29 AM UTC

PDB ID : 2E52 / pdb_00002e52
Title : Crystal structural analysis of HindIII restriction endonuclease in complex with cognate DNA at 2.0 angstrom resolution
Authors : Watanabe, N.; Sato, C.; Tanaka, I.
Deposited on : 2006-12-18
Resolution : 2.00 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0
Mogul : 2022.3.0, CSD as543be (2022)
Xtriage (Phenix) : 2.0
EDS : 3.0
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4 : 9.0.010 (Gargrove)
Density-Fitness : 1.0.12
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.00 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.




Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	10052 (2.00-2.00)
Clashscore	190562	11152 (2.00-2.00)
Ramachandran outliers	187476	11031 (2.00-2.00)
Sidechain outliers	187428	11029 (2.00-2.00)
RSRZ outliers	180081	10067 (2.00-2.00)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	300	92% 7%
1	B	300	4% 90% 9% ..
1	C	300	4% 89% 9% ..
1	D	300	2% 88% 10% ...
2	E	12	100%

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
2	F	12	 92% 8%
2	G	12	 83% 8% 8%
2	H	12	 100%
2	I	12	 67% 92% 8%
2	J	12	 67% 58% 25% 17%

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
4	ACT	C	2003	-	-	X	-

2 Entry composition

There are 5 unique types of molecules in this entry. The entry contains 12288 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Type II restriction enzyme HindIII.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	299	Total	C	N	O	S	0	0	0
			2461	1586	407	465	3			
1	C	299	Total	C	N	O	S	0	0	0
			2461	1586	407	465	3			
1	B	298	Total	C	N	O	S	0	0	0
			2452	1580	405	464	3			
1	D	298	Total	C	N	O	S	0	0	0
			2452	1580	405	464	3			

- Molecule 2 is a DNA chain called DNA (5'-D(*DGP*DCP*DCP*DAP*DAP*DGP*DCP*DTP*DTP*DGP*DGP*DC)-3').


Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	E	12	Total	C	N	O	P	0	0	0
			243	116	46	70	11			
2	F	12	Total	C	N	O	P	0	0	0
			243	116	46	70	11			
2	G	12	Total	C	N	O	P	0	0	0
			243	116	46	70	11			
2	H	12	Total	C	N	O	P	0	0	0
			243	116	46	70	11			
2	I	12	Total	C	N	O	P	0	0	0
			243	116	46	70	11			
2	J	12	Total	C	N	O	P	0	0	0
			243	116	46	70	11			

- Molecule 3 is GLYCEROL (CCD ID: GOL) (formula: C₃H₈O₃).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	A	1	Total	C	O	0	0
			6	3	3		
3	C	1	Total	C	O	0	0
			6	3	3		
3	C	1	Total	C	O	0	0
			6	3	3		
3	E	1	Total	C	O	0	0
			6	3	3		

- Molecule 4 is ACETATE ION (CCD ID: ACT) (formula: $C_2H_3O_2$).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
4	C	1	Total C O 4 2 2	0	0
4	C	1	Total C O 4 2 2	0	0
4	B	1	Total C O 4 2 2	0	0

- Molecule 5 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
5	A	224	Total O 224 224	0	0
5	C	192	Total O 192 192	0	0
5	B	216	Total O 216 216	0	0
5	D	188	Total O 188 188	0	0
5	E	37	Total O 37 37	0	0
5	F	36	Total O 36 36	0	0
5	G	35	Total O 35 35	0	0
5	H	39	Total O 39 39	0	0
5	I	1	Total O 1 1	0	0



3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Type II restriction enzyme HindIII




- Molecule 1: Type II restriction enzyme HindIII



- Molecule 1: Type II restriction enzyme HindIII



- Molecule 1: Type II restriction enzyme HindIII



- Molecule 2: DNA (5'-D(*DGP*DCP*DCP*DAP*DAP*DGP*DCP*DTP*DTP*DGP*DGP*D C)-3')

Chain E:  100%

There are no outlier residues recorded for this chain.

- Molecule 2: DNA (5'-D(*DGP*DCP*DCP*DAP*DAP*DGP*DCP*DTP*DTP*DGP*DGP*D C)-3')

Chain F:  92% 8%




- Molecule 2: DNA (5'-D(*DGP*DCP*DCP*DAP*DAP*DGP*DCP*DTP*DTP*DGP*DGP*D C)-3')

Chain G:  83% 8% 8%




- Molecule 2: DNA (5'-D(*DGP*DCP*DCP*DAP*DAP*DGP*DCP*DTP*DTP*DGP*DGP*D C)-3')

Chain H:  100%

There are no outlier residues recorded for this chain.


- Molecule 2: DNA (5'-D(*DGP*DCP*DCP*DAP*DAP*DGP*DCP*DTP*DTP*DGP*DGP*D C)-3')

Chain I:  67% 92% 8%



- Molecule 2: DNA (5'-D(*DGP*DCP*DCP*DAP*DAP*DGP*DCP*DTP*DTP*DGP*DGP*D C)-3')

Chain J:  67% 58% 25% 17%



4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	83.12Å 131.53Å 93.72Å 90.00° 111.20° 90.00°	Depositor
Resolution (Å)	39.68 – 2.00 39.68 – 2.00	Depositor EDS
% Data completeness (in resolution range)	99.9 (39.68-2.00) 99.8 (39.68-2.00)	Depositor EDS
R_{merge}	0.06	Depositor
R_{sym}	0.06	Depositor
$\langle I/\sigma(I) \rangle$ ¹	10.22 (at 2.00Å)	Xtriage
Refinement program	REFMAC 5.2.0019	Depositor
R, R_{free}	0.175 , 0.217 0.172 , 0.216	Depositor DCC
R_{free} test set	6327 reflections (5.01%)	wwPDB-VP
Wilson B-factor (Å ²)	23.2	Xtriage
Anisotropy	0.034	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.35 , 38.1	EDS
L-test for twinning ²	$\langle L \rangle = 0.50$, $\langle L^2 \rangle = 0.33$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.96	EDS
Total number of atoms	12288	wwPDB-VP
Average B, all atoms (Å ²)	23.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.18% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: GOL, ACT

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.84	0/2503	0.85	0/3362
1	B	0.86	0/2494	0.90	2/3351 (0.1%)
1	C	0.84	0/2503	0.91	5/3362 (0.1%)
1	D	0.85	0/2494	0.93	6/3351 (0.2%)
2	E	0.59	0/272	1.21	0/418
2	F	0.62	0/272	1.27	1/418 (0.2%)
2	G	0.65	0/272	1.32	2/418 (0.5%)
2	H	0.60	0/272	1.13	0/418
2	I	0.26	0/272	1.30	3/418 (0.7%)
2	J	0.30	0/272	1.08	2/418 (0.5%)
All	All	0.81	0/11626	0.95	21/15934 (0.1%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	D	0	1

There are no bond length outliers.

All (21) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	I	6	DG	O3'-P-O5'	-14.90	81.65	104.00
1	D	263	THR	N-CA-C	-8.91	95.59	109.76
2	I	6	DG	OP1-P-O3'	-7.41	85.76	108.00
2	G	4	DA	N9-C1'-C2'	6.88	123.83	113.50
1	C	116	ARG	NE-CZ-NH2	-6.62	113.24	119.20
1	D	264	ARG	CA-C-N	6.49	133.39	121.70

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	D	264	ARG	C-N-CA	6.49	133.39	121.70
1	C	261	ASN	CA-C-N	6.13	132.74	121.70
1	C	261	ASN	C-N-CA	6.13	132.74	121.70
2	J	3	DC	P-O3'-C3'	6.01	129.22	120.20
1	B	83	LYS	CA-C-N	-5.86	114.23	120.03
1	B	83	LYS	C-N-CA	-5.86	114.23	120.03
1	D	263	THR	CA-C-N	5.81	132.16	121.70
1	D	263	THR	C-N-CA	5.81	132.16	121.70
2	I	6	DG	OP2-P-O3'	-5.77	90.69	108.00
1	C	116	ARG	NE-CZ-NH1	5.74	127.23	121.50
1	D	268	ILE	N-CA-C	-5.69	105.19	110.53
2	F	4	DA	N9-C1'-C2'	5.68	122.02	113.50
1	C	262	PHE	N-CA-C	5.17	125.47	111.00
2	J	5	DA	P-O3'-C3'	5.17	127.95	120.20
2	G	9	DT	O4'-C1'-N1	-5.15	100.68	108.40

There are no chirality outliers.

All (1) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	D	262	PHE	Peptide

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2461	0	2518	15	0
1	B	2452	0	2502	16	0
1	C	2461	0	2518	31	0
1	D	2452	0	2502	27	0
2	E	243	0	136	0	0
2	F	243	0	136	0	0
2	G	243	0	136	1	0
2	H	243	0	136	0	0
2	I	243	0	136	0	0
2	J	243	0	136	3	0
3	A	6	0	8	1	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
3	C	12	0	16	1	0
3	E	6	0	8	0	0
4	B	4	0	3	0	0
4	C	8	0	6	3	0
5	A	224	0	0	2	0
5	B	216	0	0	4	0
5	C	192	0	0	3	0
5	D	188	0	0	1	0
5	E	37	0	0	1	0
5	F	36	0	0	1	0
5	G	35	0	0	0	0
5	H	39	0	0	0	0
5	I	1	0	0	0	0
All	All	12288	0	10897	84	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 4.

All (84) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:D:264:ARG:CB	1:D:265:GLU:HB2	1.69	1.23
1:D:264:ARG:HB3	1:D:265:GLU:CB	1.73	1.18
1:A:21:LYS:HE2	1:A:22:GLN:HE22	1.15	1.10
1:D:263:THR:O	1:D:266:GLU:HB2	1.54	1.04
1:C:264:ARG:HB2	1:D:264:ARG:HA	1.41	0.99
1:A:21:LYS:HE2	1:A:22:GLN:NE2	1.81	0.94
1:C:277:MET:HE1	1:D:259:ILE:HD12	1.64	0.80
2:J:3:DC:H2'	2:J:4:DA:C8	2.20	0.77
1:C:264:ARG:CB	1:D:264:ARG:HA	2.16	0.75
1:C:224:LYS:HG2	4:C:2003:ACT:H3	1.67	0.75
1:D:264:ARG:HB3	1:D:265:GLU:HB2	0.82	0.71
1:D:278:SER:O	1:D:282:GLU:HG3	1.92	0.69
1:C:262:PHE:HA	1:C:266:GLU:OE2	1.92	0.69
1:B:86:GLU:H	1:B:86:GLU:CD	2.01	0.69
1:D:263:THR:O	1:D:266:GLU:CB	2.36	0.69
1:D:21:LYS:HG3	5:F:40:HOH:O	1.94	0.67
1:C:265:GLU:O	1:C:268:ILE:HD13	1.96	0.66
1:C:261:ASN:ND2	1:C:261:ASN:O	2.25	0.66
1:B:229:THR:O	1:B:232:GLN:HG3	1.95	0.66
1:C:259:ILE:O	1:C:262:PHE:HB2	1.97	0.64

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:83:LYS:HE2	1:A:95:ILE:HD11	1.81	0.63
1:B:217:LYS:HE3	5:B:2027:HOH:O	2.00	0.61
1:A:54:LYS:HD2	1:A:55:PRO:HD2	1.82	0.61
1:C:264:ARG:HA	1:D:264:ARG:O	2.02	0.60
1:D:264:ARG:HD3	1:D:264:ARG:C	2.27	0.59
1:C:277:MET:CE	1:D:259:ILE:HD12	2.35	0.57
1:C:264:ARG:HG3	1:C:265:GLU:N	2.15	0.56
1:A:21:LYS:CE	1:A:22:GLN:HE22	2.04	0.56
1:C:110:LYS:HG2	5:C:2162:HOH:O	2.05	0.56
1:B:217:LYS:HD2	1:B:217:LYS:C	2.31	0.55
1:C:262:PHE:H	1:C:262:PHE:HD1	1.54	0.54
1:C:262:PHE:HD1	1:C:262:PHE:N	2.05	0.54
1:C:299:LEU:HD21	1:D:230:LEU:HD23	1.90	0.54
1:C:95:ILE:HD12	1:C:136:LYS:HD3	1.90	0.53
1:C:256:ILE:O	1:C:260:LYS:HG2	2.09	0.53
1:B:46:ILE:HG12	1:B:67:GLY:HA2	1.91	0.52
1:C:262:PHE:N	1:C:262:PHE:CD1	2.77	0.52
1:B:129:LEU:HA	1:B:132:TRP:CE3	2.46	0.51
1:C:272:LEU:HD23	1:C:277:MET:CE	2.41	0.50
1:C:299:LEU:O	1:D:213:ARG:NH1	2.44	0.50
1:B:21:LYS:HB2	5:E:1035:HOH:O	2.10	0.50
2:J:7:DC:H2'	2:J:8:DT:C6	2.47	0.50
1:B:132:TRP:HZ3	5:B:2080:HOH:O	1.95	0.49
1:C:186:PHE:CE1	4:C:2003:ACT:H2	2.47	0.49
1:D:264:ARG:CA	1:D:265:GLU:HB2	2.37	0.49
1:A:179:ASP:HB3	5:A:1147:HOH:O	2.12	0.49
1:D:229:THR:O	1:D:232:GLN:HG3	2.11	0.49
1:D:264:ARG:H	1:D:267:ALA:H	1.61	0.49
1:B:84:PRO:HD2	5:B:2164:HOH:O	2.13	0.48
1:B:86:GLU:CD	1:B:86:GLU:N	2.71	0.48
1:A:213:ARG:HG2	1:A:217:LYS:HE3	1.96	0.48
1:D:263:THR:O	1:D:266:GLU:N	2.48	0.47
1:A:163:ASN:HA	1:A:195:PRO:HB2	1.97	0.47
1:A:177:GLN:HE21	1:A:233:LEU:HD22	1.80	0.47
1:C:46:ILE:HG12	1:C:67:GLY:HA2	1.97	0.47
1:D:246:LEU:O	1:D:250:GLU:HG3	2.16	0.46
2:J:4:DA:H2''	2:J:5:DA:C8	2.50	0.46
1:C:224:LYS:HE2	5:C:2192:HOH:O	2.16	0.46
1:A:129:LEU:HA	1:A:132:TRP:CE3	2.50	0.45
1:A:46:ILE:HG12	1:A:67:GLY:HA2	1.99	0.45
1:D:13:GLU:O	1:D:16:THR:HG23	2.17	0.45

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:255:GLN:O	1:B:259:ILE:HD13	2.17	0.44
1:C:298:TYR:CE1	3:C:1002:GOL:H11	2.52	0.44
1:B:24:THR:HG22	1:B:65:LYS:HG2	1.99	0.44
1:D:143:ALA:O	1:D:168:SER:HA	2.18	0.44
1:C:162:GLU:OE2	5:C:2188:HOH:O	2.21	0.43
1:A:85:LEU:HD11	1:A:95:ILE:HG23	2.01	0.43
1:C:264:ARG:HB3	1:D:264:ARG:HG2	2.01	0.43
1:D:163:ASN:HA	1:D:195:PRO:HB2	2.02	0.42
1:D:46:ILE:HG12	1:D:67:GLY:HA2	2.02	0.42
1:C:11:LEU:O	1:C:15:LEU:HG	2.19	0.42
1:C:79:ASN:OD1	1:C:99:LYS:HE2	2.20	0.42
1:D:42:LEU:HD12	1:D:42:LEU:HA	1.96	0.41
1:B:89:GLY:HA2	2:G:4:DA:H5'	2.02	0.41
1:C:256:ILE:O	1:C:260:LYS:CG	2.69	0.41
1:A:110:LYS:HG2	5:A:1046:HOH:O	2.20	0.41
1:D:213:ARG:NH2	5:D:477:HOH:O	2.54	0.41
1:B:143:ALA:O	1:B:168:SER:HA	2.20	0.41
3:A:1004:GOL:H11	1:B:298:TYR:CE2	2.56	0.41
1:A:21:LYS:CE	1:A:22:GLN:NE2	2.69	0.41
1:C:224:LYS:CG	4:C:2003:ACT:H3	2.43	0.40
1:C:105:LEU:HD12	1:C:105:LEU:C	2.47	0.40
1:A:54:LYS:HD2	1:A:55:PRO:CD	2.47	0.40
1:B:217:LYS:HG3	5:B:2027:HOH:O	2.20	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	297/300 (99%)	290 (98%)	7 (2%)	0	100	100
1	B	296/300 (99%)	289 (98%)	7 (2%)	0	100	100

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	C	297/300 (99%)	286 (96%)	10 (3%)	1 (0%)	36	35
1	D	296/300 (99%)	286 (97%)	8 (3%)	2 (1%)	18	14
All	All	1186/1200 (99%)	1151 (97%)	32 (3%)	3 (0%)	36	35

All (3) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	D	264	ARG
1	D	265	GLU
1	C	262	PHE

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	275/276 (100%)	272 (99%)	3 (1%)	65	73
1	B	274/276 (99%)	263 (96%)	11 (4%)	28	27
1	C	275/276 (100%)	266 (97%)	9 (3%)	33	34
1	D	274/276 (99%)	264 (96%)	10 (4%)	31	31
All	All	1098/1104 (100%)	1065 (97%)	33 (3%)	36	38

All (33) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	86	GLU
1	A	211	PHE
1	A	261	ASN
1	C	19	GLU
1	C	131	GLU
1	C	211	PHE
1	C	235	GLN
1	C	260	LYS
1	C	261	ASN

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	C	262	PHE
1	C	265	GLU
1	C	268	ILE
1	B	2	LYS
1	B	14	ASN
1	B	34	LEU
1	B	86	GLU
1	B	131	GLU
1	B	184	ASN
1	B	201	LYS
1	B	211	PHE
1	B	217	LYS
1	B	231	ASN
1	B	299	LEU
1	D	19	GLU
1	D	25	ASN
1	D	42	LEU
1	D	87	GLN
1	D	110	LYS
1	D	201	LYS
1	D	205	SER
1	D	211	PHE
1	D	264	ARG
1	D	265	GLU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (29) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	17	ASN
1	A	18	GLN
1	A	22	GLN
1	A	37	ASN
1	A	198	GLN
1	A	231	ASN
1	A	239	ASN
1	C	87	GLN
1	C	102	ASN
1	C	163	ASN
1	C	177	GLN
1	C	190	GLN
1	C	193	ASN
1	C	198	GLN

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	C	235	GLN
1	C	239	ASN
1	B	81	ASN
1	B	163	ASN
1	B	184	ASN
1	B	190	GLN
1	B	193	ASN
1	B	209	ASN
1	B	231	ASN
1	B	232	GLN
1	D	18	GLN
1	D	22	GLN
1	D	37	ASN
1	D	81	ASN
1	D	255	GLN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no oligosaccharides in this entry.

5.6 Ligand geometry ⓘ

7 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
3	GOL	C	1002	-	5,5,5	0.84	0	5,5,5	0.68	0
3	GOL	C	1003	-	5,5,5	0.79	0	5,5,5	0.80	0
4	ACT	C	2003	-	3,3,3	0.87	0	3,3,3	1.17	0
4	ACT	B	2002	-	3,3,3	0.77	0	3,3,3	1.27	0
4	ACT	C	2001	-	3,3,3	1.02	0	3,3,3	1.30	0
3	GOL	A	1004	-	5,5,5	0.75	0	5,5,5	0.71	0
3	GOL	E	1001	-	5,5,5	0.43	0	5,5,5	0.53	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	GOL	C	1003	-	-	2/4/4/4	-
3	GOL	E	1001	-	-	0/4/4/4	-
3	GOL	A	1004	-	-	0/4/4/4	-
3	GOL	C	1002	-	-	0/4/4/4	-

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

All (2) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
3	C	1003	GOL	C1-C2-C3-O3
3	C	1003	GOL	O2-C2-C3-O3

There are no ring outliers.

3 monomers are involved in 5 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	C	1002	GOL	1	0
4	C	2003	ACT	3	0
3	A	1004	GOL	1	0

5.7 Other polymers

There are no such residues in this entry.

5.8 Polymer linkage issues ⓘ

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ > 2	OWAB(Å ²)	Q < 0.9
1	A	299/300 (99%)	-0.32	3 (1%) 79 79	11, 20, 32, 41	0
1	B	298/300 (99%)	-0.07	12 (4%) 42 41	10, 23, 35, 45	0
1	C	299/300 (99%)	-0.05	13 (4%) 40 39	11, 23, 43, 53	1 (0%)
1	D	298/300 (99%)	-0.23	6 (2%) 65 64	12, 21, 36, 48	0
2	E	12/12 (100%)	-1.22	0 100 100	12, 14, 21, 29	0
2	F	12/12 (100%)	-1.25	0 100 100	14, 15, 20, 25	0
2	G	12/12 (100%)	-1.22	0 100 100	12, 14, 21, 24	0
2	H	12/12 (100%)	-1.21	0 100 100	13, 14, 21, 29	0
2	I	12/12 (100%)	2.33	8 (66%) 0 0	22, 32, 36, 37	12 (100%)
2	J	12/12 (100%)	2.70	8 (66%) 0 0	20, 33, 39, 39	12 (100%)
All	All	1266/1272 (99%)	-0.16	50 (3%) 43 42	10, 22, 37, 53	25 (1%)

All (50) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	C	268	ILE	5.2
1	C	264	ARG	4.9
1	C	263	THR	4.4
2	J	10	DG	4.3
1	D	265	GLU	4.3
2	J	6	DG	4.0
1	B	16	THR	3.9
1	D	263	THR	3.7
2	J	9	DT	3.6
1	B	91	ALA	3.4
1	B	87	GLN	3.2
2	I	7	DC	3.2
2	I	10	DG	3.2

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
2	J	7	DC	3.1
2	I	6	DG	3.1
1	D	17	ASN	3.0
1	C	272	LEU	3.0
2	J	4	DA	3.0
1	D	264	ARG	2.9
2	J	5	DA	2.9
1	D	16	THR	2.8
1	B	14	ASN	2.8
1	B	89	GLY	2.8
1	A	1	LYS	2.7
1	B	84	PRO	2.7
1	A	17	ASN	2.6
2	I	9	DT	2.6
1	A	86	GLU	2.5
1	B	95	ILE	2.5
1	C	265	GLU	2.5
1	C	262	PHE	2.4
1	B	92	GLY	2.4
1	B	83	LYS	2.4
1	C	267	ALA	2.3
2	I	1	DG	2.3
2	I	5	DA	2.3
1	C	17	ASN	2.3
1	B	2	LYS	2.3
1	B	217	LYS	2.3
2	I	4	DA	2.2
1	C	32	TYR	2.2
1	D	267	ALA	2.2
1	C	232	GLN	2.2
2	J	3	DC	2.1
1	B	299	LEU	2.1
1	C	95	ILE	2.1
1	C	258	ILE	2.1
2	I	12	DC	2.1
1	C	257	ASN	2.0
2	J	12	DC	2.0

6.2 Non-standard residues in protein, DNA, RNA chains

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
4	ACT	C	2001	4/4	0.49	0.26	49,49,49,49	0
4	ACT	B	2002	4/4	0.62	0.25	46,46,47,47	0
4	ACT	C	2003	4/4	0.80	0.15	50,50,50,50	0
3	GOL	C	1003	6/6	0.93	0.12	22,26,27,29	0
3	GOL	E	1001	6/6	0.94	0.10	21,21,24,27	0
3	GOL	C	1002	6/6	0.94	0.11	22,23,25,25	0
3	GOL	A	1004	6/6	0.95	0.11	21,23,24,25	0

6.5 Other polymers [i](#)

There are no such residues in this entry.