



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 5, 2026 – 07:45 AM UTC

PDB ID : 3E51 / pdb_00003e51
Title : Crystal structure of HCV NS5B polymerase with a novel pyridazinone inhibitor
Authors : Han, Q.; Showalter, R.E.; Zhao, Q.; Kissinger, C.R.
Deposited on : 2008-08-12
Resolution : 1.90 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Buster-report	:	wwPDB partial adaption of 1.1.7 (2018)
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.90 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	7789 (1.90-1.90)
Clashscore	190562	8410 (1.90-1.90)
Ramachandran outliers	187476	8333 (1.90-1.90)
Sidechain outliers	187428	8333 (1.90-1.90)
RSRZ outliers	180081	7790 (1.90-1.90)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	578	 6% 89% 7% . .
1	B	578	 6% 90% 6% .

2 Entry composition

There are 3 unique types of molecules in this entry. The entry contains 9210 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called RNA-directed RNA polymerase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	557	Total	C	N	O	S	0	0	0
			4334	2735	764	804	31			
1	B	557	Total	C	N	O	S	0	0	0
			4334	2735	764	804	31			

There are 18 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	544	GLN	ARG	engineered mutation	UNP P26663
A	571	LEU	-	expression tag	UNP P26663
A	572	GLU	-	expression tag	UNP P26663
A	573	HIS	-	expression tag	UNP P26663
A	574	HIS	-	expression tag	UNP P26663
A	575	HIS	-	expression tag	UNP P26663
A	576	HIS	-	expression tag	UNP P26663
A	577	HIS	-	expression tag	UNP P26663
A	578	HIS	-	expression tag	UNP P26663
B	544	GLN	ARG	engineered mutation	UNP P26663
B	571	LEU	-	expression tag	UNP P26663
B	572	GLU	-	expression tag	UNP P26663
B	573	HIS	-	expression tag	UNP P26663
B	574	HIS	-	expression tag	UNP P26663
B	575	HIS	-	expression tag	UNP P26663
B	576	HIS	-	expression tag	UNP P26663
B	577	HIS	-	expression tag	UNP P26663
B	578	HIS	-	expression tag	UNP P26663

- Molecule 2 is N-{3-[5-hydroxy-2-(3-methylbutyl)-3-oxo-6-pyrrolidin-1-yl]-2,3-dihydropyridazin-4-yl]-1,1-dioxido-2H-1,2,4-benzothiadiazin-7-yl}methanesulfonamide (CCD ID: N35) (formula: C₂₁H₂₈N₆O₆S₂).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
2	A	1	Total	C	N	O	S	0	0
			35	21	6	6	2		
2	B	1	Total	C	N	O	S	0	0
			35	21	6	6	2		


- Molecule 3 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	236	Total	O	0	0
			236	236		
3	B	236	Total	O	0	0
			236	236		


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: RNA-directed RNA polymerase



• Molecule 1: RNA-directed RNA polymerase



4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	85.75Å 106.38Å 126.85Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	45.22 – 1.90 45.22 – 1.90	Depositor EDS
% Data completeness (in resolution range)	97.5 (45.22-1.90) 97.5 (45.22-1.90)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.36 (at 1.91Å)	Xtriage
Refinement program	REFMAC 5.2.0019	Depositor
R, R_{free}	0.222 , 0.271 0.225 , 0.273	Depositor DCC
R_{free} test set	4402 reflections (4.93%)	wwPDB-VP
Wilson B-factor (Å ²)	30.1	Xtriage
Anisotropy	0.457	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.35 , 33.6	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.33$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.96	EDS
Total number of atoms	9210	wwPDB-VP
Average B, all atoms (Å ²)	41.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The analyses of the Patterson function reveals a significant off-origin peak that is 51.62 % of the origin peak, indicating pseudo-translational symmetry. The chance of finding a peak of this or larger height randomly in a structure without pseudo-translational symmetry is equal to 5.4696e-05. The detected translational NCS is most likely also responsible for the elevated intensity ratio.*

¹ Intensities estimated from amplitudes.

² Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: N35

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.60	0/4428	0.81	0/6009
1	B	0.60	0/4428	0.82	0/6009
All	All	0.60	0/8856	0.82	0/12018

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	4334	0	4354	24	0
1	B	4334	0	4354	21	0
2	A	35	0	28	1	0
2	B	35	0	28	1	0
3	A	236	0	0	4	0
3	B	236	0	0	6	0
All	All	9210	0	8764	46	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 3.

All (46) close contacts within the same asymmetric unit are listed below, sorted by their clash

magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:331:GLU:HG2	3:B:750:HOH:O	1.92	0.69
1:B:12:THR:HG21	1:B:270:LYS:HE2	1.76	0.68
1:A:82:LEU:HD13	1:A:249:ALA:HB2	1.78	0.64
1:A:13:PRO:HG3	1:A:43:ARG:HH12	1.65	0.61
1:A:95:HIS:H	1:A:95:HIS:CD2	2.20	0.59
1:A:187:MET:HE1	1:A:292:THR:CG2	2.35	0.57
1:A:329:THR:HG23	3:A:795:HOH:O	2.04	0.57
1:A:172:LYS:HE3	1:A:560:ILE:HD13	1.87	0.57
1:A:187:MET:HE1	1:A:292:THR:HG22	1.87	0.57
1:A:308:LEU:HB3	1:A:311:CYS:SG	2.46	0.56
1:B:160:ILE:HA	1:B:282:SER:OG	2.06	0.56
1:B:445:CYS:SG	1:B:454:ILE:HD12	2.46	0.54
1:A:13:PRO:HG3	1:A:43:ARG:NH1	2.22	0.54
1:B:308:LEU:HB2	1:B:311:CYS:SG	2.50	0.52
1:B:227:THR:HB	1:B:347:SER:O	2.11	0.51
1:B:236:GLU:OE2	1:B:280:ARG:NH2	2.25	0.50
1:B:491:LYS:HE2	3:B:815:HOH:O	2.12	0.50
1:A:381:VAL:HG11	1:A:474:LEU:CD2	2.42	0.49
2:B:579:N35:N22	2:B:579:N35:O28	2.44	0.49
1:A:346:TYR:O	1:A:347:SER:HB3	2.13	0.48
1:A:22:PRO:HG2	1:A:401:ARG:HG3	1.96	0.48
1:B:31:LEU:HD11	1:B:492:LEU:HD22	1.96	0.47
1:A:451:CYS:HB2	3:A:813:HOH:O	2.14	0.47
1:A:523:LYS:HG3	1:A:534:LEU:HD12	1.96	0.46
1:A:11:ILE:HD13	1:A:159:LEU:HD22	1.97	0.46
1:B:346:TYR:O	1:B:347:SER:HB3	2.15	0.45
1:A:154:ARG:N	3:A:805:HOH:O	2.51	0.44
1:B:488:CYS:HB2	3:B:800:HOH:O	2.18	0.44
1:B:451:CYS:SG	3:B:804:HOH:O	2.61	0.43
1:B:273:ASN:ND2	1:B:275:GLY:H	2.17	0.43
1:A:455:GLU:HB3	3:A:770:HOH:O	2.18	0.43
1:B:544:GLN:HB3	3:B:812:HOH:O	2.18	0.43
1:B:182:LEU:C	1:B:182:LEU:HD23	2.44	0.43
1:A:419:LEU:C	1:A:419:LEU:HD23	2.44	0.42
1:A:182:LEU:HD11	1:A:239:ILE:HG22	1.99	0.42
1:B:22:PRO:HG2	1:B:401:ARG:HG3	2.00	0.42
1:B:86:GLU:O	1:B:90:LYS:HG2	2.20	0.42
1:A:90:LYS:HE2	1:A:111:LEU:HD11	2.01	0.42
2:A:579:N35:O28	2:A:579:N35:N22	2.51	0.42
1:A:374:HIS:CD2	1:A:476:SER:HB2	2.55	0.41
1:A:77:THR:O	1:B:24:ASN:HB2	2.21	0.41

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:409:LEU:HD21	1:A:454:ILE:HB	2.03	0.41
1:B:100:LYS:HG3	3:B:813:HOH:O	2.20	0.41
1:A:419:LEU:HD21	1:A:497:LEU:CD1	2.51	0.41
1:B:527:ASN:O	1:B:533:LYS:HE2	2.21	0.40
1:B:85:VAL:HG11	1:B:116:VAL:HG13	2.04	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	553/578 (96%)	542 (98%)	11 (2%)	0	100	100
1	B	553/578 (96%)	544 (98%)	9 (2%)	0	100	100
All	All	1106/1156 (96%)	1086 (98%)	20 (2%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	475/493 (96%)	465 (98%)	10 (2%)	47	44
1	B	475/493 (96%)	469 (99%)	6 (1%)	61	61
All	All	950/986 (96%)	934 (98%)	16 (2%)	53	52

All (16) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	82	LEU
1	A	110	ASN
1	A	148	GLN
1	A	236	GLU
1	A	308	LEU
1	A	361	GLU
1	A	381	VAL
1	A	419	LEU
1	A	514	GLN
1	A	533	LYS
1	B	106	LYS
1	B	159	LEU
1	B	182	LEU
1	B	336	LEU
1	B	352	ASP
1	B	547	LEU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (10) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	110	ASN
1	A	117	ASN
1	A	273	ASN
1	A	330	GLN
1	A	374	HIS
1	B	117	ASN
1	B	142	ASN
1	B	251	GLN
1	B	273	ASN
1	B	436	GLN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

2 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
2	N35	B	579	-	38,38,38	1.87	3 (7%)	49,58,58	2.04	12 (24%)
2	N35	A	579	-	38,38,38	1.91	4 (10%)	49,58,58	1.80	10 (20%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	N35	B	579	-	-	5/12/40/40	0/4/4/4
2	N35	A	579	-	-	3/12/40/40	0/4/4/4

All (7) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	A	579	N35	S34-N24	7.49	1.70	1.62
2	B	579	N35	S35-N27	7.03	1.72	1.63
2	A	579	N35	S35-N27	6.19	1.71	1.63
2	B	579	N35	S34-N24	6.18	1.68	1.62
2	A	579	N35	C8-C10	4.15	1.51	1.42
2	B	579	N35	C8-C10	4.06	1.50	1.42
2	A	579	N35	C9-N24	2.09	1.42	1.37

All (22) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	B	579	N35	C7-C8-C10	-8.24	117.04	121.99
2	A	579	N35	C7-C8-C10	-7.37	117.57	121.99
2	B	579	N35	C18-S35-N27	-4.35	101.67	106.56
2	B	579	N35	C8-C7-C9	3.69	123.05	118.55
2	B	579	N35	O30-S34-O29	3.61	122.11	118.44
2	B	579	N35	O30-S34-C6	-3.51	104.92	109.30
2	A	579	N35	C18-S35-N27	-3.47	102.66	106.56
2	A	579	N35	C8-C7-C9	3.16	122.41	118.55
2	A	579	N35	O30-S34-O29	3.04	121.53	118.44
2	B	579	N35	O28-C11-C7	-2.75	119.29	125.16
2	A	579	N35	O29-S34-C6	-2.72	105.90	109.30
2	B	579	N35	C20-N26-N23	2.69	118.44	113.90
2	B	579	N35	C11-N26-N23	-2.69	123.47	126.24
2	B	579	N35	O32-S35-N27	2.68	112.46	107.10
2	A	579	N35	O28-C11-C7	-2.68	119.44	125.16
2	A	579	N35	C20-N26-N23	2.60	118.30	113.90
2	B	579	N35	C11-C7-C9	-2.43	116.59	120.77
2	B	579	N35	O29-S34-N24	-2.39	106.31	108.26
2	A	579	N35	C11-N26-N23	-2.27	123.90	126.24
2	A	579	N35	O31-S35-C18	-2.26	104.69	108.26
2	B	579	N35	O33-C8-C10	2.22	120.20	116.08
2	A	579	N35	C4-N22-C9	2.19	121.32	117.58

There are no chirality outliers.

All (8) torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
2	B	579	N35	N23-C10-N25-C15
2	B	579	N35	C21-C19-C20-N26
2	A	579	N35	C5-N27-S35-O32
2	A	579	N35	C5-N27-S35-O31
2	B	579	N35	C5-N27-S35-O32
2	B	579	N35	C5-N27-S35-O31
2	A	579	N35	C5-N27-S35-C18
2	B	579	N35	C5-N27-S35-C18


There are no ring outliers.

2 monomers are involved in 2 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	B	579	N35	1	0
2	A	579	N35	1	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.





5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	557/578 (96%)	0.61	36 (6%) 25 26	28, 39, 57, 85	0
1	B	557/578 (96%)	0.67	32 (5%) 29 31	27, 40, 56, 82	0
All	All	1114/1156 (96%)	0.64	68 (6%) 27 29	27, 40, 57, 85	0

All (68) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	545	LEU	4.8
1	B	534	LEU	4.8
1	B	532	THR	4.5
1	A	542	ALA	4.4
1	B	541	ALA	4.4
1	A	537	THR	4.2
1	A	541	ALA	4.0
1	A	311	CYS	4.0
1	A	539	ILE	4.0
1	B	545	LEU	3.9
1	B	542	ALA	3.9
1	B	540	PRO	3.8
1	B	538	PRO	3.7
1	B	536	LEU	3.7
1	A	534	LEU	3.6
1	B	535	LYS	3.6
1	B	539	ILE	3.4
1	B	543	SER	3.4
1	A	381	VAL	3.4
1	B	311	CYS	3.3
1	A	536	LEU	3.3
1	B	537	THR	3.3
1	A	535	LYS	3.0
1	A	111	LEU	3.0

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	B	530	VAL	2.9
1	A	549	GLY	2.9
1	A	538	PRO	2.9
1	A	533	LYS	2.9
1	B	14	CYS	2.9
1	B	160	ILE	2.8
1	A	95	HIS	2.8
1	A	43	ARG	2.7
1	A	513	SER	2.7
1	A	497	LEU	2.6
1	A	16	ALA	2.6
1	B	497	LEU	2.6
1	A	532	THR	2.6
1	A	14	CYS	2.6
1	A	540	PRO	2.5
1	A	550	TRP	2.5
1	A	51	LYS	2.5
1	B	550	TRP	2.5
1	B	562	HIS	2.5
1	A	547	LEU	2.5
1	B	469	LEU	2.5
1	A	377	SER	2.4
1	B	32	ARG	2.4
1	A	499	VAL	2.4
1	A	543	SER	2.4
1	B	148	GLN	2.3
1	B	533	LYS	2.3
1	B	100	LYS	2.3
1	B	547	LEU	2.2
1	B	95	HIS	2.2
1	A	469	LEU	2.2
1	A	402	HIS	2.2
1	A	376	ALA	2.2
1	A	32	ARG	2.2
1	B	119	ILE	2.1
1	A	515	GLY	2.1
1	A	502	HIS	2.1
1	A	15	ALA	2.1
1	B	382	TYR	2.1
1	B	111	LEU	2.1
1	B	515	GLY	2.0
1	A	530	VAL	2.0

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	B	97	ALA	2.0
1	B	519	ALA	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)




In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q < 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
2	N35	B	579	35/35	0.96	0.09	31,34,40,41	0
2	N35	A	579	35/35	0.97	0.08	30,34,43,44	0




The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.

Electron density around N35 B 579:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)

**Electron density around N35 A 579:**

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



6.5 Other polymers [i](#)

There are no such residues in this entry.