



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 10, 2026 – 04:59 AM UTC

PDB ID : 3E59 / pdb_00003e59
Title : Crystal structure of the PvcA (PA2254) protein from *Pseudomonas aeruginosa*
Authors : Gulick, A.M.; Drake, E.J.
Deposited on : 2008-08-13
Resolution : 2.10 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.10 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	6658 (2.10-2.10)
Clashscore	190562	7164 (2.10-2.10)
Ramachandran outliers	187476	7099 (2.10-2.10)
Sidechain outliers	187428	7100 (2.10-2.10)
RSRZ outliers	180081	6662 (2.10-2.10)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	330	4% 86% 8% 6%
1	B	330	4% 78% 13% 8%
1	C	330	14% 81% 11% 8%
1	D	330	2% 85% 9% 5%

2 Entry composition

There are 4 unique types of molecules in this entry. The entry contains 10092 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Pyoverdine biosynthesis protein PvcA.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	311	Total	C	N	O	S	0	1	0
			2443	1557	435	444	7			
1	B	305	Total	C	N	O	S	0	2	0
			2378	1524	423	424	7			
1	C	303	Total	C	N	O	S	0	1	0
			2348	1503	417	421	7			
1	D	313	Total	C	N	O	S	0	2	0
			2456	1567	429	453	7			

There are 8 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	-1	GLY	-	expression tag	UNP Q9I1L5
A	0	HIS	-	expression tag	UNP Q9I1L5
B	-1	GLY	-	expression tag	UNP Q9I1L5
B	0	HIS	-	expression tag	UNP Q9I1L5
C	-1	GLY	-	expression tag	UNP Q9I1L5
C	0	HIS	-	expression tag	UNP Q9I1L5
D	-1	GLY	-	expression tag	UNP Q9I1L5
D	0	HIS	-	expression tag	UNP Q9I1L5

- Molecule 2 is PHOSPHATE ION (CCD ID: PO4) (formula: O₄P).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
2	A	1	Total	O	P	0	0
			5	4	1		
2	A	1	Total	O	P	0	0
			5	4	1		
2	B	1	Total	O	P	0	0
			5	4	1		
2	C	1	Total	O	P	0	0
			5	4	1		
2	D	1	Total	O	P	0	0
			5	4	1		
2	D	1	Total	O	P	0	0
			5	4	1		

- Molecule 3 is TRIETHYLENE GLYCOL (CCD ID: PGE) (formula: C₆H₁₄O₄).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	B	1	Total	C	O	0	0
			10	6	4		
3	B	1	Total	C	O	0	0
			10	6	4		


- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	130	Total	O	0	0
			130	130		
4	B	93	Total	O	0	0
			93	93		
4	C	77	Total	O	0	0
			77	77		
4	D	117	Total	O	0	0
			117	117		


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


• Molecule 1: Pyoverdine biosynthesis protein PvcA




• Molecule 1: Pyoverdine biosynthesis protein PvcA



• Molecule 1: Pyoverdine biosynthesis protein PvcA



● Molecule 1: Pyoverdine biosynthesis protein PvcA



4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	56.23Å 132.53Å 96.90Å 90.00° 97.93° 90.00°	Depositor
Resolution (Å)	30.00 – 2.10 30.00 – 2.10	Depositor EDS
% Data completeness (in resolution range)	96.2 (30.00-2.10) 86.8 (30.00-2.10)	Depositor EDS
R_{merge}	0.05	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.80 (at 2.10Å)	Xtriage
Refinement program	REFMAC 5.2.0005	Depositor
R, R_{free}	0.197 , 0.246 0.199 , 0.245	Depositor DCC
R_{free} test set	3549 reflections (4.79%)	wwPDB-VP
Wilson B-factor (Å ²)	39.2	Xtriage
Anisotropy	0.577	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.33 , 41.9	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.94	EDS
Total number of atoms	10092	wwPDB-VP
Average B, all atoms (Å ²)	42.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The analyses of the Patterson function reveals a significant off-origin peak that is 46.50 % of the origin peak, indicating pseudo-translational symmetry. The chance of finding a peak of this or larger height randomly in a structure without pseudo-translational symmetry is equal to 1.1352e-04. The detected translational NCS is most likely also responsible for the elevated intensity ratio.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: PGE, PO4

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.63	0/2499	0.83	0/3395
1	B	0.63	0/2435	0.86	1/3310 (0.0%)
1	C	0.64	3/2401 (0.1%)	0.89	2/3265 (0.1%)
1	D	0.64	0/2515	0.86	1/3419 (0.0%)
All	All	0.63	3/9850 (0.0%)	0.86	4/13389 (0.0%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	B	0	1

All (3) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	C	295	ARG	C-O	-7.10	1.16	1.24
1	C	215	GLY	C-N	6.17	1.42	1.33
1	C	215	GLY	C-O	5.20	1.28	1.23

All (4) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	C	309	ASN	N-CA-C	9.12	130.22	110.80
1	B	318	PRO	N-CA-CB	6.69	110.36	103.00
1	C	318	PRO	N-CA-CB	6.54	110.19	103.00
1	D	318	PRO	N-CA-CB	6.31	109.94	103.00

There are no chirality outliers.

All (1) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	B	119	ASP	Peptide

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2443	0	2388	15	0
1	B	2378	0	2334	28	0
1	C	2348	0	2281	17	0
1	D	2456	0	2390	21	0
2	A	10	0	0	0	0
2	B	5	0	0	0	0
2	C	5	0	0	0	0
2	D	10	0	0	0	0
3	B	20	0	28	0	0
4	A	130	0	0	0	0
4	B	93	0	0	3	0
4	C	77	0	0	1	0
4	D	117	0	0	0	0
All	All	10092	0	9421	78	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 4.

All (78) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:D:91:LEU:HD11	1:D:268:MET:HE3	1.38	1.05
1:D:91:LEU:CD1	1:D:268:MET:HE3	1.94	0.97
1:B:118:GLY:O	4:B:1058:HOH:O	1.82	0.95
1:A:200:ILE:HG13	1:A:232:ILE:HD11	1.49	0.94
1:D:91:LEU:HD11	1:D:268:MET:CE	2.17	0.73
1:C:37:ILE:O	1:C:41:GLU:HG2	1.89	0.72
1:C:49:ALA:O	1:C:53:GLU:HG2	1.94	0.68
1:D:65:PHE:CE1	1:D:268:MET:HE1	2.31	0.66

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:D:65:PHE:CZ	1:D:268:MET:HE1	2.32	0.65
1:A:200:ILE:HG13	1:A:232:ILE:CD1	2.24	0.65
1:B:92:ASN:HD21	1:B:146:HIS:H	1.45	0.63
1:B:33:GLU:O	1:B:37:ILE:HD12	1.98	0.63
1:D:91:LEU:CG	1:D:268:MET:HE3	2.29	0.61
1:B:92:ASN:ND2	1:B:146:HIS:H	1.98	0.61
1:D:91:LEU:CD1	1:D:268:MET:CE	2.74	0.60
1:B:282:VAL:HG12	4:B:1029:HOH:O	2.01	0.60
1:D:79:ARG:NH2	1:D:311:GLN:OE1	2.36	0.59
1:B:79:ARG:HH21	1:B:312:PRO:HD2	1.69	0.56
1:C:284:VAL:HG21	1:C:293:MET:HE2	1.86	0.56
1:C:276:LEU:HD21	1:C:281:GLY:HA3	1.87	0.56
1:A:97:ARG:HH22	1:B:19:LEU:HD13	1.72	0.55
1:A:68:LYS:HG3	1:A:75:VAL:HG11	1.88	0.54
1:A:200:ILE:CG1	1:A:232:ILE:HD11	2.32	0.54
1:C:268:MET:HA	1:C:268:MET:HE2	1.91	0.53
1:A:188:LEU:HD22	1:A:194:LEU:HD21	1.91	0.53
1:B:173:ILE:HD11	1:B:234:ARG:NE	2.24	0.52
1:C:92:ASN:HD21	1:C:146:HIS:H	1.57	0.52
1:A:308:ILE:HD13	1:B:141:GLU:HA	1.92	0.52
1:C:11:ALA:O	1:C:15:LYS:HG3	2.11	0.51
1:D:115:ARG:HG3	1:D:153:GLU:OE1	2.10	0.51
1:B:115:ARG:HD2	1:B:153:GLU:OE1	2.11	0.51
1:C:92:ASN:ND2	1:C:146:HIS:H	2.09	0.51
1:B:230:GLY:O	1:B:234:ARG:HG2	2.10	0.50
1:C:293:MET:HE3	1:C:298:VAL:HG22	1.93	0.50
1:A:161:GLN:NE2	1:A:167:GLN:OE1	2.46	0.49
1:A:97:ARG:NH2	1:B:19:LEU:HD13	2.28	0.48
1:D:230:GLY:O	1:D:234:ARG:HG2	2.14	0.48
1:A:228:ALA:O	1:A:232:ILE:HG12	2.14	0.48
1:C:269:MET:HG3	1:C:270:PRO:HD2	1.96	0.47
1:B:12:ARG:NH2	4:B:1078:HOH:O	2.47	0.47
1:B:173:ILE:HD11	1:B:234:ARG:CZ	2.45	0.47
1:B:106:ALA:C	1:B:107:LYS:HD2	2.40	0.46
1:A:60:PHE:HB2	1:A:108:ILE:HG12	1.96	0.46
1:D:60:PHE:CD2	1:D:250:ILE:HB	2.51	0.46
1:D:68:LYS:HG2	1:D:77:ASP:O	2.16	0.46
1:B:63:PRO:HA	1:B:111:CYS:O	2.15	0.46
1:B:136:ARG:HH21	1:B:151:ASN:HD22	1.64	0.45
1:B:60:PHE:CD2	1:B:250:ILE:HB	2.51	0.45
1:C:230:GLY:O	1:C:234:ARG:HG2	2.17	0.45

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:D:173:ILE:HD11	1:D:234:ARG:CZ	2.46	0.45
1:D:238:TRP:O	1:D:242:LEU:HG	2.16	0.45
1:B:107:LYS:NZ	1:B:146:HIS:HD2	2.14	0.44
1:B:284:VAL:HG21	1:B:293:MET:CG	2.47	0.44
1:A:181:GLU:H	1:A:181:GLU:CD	2.26	0.44
1:D:72:PRO:O	1:D:295:ARG:NH1	2.51	0.44
1:B:76:LEU:HD21	1:B:305:LEU:HD13	2.01	0.43
1:B:68:LYS:HG2	1:B:77:ASP:O	2.19	0.43
1:B:132[A]:GLN:HE21	1:B:151:ASN:HD21	1.66	0.43
1:C:119:ASP:OD1	1:C:119:ASP:N	2.51	0.43
1:D:295:ARG:O	1:D:298:VAL:HG12	2.19	0.43
1:D:62:LEU:HD11	1:D:254:ILE:HG23	2.00	0.43
1:A:7:ASP:O	1:A:11:ALA:HB3	2.18	0.42
1:A:112:SER:HB3	1:A:151:ASN:HD22	1.84	0.42
1:C:71:ASN:HA	1:C:72:PRO:HD3	1.93	0.42
1:C:294:LYS:O	1:C:298:VAL:HG23	2.20	0.42
1:B:200:ILE:CD1	1:B:232:ILE:HG12	2.50	0.41
1:C:9:LEU:HB3	1:C:10:PRO:HD3	2.03	0.41
1:C:60:PHE:CD2	1:C:250:ILE:HB	2.56	0.41
1:C:282:VAL:HG12	4:C:362:HOH:O	2.18	0.41
1:B:132[A]:GLN:NE2	1:B:151:ASN:HD21	2.19	0.41
1:D:202:ARG:O	1:D:206:GLU:HG3	2.20	0.41
1:D:91:LEU:HG	1:D:268:MET:HE3	2.01	0.41
1:B:200:ILE:HD13	1:B:232:ILE:CG1	2.51	0.40
1:A:9:LEU:HB2	1:A:10:PRO:HD3	2.03	0.40
1:D:161:GLN:HG3	1:D:171:LEU:HD11	2.02	0.40
1:B:162:ARG:HE	1:B:162:ARG:HB3	1.62	0.40
1:D:70:PRO:O	1:D:72:PRO:HD3	2.22	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	310/330 (94%)	301 (97%)	8 (3%)	1 (0%)	36	36
1	B	303/330 (92%)	297 (98%)	5 (2%)	1 (0%)	36	36
1	C	300/330 (91%)	293 (98%)	5 (2%)	2 (1%)	18	15
1	D	313/330 (95%)	307 (98%)	6 (2%)	0	100	100
All	All	1226/1320 (93%)	1198 (98%)	24 (2%)	4 (0%)	36	36

All (4) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	8	THR
1	B	67	ALA
1	C	67	ALA
1	C	309	ASN

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	248/272 (91%)	240 (97%)	8 (3%)	34	38
1	B	239/272 (88%)	229 (96%)	10 (4%)	26	28
1	C	233/272 (86%)	228 (98%)	5 (2%)	47	54
1	D	250/272 (92%)	245 (98%)	5 (2%)	48	56
All	All	970/1088 (89%)	942 (97%)	28 (3%)	37	42

All (28) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	30	SER
1	A	39	ARG
1	A	68	LYS
1	A	116	VAL
1	A	182	SER
1	A	188	LEU
1	A	198	ARG

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	A	307	GLN
1	B	15	LYS
1	B	37	ILE
1	B	115	ARG
1	B	137	LEU
1	B	206	GLU
1	B	209	LEU
1	B	212	ASP
1	B	223	ASP
1	B	276	LEU
1	B	307	GLN
1	C	84	GLU
1	C	110	VAL
1	C	185	GLU
1	C	202	ARG
1	C	206	GLU
1	D	12	ARG
1	D	115	ARG
1	D	140	GLU
1	D	251	ARG
1	D	263	LYS

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (14) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	96	GLN
1	A	151	ASN
1	A	285	ASN
1	B	92	ASN
1	B	146	HIS
1	B	151	ASN
1	B	307	GLN
1	C	132	GLN
1	C	151	ASN
1	C	307	GLN
1	D	96	GLN
1	D	99	GLN
1	D	164	ASN
1	D	221	GLN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no oligosaccharides in this entry.

5.6 Ligand geometry ⓘ

8 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
2	PO4	A	329	-	4,4,4	0.76	0	6,6,6	0.72	0
3	PGE	B	991	-	9,9,9	0.49	0	8,8,8	0.26	0
2	PO4	C	329	-	4,4,4	0.96	0	6,6,6	0.76	0
2	PO4	B	329	-	4,4,4	1.31	1 (25%)	6,6,6	0.47	0
2	PO4	D	329	-	4,4,4	0.79	0	6,6,6	1.02	0
2	PO4	A	330	-	4,4,4	0.88	0	6,6,6	0.47	0
2	PO4	D	330	-	4,4,4	0.85	0	6,6,6	0.54	0
3	PGE	B	992	-	9,9,9	0.52	0	8,8,8	0.17	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	PGE	B	992	-	-	4/7/7/7	-
3	PGE	B	991	-	-	4/7/7/7	-

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	B	329	PO4	P-O4	-2.04	1.48	1.54

There are no bond angle outliers.

There are no chirality outliers.

All (8) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
3	B	992	PGE	O3-C5-C6-O4
3	B	991	PGE	O1-C1-C2-O2
3	B	992	PGE	O2-C3-C4-O3
3	B	992	PGE	C1-C2-O2-C3
3	B	991	PGE	O2-C3-C4-O3
3	B	992	PGE	C3-C4-O3-C5
3	B	991	PGE	O3-C5-C6-O4
3	B	991	PGE	C4-C3-O2-C2

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	311/330 (94%)	0.53	13 (4%) 40 42	19, 40, 53, 71	1 (0%)
1	B	305/330 (92%)	0.53	14 (4%) 37 39	17, 42, 57, 69	2 (0%)
1	C	303/330 (91%)	0.94	45 (14%) 5 5	23, 44, 63, 74	1 (0%)
1	D	313/330 (94%)	0.35	8 (2%) 57 60	19, 39, 53, 70	2 (0%)
All	All	1232/1320 (93%)	0.59	80 (6%) 25 26	17, 41, 59, 74	6 (0%)

All (80) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	118	GLY	6.8
1	C	219	ALA	4.9
1	C	300	GLU	4.4
1	A	7	ASP	4.4
1	C	288	ASP	4.3
1	C	289	ARG	4.2
1	C	286	THR	4.2
1	C	291	VAL	4.2
1	A	8	THR	4.1
1	C	222	ARG	4.1
1	C	218	THR	3.8
1	C	118	GLY	3.7
1	C	275	TRP	3.5
1	C	215	GLY	3.5
1	C	307	GLN	3.3
1	B	212	ASP	3.3
1	C	119	ASP	3.2
1	C	271	THR	3.2
1	C	281	GLY	3.1
1	A	119	ASP	3.1
1	A	118	GLY	3.1

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	286	THR	3.0
1	B	8	THR	3.0
1	C	140	GLU	2.9
1	C	221	GLN	2.8
1	D	167	GLN	2.8
1	B	275	TRP	2.8
1	C	33	GLU	2.8
1	C	298	VAL	2.7
1	C	309	ASN	2.7
1	C	53	GLU	2.7
1	C	301	LEU	2.7
1	C	274	ASP	2.6
1	C	220	LEU	2.6
1	C	117	PHE	2.5
1	D	30	SER	2.5
1	B	140	GLU	2.5
1	D	318	PRO	2.5
1	C	284	VAL	2.4
1	B	276	LEU	2.4
1	A	93	HIS	2.4
1	C	290	PHE	2.4
1	B	35	ASP	2.4
1	B	219	ALA	2.4
1	D	226	GLU	2.3
1	C	308	ILE	2.3
1	B	24	TYR	2.3
1	A	280	HIS	2.3
1	B	43	VAL	2.3
1	C	292	LEU	2.3
1	B	300	GLU	2.3
1	A	291	VAL	2.3
1	C	279	TRP	2.3
1	C	318	PRO	2.2
1	C	302	GLY	2.2
1	A	177	ALA	2.2
1	A	31	ALA	2.2
1	C	199	ALA	2.2
1	C	317	LEU	2.2
1	C	213	TYR	2.2
1	B	279	TRP	2.2
1	C	265	GLY	2.1
1	C	224	ALA	2.1

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	D	119	ASP	2.1
1	C	203	PHE	2.1
1	C	198	ARG	2.1
1	C	15	LYS	2.1
1	A	29	GLN	2.1
1	A	211	PRO	2.1
1	B	296	SER	2.1
1	C	299	LEU	2.1
1	D	276	LEU	2.1
1	B	310	GLY	2.1
1	C	306	VAL	2.1
1	C	260	ASP	2.0
1	D	163	ASP	2.0
1	C	54	ALA	2.0
1	C	305	LEU	2.0
1	D	29	GLN	2.0
1	A	55	GLY	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
2	PO4	D	330	5/5	0.69	0.14	84,84,84,85	0
2	PO4	A	330	5/5	0.78	0.12	81,82,82,82	0
3	PGE	B	991	10/10	0.80	0.15	59,61,63,64	0
3	PGE	B	992	10/10	0.80	0.13	51,57,60,60	0
2	PO4	C	329	5/5	0.84	0.13	25,34,38,39	5
2	PO4	B	329	5/5	0.89	0.20	30,34,39,40	5

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
2	PO4	D	329	5/5	0.94	0.18	39,48,49,49	0
2	PO4	A	329	5/5	0.95	0.14	31,37,38,39	0

6.5 Other polymers [i](#)

There are no such residues in this entry.