



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 20, 2026 – 06:27 AM UTC

PDB ID : 4E5S / pdb_00004e5s
Title : Crystal structure of MccFlike protein (BA_5613) from Bacillus anthracis str. Ames
Authors : Nocek, B.; Anderson, W.F.; Joachimiak, A.; Center for Structural Genomics of Infectious Diseases (CSGID)
Deposited on : 2012-03-14
Resolution : 1.95 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0
Mogul : 2022.3.0, CSD as543be (2022)
Xtriage (Phenix) : 2.0
EDS : 3.0
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4 : 9.0.010 (Gargrove)
Density-Fitness : 1.0.12
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.95 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	3494 (1.96-1.96)
Clashscore	190562	3612 (1.96-1.96)
Ramachandran outliers	187476	3587 (1.96-1.96)
Sidechain outliers	187428	3587 (1.96-1.96)
RSRZ outliers	180081	3495 (1.96-1.96)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	331	2% 84% 15% ..
1	B	331	86% 13% .
1	C	331	90% 10%
1	D	331	2% 85% 13% ..

2 Entry composition

There are 3 unique types of molecules in this entry. The entry contains 11229 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called MccFlike protein (BA_5613).

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
1	A	329	Total	C	N	O	S	Se	0	0	0
			2567	1631	427	498	4	7			
1	B	329	Total	C	N	O	S	Se	0	3	0
			2586	1644	429	502	4	7			
1	C	330	Total	C	N	O	S	Se	0	4	0
			2590	1646	432	501	4	7			
1	D	329	Total	C	N	O	S	Se	0	1	0
			2549	1618	424	496	4	7			

There are 12 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	-2	SER	-	expression tag	UNP Q81JT5
A	-1	ASN	-	expression tag	UNP Q81JT5
A	0	ALA	-	expression tag	UNP Q81JT5
B	-2	SER	-	expression tag	UNP Q81JT5
B	-1	ASN	-	expression tag	UNP Q81JT5
B	0	ALA	-	expression tag	UNP Q81JT5
C	-2	SER	-	expression tag	UNP Q81JT5
C	-1	ASN	-	expression tag	UNP Q81JT5
C	0	ALA	-	expression tag	UNP Q81JT5
D	-2	SER	-	expression tag	UNP Q81JT5
D	-1	ASN	-	expression tag	UNP Q81JT5
D	0	ALA	-	expression tag	UNP Q81JT5

- Molecule 2 is GLYCEROL (CCD ID: GOL) (formula: C₃H₈O₃).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
2	A	1	Total	C	O	0	0
			6	3	3		


- Molecule 3 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	262	Total	O	0	0
			262	262		
3	B	321	Total	O	0	0
			321	321		
3	C	249	Total	O	0	0
			249	249		
3	D	99	Total	O	0	0
			99	99		


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: MccFlike protein (BA_5613)




- Molecule 1: MccFlike protein (BA_5613)




- Molecule 1: MccFlike protein (BA_5613)



- Molecule 1: MccFlike protein (BA_5613)





4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	74.58Å 85.80Å 125.15Å 90.00° 92.00° 90.00°	Depositor
Resolution (Å)	32.52 – 1.95 32.52 – 1.95	Depositor EDS
% Data completeness (in resolution range)	68.5 (32.52-1.95) 86.0 (32.52-1.95)	Depositor EDS
R_{merge}	0.07	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	4.12 (at 1.95Å)	Xtriage
Refinement program	PHENIX (phenix.refine: 1.7.3_928)/REFMAC	Depositor
R, R_{free}	0.153 , 0.202 0.155 , 0.202	Depositor DCC
R_{free} test set	5739 reflections (5.01%)	wwPDB-VP
Wilson B-factor (Å ²)	22.6	Xtriage
Anisotropy	0.140	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.32 , 51.3	EDS
L-test for twinning ²	$\langle L \rangle = 0.48$, $\langle L^2 \rangle = 0.31$	Xtriage
Estimated twinning fraction	0.027 for h,-k,-l	Xtriage
F_o, F_c correlation	0.96	EDS
Total number of atoms	11229	wwPDB-VP
Average B, all atoms (Å ²)	33.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.53% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section:
GOL

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	1.14	7/2613 (0.3%)	1.17	4/3527 (0.1%)
1	B	1.20	12/2641 (0.5%)	1.22	9/3565 (0.3%)
1	C	1.07	1/2645 (0.0%)	1.09	2/3569 (0.1%)
1	D	0.95	1/2599 (0.0%)	1.07	5/3516 (0.1%)
All	All	1.10	21/10498 (0.2%)	1.14	20/14177 (0.1%)

All (21) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	B	37	THR	C-O	-8.54	1.14	1.24
1	A	201	ILE	C-O	-7.84	1.15	1.23
1	B	228	GLU	C-O	-7.36	1.14	1.23
1	B	224	ILE	C-O	-7.01	1.16	1.24
1	A	151	LEU	C-O	-7.00	1.16	1.24
1	B	36	LEU	C-O	-6.74	1.16	1.24
1	A	200	ILE	C-O	-6.27	1.15	1.24
1	B	42	HIS	CG-ND1	-6.27	1.31	1.38
1	A	206	SER	C-O	-6.24	1.16	1.24
1	B	225	LEU	C-O	-6.20	1.16	1.24
1	B	226	PHE	C-O	-5.48	1.17	1.24
1	A	204	ASN	C-O	-5.44	1.17	1.24
1	B	40	GLY	C-O	-5.41	1.17	1.24
1	B	38	GLU	C-O	-5.38	1.17	1.24
1	B	41	PHE	C-O	-5.34	1.17	1.23
1	B	39	LEU	C-O	-5.25	1.17	1.24
1	A	199	ASP	C-O	-5.22	1.17	1.23
1	D	244	TYR	C-O	5.15	1.30	1.24
1	A	205	MSE	C-O	-5.13	1.18	1.24
1	C	222	ASP	CG-OD2	5.11	1.35	1.25
1	B	42	HIS	CD2-NE2	-5.08	1.32	1.37

All (20) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	D	300	THR	N-CA-C	-7.07	101.41	110.53
1	B	217	MSE	CA-C-N	-6.48	113.61	120.03
1	B	217	MSE	C-N-CA	-6.48	113.61	120.03
1	B	44	THR	N-CA-CB	-6.19	100.00	111.52
1	B	309	GLY	N-CA-C	-5.88	106.98	115.27
1	A	298	GLY	N-CA-C	-5.84	105.13	111.21
1	A	222	ASP	N-CA-C	-5.69	106.29	113.23
1	B	111	SER	CB-CA-C	-5.61	110.12	116.63
1	B	165	SER	N-CA-C	-5.60	103.52	110.41
1	D	299	HIS	N-CA-C	5.38	119.94	113.17
1	A	2	LEU	CA-C-N	5.34	125.28	119.78
1	A	2	LEU	C-N-CA	5.34	125.28	119.78
1	C	111	SER	CB-CA-C	-5.32	110.46	116.63
1	B	2	LEU	CA-C-N	-5.31	114.77	120.03
1	B	2	LEU	C-N-CA	-5.31	114.77	120.03
1	B	83	LEU	CB-CA-C	-5.30	100.78	111.17
1	D	39	LEU	N-CA-C	-5.25	106.93	113.38
1	D	241	PHE	N-CA-C	-5.18	105.64	111.28
1	C	175	ILE	CB-CA-C	-5.17	105.42	111.94
1	D	71	PHE	N-CA-C	5.09	116.83	111.28

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2567	0	2538	34	0
1	B	2586	0	2561	15	0
1	C	2590	0	2558	16	0
1	D	2549	0	2484	22	0
2	A	6	0	8	1	0
3	A	262	0	0	1	0
3	B	321	0	0	1	0
3	C	249	0	0	2	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
3	D	99	0	0	0	0
All	All	11229	0	10149	83	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 4.

All (83) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:11:GLU:OE2	1:B:44:THR:HG21	1.84	0.78
1:B:13:ARG:HE	1:B:44:THR:CG2	1.98	0.76
1:A:23:ILE:HD11	1:A:52:ILE:HD11	1.71	0.72
1:D:200:ILE:HD11	1:D:312:ALA:HB2	1.76	0.68
1:C:54:ARG:NE	1:D:278:GLU:OE2	2.29	0.66
1:B:13:ARG:HE	1:B:44:THR:HG23	1.63	0.64
1:D:39:LEU:HD22	1:D:151:LEU:HD11	1.80	0.63
1:D:319:GLU:O	1:D:320:LYS:CB	2.45	0.63
1:A:23:ILE:HD13	1:A:52:ILE:HD12	1.81	0.62
1:C:200:ILE:HD11	1:C:312:ALA:HB2	1.82	0.61
1:C:34:LYS:O	1:C:38:GLU:HG3	2.03	0.59
1:D:70:ALA:HB1	1:D:79:ILE:HG12	1.84	0.59
1:D:206:SER:N	1:D:229:GLU:OE1	2.33	0.59
1:D:311:ARG:HB3	1:D:327:THR:HG23	1.86	0.57
1:A:22:SER:O	2:A:501:GOL:H2	2.04	0.57
1:A:23:ILE:CD1	1:A:52:ILE:CD1	2.82	0.57
1:D:165:SER:OG	1:D:167:THR:O	2.23	0.57
1:A:23:ILE:CD1	1:A:52:ILE:HD11	2.34	0.57
1:B:28:ASN:HB3	3:B:604:HOH:O	2.04	0.56
1:C:221[B]:LYS:NZ	3:C:630:HOH:O	2.39	0.55
1:D:307:PRO:HG3	1:D:328:HIS:HB3	1.89	0.55
1:A:245:LEU:O	1:A:249:MSE:HG3	2.08	0.54
1:A:279:MSE:HE2	1:A:279:MSE:C	2.32	0.54
1:A:197:THR:OG1	1:A:313:THR:HG22	2.08	0.54
1:A:72:ARG:HD3	3:A:790:HOH:O	2.09	0.53
1:B:246:HIS:HA	1:B:249:MSE:HE2	1.90	0.52
1:A:279:MSE:HE3	1:A:283:LYS:HE2	1.91	0.52
1:C:86:TYR:CD2	1:D:239:LYS:HD3	2.45	0.51
1:A:23:ILE:HD13	1:A:52:ILE:CD1	2.40	0.51
1:A:313:THR:HG23	1:A:326:LEU:HD11	1.93	0.50
1:A:21:LEU:HA	1:A:82:THR:HG21	1.94	0.50
1:D:66:ASP:O	1:D:69:GLU:HG3	2.11	0.50

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:5:LYS:HE3	1:A:156:SER:O	2.11	0.49
1:A:26:THR:O	1:A:30:ARG:HG3	2.12	0.49
1:A:176:ASP:OD1	1:A:178:GLU:HG2	2.12	0.49
1:A:8:LYS:HD3	1:A:40:GLY:HA3	1.95	0.48
1:D:238:LEU:CD2	1:D:275:ASP:HB3	2.45	0.46
1:B:230:ASP:OD1	1:B:230:ASP:C	2.59	0.46
1:C:273:ILE:O	1:C:277:GLN:HG3	2.16	0.46
1:A:31:LEU:HD21	1:A:140:GLU:HG2	1.98	0.46
1:C:39:LEU:HD22	1:C:151:LEU:HD11	1.98	0.45
1:B:21:LEU:HA	1:B:82:THR:HG21	1.99	0.45
1:D:259:GLY:HA2	1:D:289:ILE:HD12	1.98	0.45
1:A:279:MSE:HE3	1:A:283:LYS:CE	2.47	0.45
1:A:279:MSE:HE1	1:A:280:ILE:HD13	1.97	0.45
1:B:193:GLU:HG2	1:B:317:SER:HA	1.99	0.45
1:D:144:GLU:HG3	1:D:145:TYR:N	2.31	0.45
1:D:283:LYS:HA	1:D:284:PRO:HD2	1.84	0.44
1:A:230:ASP:C	1:A:230:ASP:OD1	2.61	0.44
1:A:23:ILE:CD1	1:A:52:ILE:HD12	2.48	0.44
1:A:240:THR:O	1:A:243:ARG:HG2	2.18	0.44
1:B:13:ARG:HA	1:B:44:THR:HG22	2.00	0.43
1:B:238:LEU:CD2	1:B:275:ASP:HB3	2.48	0.43
1:D:227:LEU:O	1:D:262:ILE:HA	2.18	0.43
1:A:278:GLU:OE2	1:B:54:ARG:NE	2.45	0.43
1:C:238:LEU:HD11	1:C:279:MSE:HE2	1.99	0.43
1:A:38:GLU:HA	1:C:8:LYS:HD3	2.00	0.43
1:C:29[B]:ARG:HE	1:C:29[B]:ARG:HB3	1.60	0.43
1:B:106:PHE:CE2	1:B:108:CYS:HB2	2.54	0.42
1:B:195:GLU:HG2	1:B:315:ILE:HG12	2.01	0.42
1:C:148:ASP:O	1:C:152:GLN:HG3	2.19	0.42
1:C:28:ASN:HA	1:C:31:LEU:HD12	2.00	0.42
1:A:222:ASP:HA	1:A:258:LYS:HE2	2.01	0.42
1:A:225:LEU:HD13	1:A:248:LEU:HD21	2.02	0.42
1:B:77:LYS:HA	1:B:77:LYS:HD3	1.82	0.42
1:C:72[A]:ARG:NH2	3:C:441:HOH:O	2.47	0.42
1:A:245:LEU:HD23	1:A:279:MSE:HE1	2.02	0.41
1:C:82:THR:HG22	1:C:83:LEU:HB2	2.02	0.41
1:A:2:LEU:HD22	1:A:328:HIS:C	2.45	0.41
1:A:140:GLU:HG3	1:A:177:GLN:CD	2.46	0.41
1:A:23:ILE:HD12	1:A:23:ILE:HA	1.84	0.41
1:C:283:LYS:HA	1:C:284:PRO:HD2	1.85	0.41
1:D:21:LEU:HA	1:D:82:THR:HG21	2.02	0.41

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:D:228:GLU:HG2	1:D:298:GLY:HA2	2.03	0.41
1:A:15:ILE:HG22	1:A:46:SER:HB3	2.03	0.41
1:A:279:MSE:HE2	1:A:280:ILE:N	2.36	0.41
1:A:238:LEU:CD2	1:A:275:ASP:HB3	2.50	0.40
1:D:8:LYS:O	1:D:40:GLY:O	2.39	0.40
1:B:61:SER:O	1:B:65:GLN:HG3	2.21	0.40
1:C:201:ILE:HD13	1:C:201:ILE:HG21	1.87	0.40
1:D:123:LYS:HA	1:D:123:LYS:HD2	1.88	0.40
1:D:146:THR:HA	1:D:303:ILE:O	2.21	0.40
1:D:73:ASP:HA	1:D:74:PRO:HD3	1.93	0.40

There are no symmetry-related clashes.

5.3 Torsion angles ⓘ

5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	327/331 (99%)	320 (98%)	7 (2%)	0	100	100
1	B	330/331 (100%)	323 (98%)	7 (2%)	0	100	100
1	C	332/331 (100%)	325 (98%)	7 (2%)	0	100	100
1	D	328/331 (99%)	312 (95%)	15 (5%)	1 (0%)	36	28
All	All	1317/1324 (100%)	1280 (97%)	36 (3%)	1 (0%)	48	41

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	D	320	LYS

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	286/286 (100%)	280 (98%)	6 (2%)	47	41
1	B	289/286 (101%)	282 (98%)	7 (2%)	43	36
1	C	287/286 (100%)	282 (98%)	5 (2%)	53	49
1	D	281/286 (98%)	274 (98%)	7 (2%)	42	34
All	All	1143/1144 (100%)	1118 (98%)	25 (2%)	45	40

All (25) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	23	ILE
1	A	161	GLU
1	A	227	LEU
1	A	255	LYS
1	A	267	LYS
1	A	279	MSE
1	B	44	THR
1	B	151	LEU
1	B	209	ASN
1	B	215	SER
1	B	227	LEU
1	B	324	THR
1	B	327	THR
1	C	23	ILE
1	C	35	ARG
1	C	156	SER
1	C	159	PRO
1	C	285	GLU
1	D	35	ARG
1	D	69	GLU
1	D	229	GLU
1	D	278	GLU
1	D	279	MSE
1	D	314	ILE
1	D	327	THR

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (8) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	157	ASN
1	A	212	GLN
1	B	157	ASN
1	C	89	ASN
1	D	28	ASN
1	D	42	HIS
1	D	48	HIS
1	D	253	ASN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

1 ligand is modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
2	GOL	A	501	-	5,5,5	0.20	0	5,5,5	1.02	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the

Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	GOL	A	501	-	-	3/4/4/4	-

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

All (3) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	A	501	GOL	O1-C1-C2-C3
2	A	501	GOL	C1-C2-C3-O3
2	A	501	GOL	O2-C2-C3-O3

There are no ring outliers.

1 monomer is involved in 1 short contact:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	A	501	GOL	1	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	322/331 (97%)	-0.40	3 (0%) 81 85	16, 25, 46, 75	0
1	B	322/331 (97%)	-0.55	0 100 100	14, 23, 40, 53	3 (0%)
1	C	323/331 (97%)	-0.37	0 100 100	14, 28, 45, 63	4 (1%)
1	D	322/331 (97%)	0.45	6 (1%) 66 74	29, 47, 68, 82	1 (0%)
All	All	1289/1324 (97%)	-0.22	9 (0%) 84 88	14, 29, 58, 82	8 (0%)

All (9) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	D	0	ALA	3.5
1	A	222	ASP	3.1
1	D	6	LEU	2.9
1	D	319	GLU	2.5
1	D	313	THR	2.4
1	D	183	ILE	2.3
1	A	0	ALA	2.1
1	A	157	ASN	2.0
1	D	43	VAL	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
2	GOL	A	501	6/6	0.89	0.10	45,52,55,58	0

6.5 Other polymers

There are no such residues in this entry.