



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 10, 2026 – 03:51 AM UTC

PDB ID : 6E57 / pdb_00006e57
Title : Bacteroides ovatus mixed-linkage glucan utilization locus (MLGUL) SGBP-B
in complex with mixed-linkage heptasaccharide
Authors : Koropatkin, N.M.; Schnizlein, M.; Bahr, C.M.E.
Deposited on : 2018-07-19
Resolution : 2.71 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.71 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	4348 (2.74-2.70)
Clashscore	190562	4665 (2.74-2.70)
Ramachandran outliers	187476	4584 (2.74-2.70)
Sidechain outliers	187428	4585 (2.74-2.70)
RSRZ outliers	180081	4348 (2.74-2.70)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	420	
1	B	420	
1	C	420	
1	D	420	
2	E	5	

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
2	F	5	 60%40%
2	G	5	 60%40%

2 Entry composition


There are 6 unique types of molecules in this entry. The entry contains 12333 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called surface glycan binding protein B.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	399	Total	C	N	O	S	0	0	0
			3050	1910	495	637	8			
1	B	385	Total	C	N	O	S	0	0	0
			2941	1842	479	612	8			
1	C	390	Total	C	N	O	S	0	0	0
			2979	1865	486	620	8			
1	D	388	Total	C	N	O	S	0	0	0
			2963	1856	484	615	8			

- Molecule 2 is an oligosaccharide called beta-D-glucopyranose-(1-4)-beta-D-glucopyranose-(1-4)-beta-D-glucopyranose-(1-4)-beta-D-glucopyranose.




Mol	Chain	Residues	Atoms			ZeroOcc	AltConf	Trace
2	E	5	Total	C	O	0	0	0
			56	30	26			
2	F	5	Total	C	O	0	0	0
			56	30	26			
2	G	5	Total	C	O	0	0	0
			56	30	26			

- Molecule 3 is 1,2-ETHANEDIOL (CCD ID: EDO) (formula: C₂H₆O₂).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	A	1	Total	C	O	0	0
			4	2	2		
3	A	1	Total	C	O	0	0
			4	2	2		
3	B	1	Total	C	O	0	0
			4	2	2		
3	C	1	Total	C	O	0	0
			4	2	2		
3	D	1	Total	C	O	0	0
			4	2	2		

- Molecule 4 is PHOSPHATE ION (CCD ID: PO4) (formula: O_4P).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
4	A	1	Total O P 5 4 1	0	0
4	A	1	Total O P 5 4 1	0	0
4	A	1	Total O P 5 4 1	0	0
4	A	1	Total O P 5 4 1	0	0
4	B	1	Total O P 5 4 1	0	0
4	B	1	Total O P 5 4 1	0	0
4	B	1	Total O P 5 4 1	0	0
4	C	1	Total O P 5 4 1	0	0
4	C	1	Total O P 5 4 1	0	0
4	C	1	Total O P 5 4 1	0	0
4	C	1	Total O P 5 4 1	0	0
4	D	1	Total O P 5 4 1	0	0
4	D	1	Total O P 5 4 1	0	0
4	D	1	Total O P 5 4 1	0	0

- Molecule 5 is SODIUM ION (CCD ID: NA) (formula: Na).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
5	A	1	Total Na 1 1	0	0
5	C	1	Total Na 1 1	0	0

- Molecule 6 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
6	A	48	Total O 48 48	0	0
6	B	19	Total O 19 19	0	0

Continued on next page...


Continued from previous page...

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
6	C	45	Total 45	O 45	0	0
6	D	28	Total 28	O 28	0	0


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: surface glycan binding protein B




- Molecule 1: surface glycan binding protein B




- Molecule 1: surface glycan binding protein B



- Molecule 1: surface glycan binding protein B





- Molecule 2: beta-D-glucopyranose-(1-4)-beta-D-glucopyranose-(1-4)-beta-D-glucopyranose-(1-4)-beta-D-glucopyranose-(1-4)-beta-D-glucopyranose

Chain E: 80% 20%



- Molecule 2: beta-D-glucopyranose-(1-4)-beta-D-glucopyranose-(1-4)-beta-D-glucopyranose-(1-4)-beta-D-glucopyranose-(1-4)-beta-D-glucopyranose

Chain F: 60% 40%



- Molecule 2: beta-D-glucopyranose-(1-4)-beta-D-glucopyranose-(1-4)-beta-D-glucopyranose-(1-4)-beta-D-glucopyranose-(1-4)-beta-D-glucopyranose

Chain G: 60% 40%



4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	156.44Å 243.65Å 76.06Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	78.34 – 2.71 78.34 – 2.71	Depositor EDS
% Data completeness (in resolution range)	97.7 (78.34-2.71) 97.7 (78.34-2.71)	Depositor EDS
R_{merge}	0.12	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.36 (at 2.69Å)	Xtriage
Refinement program	REFMAC 5.8.0222	Depositor
R, R_{free}	0.194 , 0.238 (Not available) , 0.238	Depositor DCC
R_{free} test set	3927 reflections (5.03%)	wwPDB-VP
Wilson B-factor (Å ²)	49.9	Xtriage
Anisotropy	0.113	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.34 , 49.0	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.94	EDS
Total number of atoms	12333	wwPDB-VP
Average B, all atoms (Å ²)	58.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 2.76% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: EDO, BGC, NA, PO4

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.56	0/3106	0.81	0/4240
1	B	0.56	0/2994	0.81	0/4087
1	C	0.56	0/3033	0.80	0/4139
1	D	0.55	0/3017	0.82	0/4117
All	All	0.56	0/12150	0.81	0/16583

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	1
1	D	0	1
All	All	0	2

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

All (2) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	143	ARG	Sidechain
1	D	118	LYS	Peptide

5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3050	0	2952	12	0
1	B	2941	0	2847	7	0
1	C	2979	0	2890	7	0
1	D	2963	0	2877	9	0
2	E	56	0	48	0	0
2	F	56	0	48	0	0
2	G	56	0	48	0	0
3	A	8	0	12	0	0
3	B	4	0	6	0	0
3	C	4	0	6	0	0
3	D	4	0	6	0	0
4	A	20	0	0	0	0
4	B	15	0	0	0	0
4	C	20	0	0	0	0
4	D	15	0	0	0	0
5	A	1	0	0	0	0
5	C	1	0	0	0	0
6	A	48	0	0	0	0
6	B	19	0	0	0	0
6	C	45	0	0	0	0
6	D	28	0	0	0	0
All	All	12333	0	11740	34	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 1.

All (34) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:C:33:GLU:HA	1:C:34:ASN:HB3	1.56	0.86
1:C:33:GLU:HA	1:C:34:ASN:CB	2.19	0.71
1:B:70:THR:HG22	1:B:71:PRO:HD3	1.75	0.67
1:D:33:ASN:HA	1:D:34:ASP:CB	2.34	0.56
1:A:143:ARG:HH11	1:A:143:ARG:CG	2.21	0.53
1:B:333:PHE:HB3	1:B:407:CYS:HB2	1.91	0.52

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:223:VAL:HG13	1:A:238:ILE:HG23	1.94	0.50
1:A:333:PHE:HB3	1:A:407:CYS:HB2	1.94	0.49
1:D:33:ASN:HA	1:D:34:ASP:HB2	1.93	0.49
1:C:33:GLU:CA	1:C:34:ASN:CB	2.90	0.48
1:C:334:PHE:HB3	1:C:408:CYS:HB2	1.95	0.48
1:A:343:GLY:HA3	1:B:302:GLU:HB2	1.95	0.47
1:A:222:LEU:HB3	1:A:241:ILE:HG22	1.95	0.47
1:D:33:ASN:CA	1:D:34:ASP:HB2	2.45	0.47
1:B:232:ALA:HB2	1:B:309:THR:HG22	1.95	0.47
1:D:114:VAL:O	1:D:115:LYS:C	2.59	0.46
1:B:223:VAL:HG13	1:B:238:ILE:HG23	1.96	0.46
1:A:143:ARG:HH11	1:A:143:ARG:HG2	1.82	0.45
1:A:48:ASN:HD22	1:A:48:ASN:C	2.25	0.45
1:A:59:ARG:NH1	1:A:130:LEU:HD12	2.33	0.44
1:A:80:PHE:HB2	1:A:108:LYS:HB2	1.99	0.44
1:C:68:LEU:HD21	1:C:110:ILE:HG21	1.99	0.43
1:D:373:THR:HG21	1:D:398:LYS:HE3	2.01	0.43
1:B:250:THR:O	1:B:281:MET:HA	2.20	0.42
1:A:77:VAL:HG22	1:A:93:ASP:O	2.19	0.42
1:D:115:LYS:HA	1:D:116:GLY:HA2	1.68	0.42
1:A:173:ASP:N	1:A:174:PRO:CD	2.83	0.42
1:B:234:VAL:HG22	1:B:235:ASP:N	2.35	0.41
1:D:109:ILE:O	1:D:120:SER:HA	2.19	0.41
1:D:333:PHE:HB3	1:D:407:CYS:HB2	2.02	0.41
1:D:49:GLY:H	1:D:50:GLN:HA	1.85	0.41
1:C:224:VAL:HG13	1:C:239:ILE:HG23	2.03	0.41
1:A:250:THR:O	1:A:281:MET:HA	2.21	0.41
1:C:44:PHE:HD2	1:C:122:ARG:HH11	1.69	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	397/420 (94%)	380 (96%)	17 (4%)	0	100	100
1	B	381/420 (91%)	361 (95%)	19 (5%)	1 (0%)	36	59
1	C	388/420 (92%)	369 (95%)	14 (4%)	5 (1%)	9	23
1	D	386/420 (92%)	360 (93%)	23 (6%)	3 (1%)	16	35
All	All	1552/1680 (92%)	1470 (95%)	73 (5%)	9 (1%)	21	42

All (9) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	C	34	ASN
1	C	116	LYS
1	C	261	SER
1	D	115	LYS
1	D	117	LYS
1	C	36	ASP
1	D	34	ASP
1	B	234	VAL
1	C	117	GLY

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	340/358 (95%)	333 (98%)	7 (2%)	47	73
1	B	328/358 (92%)	317 (97%)	11 (3%)	32	60
1	C	332/358 (93%)	325 (98%)	7 (2%)	47	73
1	D	330/358 (92%)	323 (98%)	7 (2%)	47	73
All	All	1330/1432 (93%)	1298 (98%)	32 (2%)	43	70

All (32) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	23	GLU
1	A	59	ARG

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	A	77	VAL
1	A	115	LYS
1	A	120	SER
1	A	143	ARG
1	A	408	VAL
1	B	37	HIS
1	B	59	ARG
1	B	69	VAL
1	B	70	THR
1	B	87	GLU
1	B	97	ASN
1	B	170	THR
1	B	244	GLU
1	B	297	ASN
1	B	322	TRP
1	B	408	VAL
1	C	33	GLU
1	C	64	LEU
1	C	73	LYS
1	C	216	VAL
1	C	323	TRP
1	C	346	THR
1	C	399	LYS
1	D	48	ASN
1	D	50	GLN
1	D	106	ASN
1	D	107	LEU
1	D	215	VAL
1	D	287	SER
1	D	322	TRP

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (29) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	48	ASN
1	A	62	ASN
1	A	97	ASN
1	A	128	ASN
1	A	157	ASN
1	A	203	GLN
1	A	209	GLN
1	A	393	GLN

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	A	400	GLN
1	B	62	ASN
1	B	128	ASN
1	B	157	ASN
1	B	261	ASN
1	B	295	ASN
1	B	303	GLN
1	B	330	ASN
1	C	38	HIS
1	C	63	ASN
1	C	158	ASN
1	C	161	ASN
1	C	204	GLN
1	C	210	GLN
1	C	296	ASN
1	C	304	GLN
1	C	394	GLN
1	D	157	ASN
1	D	261	ASN
1	D	297	ASN
1	D	400	GLN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

15 monosaccharides are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
2	BGC	E	1	-	12,12,12	0.44	0	17,17,17	0.79	0
2	BGC	E	2	-	11,11,12	0.19	0	15,15,17	0.82	1 (6%)
2	BGC	E	3	-	11,11,12	0.26	0	15,15,17	0.58	0
2	BGC	E	4	-	11,11,12	0.29	0	15,15,17	0.59	0
2	BGC	E	5	-	11,11,12	0.25	0	15,15,17	0.61	0
2	BGC	F	1	-	12,12,12	0.57	0	17,17,17	1.01	2 (11%)
2	BGC	F	2	-	11,11,12	0.36	0	15,15,17	0.41	0
2	BGC	F	3	-	11,11,12	0.41	0	15,15,17	0.59	0
2	BGC	F	4	-	11,11,12	0.33	0	15,15,17	0.83	1 (6%)
2	BGC	F	5	-	11,11,12	0.43	0	15,15,17	0.77	0
2	BGC	G	1	-	12,12,12	0.61	0	17,17,17	0.90	0
2	BGC	G	2	-	11,11,12	0.47	0	15,15,17	0.71	0
2	BGC	G	3	-	11,11,12	0.43	0	15,15,17	0.64	0
2	BGC	G	4	-	11,11,12	0.47	0	15,15,17	0.96	1 (6%)
2	BGC	G	5	-	11,11,12	0.42	0	15,15,17	0.94	1 (6%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	BGC	E	1	-	-	2/2/22/22	0/1/1/1
2	BGC	E	2	-	-	1/2/19/22	0/1/1/1
2	BGC	E	3	-	-	0/2/19/22	0/1/1/1
2	BGC	E	4	-	-	0/2/19/22	0/1/1/1
2	BGC	E	5	-	-	0/2/19/22	0/1/1/1
2	BGC	F	1	-	-	2/2/22/22	0/1/1/1
2	BGC	F	2	-	-	1/2/19/22	0/1/1/1
2	BGC	F	3	-	-	2/2/19/22	0/1/1/1
2	BGC	F	4	-	-	2/2/19/22	0/1/1/1
2	BGC	F	5	-	-	2/2/19/22	0/1/1/1
2	BGC	G	1	-	-	2/2/22/22	0/1/1/1
2	BGC	G	2	-	-	1/2/19/22	0/1/1/1
2	BGC	G	3	-	-	0/2/19/22	0/1/1/1
2	BGC	G	4	-	-	2/2/19/22	0/1/1/1
2	BGC	G	5	-	-	1/2/19/22	0/1/1/1

There are no bond length outliers.

All (6) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	F	1	BGC	C3-C4-C5	2.58	114.91	110.23
2	G	4	BGC	C1-O5-C5	2.48	115.51	112.19
2	G	5	BGC	C1-C2-C3	2.39	113.12	109.64
2	F	4	BGC	C1-O5-C5	2.24	115.19	112.19
2	F	1	BGC	O5-C5-C4	2.13	113.55	109.70
2	E	2	BGC	O5-C5-C6	2.09	111.73	107.66

There are no chirality outliers.


All (18) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	F	5	BGC	O5-C5-C6-O6
2	F	4	BGC	C4-C5-C6-O6
2	F	3	BGC	C4-C5-C6-O6
2	F	4	BGC	O5-C5-C6-O6
2	E	1	BGC	C4-C5-C6-O6
2	F	5	BGC	C4-C5-C6-O6
2	E	1	BGC	O5-C5-C6-O6
2	G	1	BGC	O5-C5-C6-O6
2	F	3	BGC	O5-C5-C6-O6
2	F	1	BGC	C4-C5-C6-O6
2	G	1	BGC	C4-C5-C6-O6
2	G	4	BGC	O5-C5-C6-O6
2	F	2	BGC	O5-C5-C6-O6
2	F	1	BGC	O5-C5-C6-O6
2	G	2	BGC	O5-C5-C6-O6
2	G	5	BGC	O5-C5-C6-O6
2	G	4	BGC	C4-C5-C6-O6
2	E	2	BGC	O5-C5-C6-O6

There are no ring outliers.

No monomer is involved in short contacts.

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for oligosaccharide.



5.6 Ligand geometry [i](#)

Of 21 ligands modelled in this entry, 2 are monoatomic - leaving 19 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
3	EDO	B	504	-	3,3,3	0.44	0	2,2,2	0.24	0
4	PO4	A	510	-	4,4,4	0.84	0	6,6,6	0.54	0
4	PO4	A	511	-	4,4,4	0.89	0	6,6,6	0.45	0
4	PO4	C	507	-	4,4,4	0.85	0	6,6,6	0.45	0
3	EDO	C	511	-	3,3,3	0.46	0	2,2,2	0.27	0
3	EDO	A	507	-	3,3,3	0.49	0	2,2,2	0.27	0
4	PO4	D	508	-	4,4,4	0.92	0	6,6,6	0.45	0
4	PO4	D	509	-	4,4,4	0.87	0	6,6,6	0.51	0
4	PO4	B	501	-	4,4,4	0.97	0	6,6,6	0.49	0
4	PO4	C	508	-	4,4,4	0.95	0	6,6,6	0.45	0
4	PO4	A	508	-	4,4,4	0.89	0	6,6,6	0.48	0
4	PO4	C	509	-	4,4,4	0.90	0	6,6,6	0.47	0
4	PO4	A	509	-	4,4,4	0.94	0	6,6,6	0.49	0
3	EDO	A	506	-	3,3,3	0.45	0	2,2,2	0.34	0
4	PO4	B	502	-	4,4,4	0.91	0	6,6,6	0.44	0
4	PO4	C	506	-	4,4,4	0.98	0	6,6,6	0.38	0
4	PO4	B	503	-	4,4,4	0.81	0	6,6,6	0.49	0
3	EDO	D	506	-	3,3,3	0.47	0	2,2,2	0.32	0
4	PO4	D	507	-	4,4,4	0.87	0	6,6,6	0.47	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	EDO	B	504	-	-	0/1/1/1	-
3	EDO	A	507	-	-	1/1/1/1	-
3	EDO	D	506	-	-	1/1/1/1	-
3	EDO	C	511	-	-	0/1/1/1	-
3	EDO	A	506	-	-	0/1/1/1	-

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

All (2) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
3	A	507	EDO	O1-C1-C2-O2
3	D	506	EDO	O1-C1-C2-O2

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	399/420 (95%)	-0.21	7 (1%) 67 65	26, 49, 82, 100	0
1	B	385/420 (91%)	0.26	16 (4%) 40 37	30, 65, 105, 123	0
1	C	390/420 (92%)	-0.09	5 (1%) 75 73	32, 58, 86, 108	0
1	D	388/420 (92%)	-0.01	8 (2%) 63 60	33, 55, 92, 103	0
All	All	1562/1680 (92%)	-0.01	36 (2%) 61 58	26, 56, 93, 123	0

All (36) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	D	115	LYS	4.8
1	A	22	THR	4.8
1	B	232	ALA	4.6
1	B	114	VAL	4.2
1	C	32	THR	3.9
1	B	234	VAL	3.6
1	C	116	LYS	3.5
1	D	33	ASN	3.3
1	B	260	SER	3.2
1	C	118	LYS	3.1
1	B	119	THR	3.1
1	A	31	THR	3.1
1	B	76	THR	3.0
1	D	74	TYR	2.9
1	B	74	TYR	2.9
1	D	261	ASN	2.8
1	D	49	GLY	2.7
1	B	261	ASN	2.6
1	B	37	HIS	2.6
1	D	173	ASP	2.5
1	B	69	VAL	2.5

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	30	VAL	2.4
1	D	251	ILE	2.3
1	B	89	GLY	2.3
1	D	256	VAL	2.3
1	A	72	LYS	2.3
1	B	88	SER	2.2
1	B	241	ILE	2.2
1	B	47	THR	2.1
1	C	120	THR	2.1
1	B	118	LYS	2.1
1	A	29	THR	2.1
1	C	50	GLY	2.1
1	B	48	ASN	2.0
1	A	252	GLY	2.0
1	A	49	GLY	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q<0.9' lists the number of atoms with occupancy less than 0.9.


Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
2	BGC	F	1	12/12	0.55	0.15	87,107,112,117	0
2	BGC	G	1	12/12	0.57	0.14	99,109,116,121	0
2	BGC	G	2	11/12	0.59	0.14	90,93,97,97	0
2	BGC	G	3	11/12	0.74	0.15	80,84,88,91	0
2	BGC	F	2	11/12	0.82	0.13	85,92,98,98	0
2	BGC	G	5	11/12	0.88	0.12	80,82,86,87	0
2	BGC	F	5	11/12	0.89	0.13	76,79,84,85	0
2	BGC	G	4	11/12	0.90	0.12	73,76,80,85	0
2	BGC	F	3	11/12	0.92	0.10	82,85,90,92	0
2	BGC	F	4	11/12	0.93	0.09	75,79,81,83	0
2	BGC	E	5	11/12	0.95	0.07	42,45,49,52	0
2	BGC	E	3	11/12	0.96	0.07	38,39,40,41	0
2	BGC	E	1	12/12	0.96	0.07	41,45,53,54	0

Continued on next page...

Continued from previous page...




Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
2	BGC	E	2	11/12	0.96	0.07	37,40,42,42	0
2	BGC	E	4	11/12	0.97	0.06	37,39,41,41	0

The following is a graphical depiction of the model fit to experimental electron density for oligosaccharide. Each fit is shown from different orientation to approximate a three-dimensional view.






Electron density around Chain F:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)

**Electron density around Chain G:**

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
3	EDO	C	511	4/4	0.62	0.32	73,75,77,77	0
3	EDO	A	506	4/4	0.72	0.24	75,77,78,79	0
3	EDO	B	504	4/4	0.83	0.21	61,63,65,65	0
4	PO4	B	502	5/5	0.85	0.17	88,89,95,98	0
4	PO4	D	509	5/5	0.88	0.22	87,97,105,108	0
3	EDO	D	506	4/4	0.89	0.19	59,60,60,62	0
4	PO4	A	510	5/5	0.91	0.13	93,100,105,107	0
3	EDO	A	507	4/4	0.92	0.16	55,61,62,65	0
4	PO4	A	511	5/5	0.93	0.18	76,82,86,91	0
4	PO4	C	509	5/5	0.94	0.17	86,88,91,94	0
4	PO4	C	507	5/5	0.94	0.15	76,76,89,90	0
4	PO4	D	508	5/5	0.95	0.20	85,93,97,101	0
4	PO4	B	501	5/5	0.95	0.13	68,72,75,77	0
5	NA	A	512	1/1	0.95	0.12	59,59,59,59	0
4	PO4	A	509	5/5	0.96	0.11	59,69,76,78	0
4	PO4	B	503	5/5	0.97	0.15	52,58,60,62	0
4	PO4	D	507	5/5	0.97	0.10	50,51,54,54	0
4	PO4	C	506	5/5	0.97	0.11	43,48,51,51	0
4	PO4	A	508	5/5	0.97	0.09	57,61,63,65	0
4	PO4	C	508	5/5	0.97	0.10	79,81,86,87	0
5	NA	C	510	1/1	0.97	0.09	54,54,54,54	0

6.5 Other polymers [i](#)

There are no such residues in this entry.