



wwPDB X-ray Structure Validation Summary Report ⓘ

Mar 17, 2026 – 09:02 PM UTC

PDB ID : 6E5B / pdb_00006e5b
Title : Human Immunoproteasome 20S particle in complex with compound 1
Authors : Steinbacher, S.; Augustin, M.; Blaesse, M.; Harris, S.F.
Deposited on : 2018-07-19
Resolution : 2.77 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Buster-report	:	wwPDB partial adaption of 1.1.7 (2018)
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.77 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	5248 (2.80-2.76)
Clashscore	190562	5693 (2.80-2.76)
Ramachandran outliers	187476	5590 (2.80-2.76)
Sidechain outliers	187428	5592 (2.80-2.76)
RSRZ outliers	180081	5251 (2.80-2.76)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	234	3% 96% ..
1	O	234	3% 96% ..
2	B	261	7% 93% 6%
2	P	261	2% 94% 6%
3	C	248	4% 89% 8%

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
3	Q	248	 5% 93% 6%
4	D	241	 5% 94% 5%
4	R	241	 6% 94% 2%
5	E	263	 4% 87% 10%
5	S	263	 4% 87% 10%
6	F	255	 3% 94% 2%
6	T	255	 4% 94% 5%
7	G	246	 4% 96% 2%
7	U	246	 4% 97% 2%
8	H	273	 2% 79% 20%
8	V	273	 3% 79% 20%
9	I	205	 1% 99% 0%
9	W	205	 4% 99% 1%
10	J	201	 2% 98% 2%
10	X	201	 3% 98% 2%
11	K	276	 3% 72% 26%
11	Y	276	 1% 71% 26%
12	L	241	 2% 85% 12%
12	Z	241	 1% 85% 12%
13	M	264	 0% 81% 19%
13	a	264	 1% 81% 19%
14	N	219	 1% 90% 9%
14	b	219	 0% 90% 9%

2 Entry composition

There are 18 unique types of molecules in this entry. The entry contains 48431 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Proteasome subunit alpha type-2.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	229	Total	C	N	O	S	22	0	0
			1787	1144	301	336	6			
1	O	229	Total	C	N	O	S	22	0	0
			1787	1144	301	336	6			

- Molecule 2 is a protein called Proteasome subunit alpha type-4.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	B	245	Total	C	N	O	S	40	0	0
			1932	1221	332	369	10			
2	P	246	Total	C	N	O	S	16	0	0
			1940	1225	333	372	10			

- Molecule 3 is a protein called Proteasome subunit alpha type-7.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
3	C	227	Total	C	N	O	S	17	0	0
			1798	1132	318	343	5			
3	Q	234	Total	C	N	O	S	23	0	0
			1848	1160	327	356	5			

- Molecule 4 is a protein called Proteasome subunit alpha type-5.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
4	D	230	Total	C	N	O	S	21	0	0
			1758	1103	291	353	11			
4	R	232	Total	C	N	O	S	12	0	0
			1769	1110	293	355	11			

- Molecule 5 is a protein called Proteasome subunit alpha type-1.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
5	E	236	Total	C	N	O	S	0	0	0
			1857	1162	334	350	11			
5	S	236	Total	C	N	O	S	6	0	0
			1857	1162	334	350	11			

- Molecule 6 is a protein called Proteasome subunit alpha type-3.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
6	F	244	Total	C	N	O	S	21	0	0
			1911	1209	325	366	11			
6	T	241	Total	C	N	O	S	5	0	0
			1885	1195	322	357	11			

- Molecule 7 is a protein called Proteasome subunit alpha type-6.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
7	G	241	Total	C	N	O	S	13	0	0
			1875	1191	311	360	13			
7	U	242	Total	C	N	O	S	17	0	0
			1886	1197	315	361	13			

- Molecule 8 is a protein called Proteasome subunit beta type-10.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
8	H	218	Total	C	N	O	S	7	0	0
			1601	1004	284	305	8			
8	V	218	Total	C	N	O	S	7	0	0
			1601	1004	284	305	8			

- Molecule 9 is a protein called Proteasome subunit beta type-3.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
9	I	204	Total	C	N	O	S	7	0	0
			1591	1013	265	294	19			
9	W	204	Total	C	N	O	S	0	0	0
			1591	1013	265	294	19			

- Molecule 10 is a protein called Proteasome subunit beta type-2.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
10	J	196	Total	C	N	O	S	7	0	0
			1571	1006	267	289	9			

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
10	X	197	Total	C	N	O	S	13	0	0
			1578	1011	268	290	9			

- Molecule 11 is a protein called Proteasome subunit beta type-8.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
11	K	203	Total	C	N	O	S	0	0	0
			1576	984	276	301	15			
11	Y	203	Total	C	N	O	S	0	0	0
			1576	984	276	301	15			

- Molecule 12 is a protein called Proteasome subunit beta type-1.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
12	L	213	Total	C	N	O	S	6	0	0
			1654	1047	284	313	10			
12	Z	213	Total	C	N	O	S	4	0	0
			1654	1047	284	313	10			


- Molecule 13 is a protein called Proteasome subunit beta type-4.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
13	M	214	Total	C	N	O	S	2	0	0
			1673	1055	289	317	12			
13	a	214	Total	C	N	O	S	0	0	0
			1673	1055	289	317	12			

- Molecule 14 is a protein called Proteasome subunit beta type-9.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
14	N	199	Total	C	N	O	S	3	0	0
			1493	939	254	291	9			
14	b	199	Total	C	N	O	S	7	0	0
			1493	939	254	291	9			

- Molecule 15 is THIOCYANATE ION (CCD ID: SCN) (formula: CNS).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
15	E	1	Total	C	N	S	0	0
			3	1	1	1		
15	L	1	Total	C	N	S	0	0
			3	1	1	1		
15	Z	1	Total	C	N	S	0	0
			3	1	1	1		

- Molecule 16 is SODIUM ION (CCD ID: NA) (formula: Na).


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
16	G	1	Total	Na	0	0
			1	1		
16	H	1	Total	Na	0	0
			1	1		
16	I	2	Total	Na	0	0
			2	2		
16	K	1	Total	Na	0	0
			1	1		
16	L	2	Total	Na	0	0
			2	2		
16	N	1	Total	Na	0	0
			1	1		
16	U	1	Total	Na	0	0
			1	1		
16	W	1	Total	Na	0	0
			1	1		
16	Z	1	Total	Na	0	0
			1	1		

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
16	b	1	Total	Na	0	0
			1	1		

- Molecule 17 is [(1R)-3-methyl-1-[[1-[2-[(2,3,4-trimethoxyphenyl)carbonylamino]ethyl]-1,2,3-triazol-4-yl]carbonylamino]butyl]-tris(oxidanyl)boron (CCD ID: HUJ) (formula: C₂₀H₃₁BN₅O₈) (labeled as "Ligand of Interest" by depositor).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
17	K	1	Total	B	C	N	O	2	0
			33	1	20	5	7		
17	N	1	Total	B	C	N	O	11	0
			33	1	20	5	7		
17	Y	1	Total	B	C	N	O	2	0
			33	1	20	5	7		
17	b	1	Total	B	C	N	O	0	0
			33	1	20	5	7		

- Molecule 18 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
18	B	2	Total	O	0	0
			2	2		
18	C	1	Total	O	0	0
			1	1		
18	E	1	Total	O	0	0
			1	1		

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
18	F	1	Total O 1 1	0	0
18	H	2	Total O 2 2	0	0
18	I	4	Total O 4 4	0	0
18	J	1	Total O 1 1	0	0
18	K	4	Total O 4 4	0	0
18	L	2	Total O 2 2	0	0
18	M	4	Total O 4 4	0	0
18	N	3	Total O 3 3	0	0
18	O	1	Total O 1 1	0	0
18	P	2	Total O 2 2	0	0
18	Q	1	Total O 1 1	0	0
18	R	4	Total O 4 4	0	0
18	S	4	Total O 4 4	0	0
18	T	2	Total O 2 2	0	0
18	U	1	Total O 1 1	0	0
18	V	5	Total O 5 5	0	0
18	W	3	Total O 3 3	0	0
18	X	3	Total O 3 3	0	0
18	Y	5	Total O 5 5	0	0
18	Z	4	Total O 4 4	0	0
18	a	2	Total O 2 2	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
18	b	1	Total	O	0	0
			1	1		


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Proteasome subunit alpha type-2





- Molecule 1: Proteasome subunit alpha type-2




- Molecule 2: Proteasome subunit alpha type-4




- Molecule 2: Proteasome subunit alpha type-4




- Molecule 3: Proteasome subunit alpha type-7




- Molecule 3: Proteasome subunit alpha type-7





- Molecule 4: Proteasome subunit alpha type-5





- Molecule 4: Proteasome subunit alpha type-5



- Molecule 5: Proteasome subunit alpha type-1




- Molecule 5: Proteasome subunit alpha type-1




GLU
HIS

- Molecule 6: Proteasome subunit alpha type-3



- Molecule 6: Proteasome subunit alpha type-3








- Molecule 7: Proteasome subunit alpha type-6






- Molecule 7: Proteasome subunit alpha type-6



- Molecule 8: Proteasome subunit beta type-10




- Molecule 8: Proteasome subunit beta type-10




- Molecule 9: Proteasome subunit beta type-3




- Molecule 9: Proteasome subunit beta type-3




- Molecule 10: Proteasome subunit beta type-2




- Molecule 10: Proteasome subunit beta type-2




- Molecule 11: Proteasome subunit beta type-8





- Molecule 11: Proteasome subunit beta type-8





- Molecule 12: Proteasome subunit beta type-1





- Molecule 12: Proteasome subunit beta type-1





• Molecule 13: Proteasome subunit beta type-4





• Molecule 13: Proteasome subunit beta type-4



• Molecule 14: Proteasome subunit beta type-9



• Molecule 14: Proteasome subunit beta type-9



4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	118.57Å 203.14Å 161.49Å 90.00° 106.82° 90.00°	Depositor
Resolution (Å)	154.58 – 2.77 154.58 – 2.77	Depositor EDS
% Data completeness (in resolution range)	97.0 (154.58-2.77) 97.0 (154.58-2.77)	Depositor EDS
R_{merge}	0.08	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.26 (at 2.77Å)	Xtriage
Refinement program	REFMAC 5.8.0155	Depositor
R, R_{free}	0.248 , 0.291 0.248 , 0.293	Depositor DCC
R_{free} test set	646 reflections (0.35%)	wwPDB-VP
Wilson B-factor (Å ²)	54.0	Xtriage
Anisotropy	0.693	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.28 , 29.4	EDS
L-test for twinning ²	$\langle L \rangle = 0.45$, $\langle L^2 \rangle = 0.28$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.91	EDS
Total number of atoms	48431	wwPDB-VP
Average B, all atoms (Å ²)	73.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 6.09% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: HUI, NA, SCN

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.79	0/1826	1.04	0/2474
1	O	0.79	0/1826	1.04	0/2474
2	B	0.76	0/1961	1.01	2/2640 (0.1%)
2	P	0.76	0/1969	1.00	1/2651 (0.0%)
3	C	0.83	0/1822	1.04	0/2458
3	Q	0.83	1/1873 (0.1%)	1.04	0/2526
4	D	0.80	0/1783	1.01	0/2406
4	R	0.80	0/1796	1.01	1/2426 (0.0%)
5	E	0.78	1/1891 (0.1%)	1.00	0/2555
5	S	0.78	1/1891 (0.1%)	1.00	0/2555
6	F	0.79	0/1946	1.02	0/2620
6	T	0.78	0/1920	1.02	0/2585
7	G	0.77	0/1909	1.00	0/2581
7	U	0.76	0/1920	1.00	0/2595
8	H	0.76	0/1625	1.01	2/2207 (0.1%)
8	V	0.76	0/1625	1.01	0/2207
9	I	0.75	0/1620	0.97	0/2184
9	W	0.75	0/1620	0.97	0/2184
10	J	0.79	0/1603	0.96	0/2168
10	X	0.79	0/1611	0.96	0/2180
11	K	0.76	0/1609	0.99	0/2170
11	Y	0.76	0/1609	0.99	0/2170
12	L	0.77	0/1684	0.99	2/2268 (0.1%)
12	Z	0.76	0/1684	0.99	2/2268 (0.1%)
13	M	0.77	0/1706	0.97	0/2309
13	a	0.76	0/1706	0.97	0/2309
14	N	0.77	0/1521	0.99	0/2062
14	b	0.76	0/1521	1.00	0/2062
All	All	0.78	3/49077 (0.0%)	1.00	10/66294 (0.0%)

All (3) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
5	E	226	ASP	CA-C	6.11	1.55	1.52
5	S	226	ASP	CA-C	5.91	1.55	1.52
3	Q	117	ARG	CD-NE	5.55	1.54	1.46

The worst 5 of 10 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	B	226	ARG	CD-NE-CZ	6.73	133.82	124.40
4	R	238	ILE	N-CA-C	5.68	121.15	109.34
2	P	32	GLY	N-CA-C	5.39	118.12	111.93
12	Z	102	PHE	CA-C-N	5.25	126.40	119.84
12	Z	102	PHE	C-N-CA	5.25	126.40	119.84

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1787	0	1782	2	0
1	O	1787	0	1782	2	0
2	B	1932	0	1947	0	0
2	P	1940	0	1951	0	0
3	C	1798	0	1820	2	0
3	Q	1848	0	1865	1	0
4	D	1758	0	1742	1	0
4	R	1769	0	1753	2	0
5	E	1857	0	1845	1	0
5	S	1857	0	1845	1	0
6	F	1911	0	1887	4	0
6	T	1885	0	1871	1	0
7	G	1875	0	1879	1	0
7	U	1886	0	1892	1	0
8	H	1601	0	1634	1	0
8	V	1601	0	1634	1	0
9	I	1591	0	1609	0	0
9	W	1591	0	1609	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
10	J	1571	0	1573	0	0
10	X	1578	0	1580	0	0
11	K	1576	0	1522	1	0
11	Y	1576	0	1522	1	0
12	L	1654	0	1656	3	0
12	Z	1654	0	1656	3	0
13	M	1673	0	1650	0	0
13	a	1673	0	1650	0	0
14	N	1493	0	1457	0	0
14	b	1493	0	1457	0	0
15	E	3	0	0	0	0
15	L	3	0	0	0	0
15	Z	3	0	0	0	0
16	G	1	0	0	0	0
16	H	1	0	0	0	0
16	I	2	0	0	0	0
16	K	1	0	0	0	0
16	L	2	0	0	0	0
16	N	1	0	0	0	0
16	U	1	0	0	0	0
16	W	1	0	0	0	0
16	Z	1	0	0	0	0
16	b	1	0	0	0	0
17	K	33	0	0	0	0
17	N	33	0	0	0	0
17	Y	33	0	0	0	0
17	b	33	0	0	0	0
18	B	2	0	0	0	0
18	C	1	0	0	0	0
18	E	1	0	0	0	0
18	F	1	0	0	0	0
18	H	2	0	0	0	0
18	I	4	0	0	0	0
18	J	1	0	0	0	0
18	K	4	0	0	0	0
18	L	2	0	0	0	0
18	M	4	0	0	0	0
18	N	3	0	0	0	0
18	O	1	0	0	0	0
18	P	2	0	0	0	0
18	Q	1	0	0	0	0
18	R	4	0	0	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
18	S	4	0	0	0	0
18	T	2	0	0	0	0
18	U	1	0	0	0	0
18	V	5	0	0	0	0
18	W	3	0	0	0	0
18	X	3	0	0	0	0
18	Y	5	0	0	0	0
18	Z	4	0	0	0	0
18	a	2	0	0	0	0
18	b	1	0	0	0	0
All	All	48431	0	48070	29	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 0.

The worst 5 of 29 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
12:L:145:LEU:HD21	12:L:182:ALA:HB2	1.78	0.65
12:Z:145:LEU:HD21	12:Z:182:ALA:HB2	1.79	0.64
1:A:110:VAL:HG22	1:A:135:ILE:HD13	1.85	0.59
1:O:110:VAL:HG22	1:O:135:ILE:HD13	1.85	0.58
4:R:107:MET:HE3	4:R:112:VAL:HG22	1.87	0.55

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	227/234 (97%)	215 (95%)	11 (5%)	1 (0%)	30 57
1	O	227/234 (97%)	215 (95%)	12 (5%)	0	100 100

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
2	B	241/261 (92%)	233 (97%)	8 (3%)	0	100	100
2	P	242/261 (93%)	234 (97%)	8 (3%)	0	100	100
3	C	221/248 (89%)	212 (96%)	9 (4%)	0	100	100
3	Q	230/248 (93%)	220 (96%)	10 (4%)	0	100	100
4	D	226/241 (94%)	212 (94%)	12 (5%)	2 (1%)	14	37
4	R	230/241 (95%)	213 (93%)	15 (6%)	2 (1%)	14	37
5	E	234/263 (89%)	221 (94%)	12 (5%)	1 (0%)	30	57
5	S	234/263 (89%)	223 (95%)	10 (4%)	1 (0%)	30	57
6	F	242/255 (95%)	237 (98%)	5 (2%)	0	100	100
6	T	239/255 (94%)	235 (98%)	4 (2%)	0	100	100
7	G	239/246 (97%)	234 (98%)	5 (2%)	0	100	100
7	U	240/246 (98%)	236 (98%)	4 (2%)	0	100	100
8	H	214/273 (78%)	205 (96%)	9 (4%)	0	100	100
8	V	214/273 (78%)	203 (95%)	11 (5%)	0	100	100
9	I	202/205 (98%)	192 (95%)	10 (5%)	0	100	100
9	W	202/205 (98%)	193 (96%)	8 (4%)	1 (0%)	24	52
10	J	194/201 (96%)	188 (97%)	6 (3%)	0	100	100
10	X	195/201 (97%)	189 (97%)	6 (3%)	0	100	100
11	K	201/276 (73%)	193 (96%)	7 (4%)	1 (0%)	24	52
11	Y	201/276 (73%)	192 (96%)	8 (4%)	1 (0%)	24	52
12	L	211/241 (88%)	204 (97%)	7 (3%)	0	100	100
12	Z	211/241 (88%)	204 (97%)	7 (3%)	0	100	100
13	M	212/264 (80%)	203 (96%)	9 (4%)	0	100	100
13	a	212/264 (80%)	203 (96%)	9 (4%)	0	100	100
14	N	197/219 (90%)	187 (95%)	10 (5%)	0	100	100
14	b	197/219 (90%)	187 (95%)	10 (5%)	0	100	100
All	All	6135/6854 (90%)	5883 (96%)	242 (4%)	10 (0%)	43	70

5 of 10 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
4	R	238	ILE
4	D	239	LYS

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
5	E	235	GLY
11	K	132	ASN
11	Y	132	ASN

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	188/191 (98%)	188 (100%)	0	100	100
1	O	188/191 (98%)	187 (100%)	1 (0%)	81	92
2	B	205/221 (93%)	205 (100%)	0	100	100
2	P	206/221 (93%)	206 (100%)	0	100	100
3	C	193/211 (92%)	191 (99%)	2 (1%)	68	86
3	Q	198/211 (94%)	197 (100%)	1 (0%)	81	92
4	D	193/203 (95%)	192 (100%)	1 (0%)	81	92
4	R	194/203 (96%)	194 (100%)	0	100	100
5	E	202/224 (90%)	200 (99%)	2 (1%)	68	86
5	S	202/224 (90%)	200 (99%)	2 (1%)	68	86
6	F	201/212 (95%)	200 (100%)	1 (0%)	81	92
6	T	198/212 (93%)	197 (100%)	1 (0%)	81	92
7	G	205/210 (98%)	203 (99%)	2 (1%)	68	86
7	U	206/210 (98%)	204 (99%)	2 (1%)	68	86
8	H	170/217 (78%)	170 (100%)	0	100	100
8	V	170/217 (78%)	170 (100%)	0	100	100
9	I	173/174 (99%)	172 (99%)	1 (1%)	78	91
9	W	173/174 (99%)	172 (99%)	1 (1%)	78	91
10	J	167/171 (98%)	167 (100%)	0	100	100
10	X	168/171 (98%)	167 (99%)	1 (1%)	78	91
11	K	165/222 (74%)	163 (99%)	2 (1%)	63	84

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
11	Y	165/222 (74%)	162 (98%)	3 (2%)	51	79
12	L	178/199 (89%)	176 (99%)	2 (1%)	65	85
12	Z	178/199 (89%)	176 (99%)	2 (1%)	65	85
13	M	177/215 (82%)	177 (100%)	0	100	100
13	a	177/215 (82%)	177 (100%)	0	100	100
14	N	153/166 (92%)	152 (99%)	1 (1%)	76	90
14	b	153/166 (92%)	152 (99%)	1 (1%)	76	90
All	All	5146/5672 (91%)	5117 (99%)	29 (1%)	78	91

5 of 29 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	O	58	GLU
12	Z	194	ARG
5	S	197	GLU
11	Y	116	GLU
5	S	83	LEU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 59 such sidechains are listed below:

Mol	Chain	Res	Type
13	M	208	ASN
12	Z	131	GLN
4	R	211	ASN
11	Y	175	ASN
10	X	55	GLN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 19 ligands modelled in this entry, 12 are monoatomic - leaving 7 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# $ Z > 2$	Counts	RMSZ	# $ Z > 2$
17	HUJ	Y	901	11	30,34,35	1.35	2 (6%)	40,46,49	1.58	11 (27%)
15	SCN	L	301	-	1,2,2	0.19	0	0,1,1	-	-
17	HUJ	N	901	14	30,34,35	1.54	5 (16%)	40,46,49	1.69	9 (22%)
17	HUJ	b	901	14	30,34,35	1.22	3 (10%)	40,46,49	1.59	9 (22%)
15	SCN	Z	301	-	1,2,2	0.23	0	0,1,1	-	-
15	SCN	E	301	-	1,2,2	0.50	0	0,1,1	-	-
17	HUJ	K	901	11	30,34,35	1.35	2 (6%)	40,46,49	1.56	9 (22%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
17	HUJ	N	901	14	-	13/27/32/34	0/2/2/2
17	HUJ	Y	901	11	-	11/27/32/34	0/2/2/2
17	HUJ	K	901	11	-	7/27/32/34	0/2/2/2
17	HUJ	b	901	14	-	10/27/32/34	0/2/2/2

The worst 5 of 12 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
17	Y	901	HUJ	N3-N2	5.54	1.41	1.32
17	K	901	HUJ	N3-N2	5.52	1.41	1.32
17	N	901	HUJ	N3-N2	5.39	1.40	1.32

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
17	b	901	HUJ	N3-N2	5.22	1.40	1.32
17	N	901	HUJ	C12-C11	-4.47	1.33	1.39

The worst 5 of 38 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
17	b	901	HUJ	C8-N1-N2	-4.10	112.55	120.78
17	b	901	HUJ	C8-N1-C7	4.09	136.48	128.83
17	N	901	HUJ	C8-N1-C7	3.87	136.06	128.83
17	N	901	HUJ	C8-N1-N2	-3.61	113.53	120.78
17	Y	901	HUJ	C6-C5-N	3.48	120.17	115.22

There are no chirality outliers.



5 of 41 torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
17	N	901	HUJ	N-C5-C6-C7
17	N	901	HUJ	N-C5-C6-N3
17	N	901	HUJ	O-C5-C6-C7
17	N	901	HUJ	O-C5-C6-N3
17	N	901	HUJ	C9-C8-N1-C7

There are no ring outliers.

No monomer is involved in short contacts.

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.





5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2			OWAB(Å ²)	Q<0.9
1	A	229/234 (97%)	0.68	7 (3%)	51	44	47, 72, 112, 125	6 (2%)
1	O	229/234 (97%)	0.67	6 (2%)	57	49	34, 70, 107, 141	5 (2%)
2	B	245/261 (93%)	0.72	17 (6%)	23	18	45, 74, 117, 176	12 (4%)
2	P	246/261 (94%)	0.64	4 (1%)	70	63	41, 75, 111, 166	5 (2%)
3	C	227/248 (91%)	0.78	10 (4%)	39	33	48, 79, 125, 175	5 (2%)
3	Q	234/248 (94%)	0.71	12 (5%)	33	27	37, 78, 122, 161	6 (2%)
4	D	230/241 (95%)	0.68	11 (4%)	35	30	44, 79, 112, 153	7 (3%)
4	R	232/241 (96%)	0.76	14 (6%)	27	23	42, 81, 116, 153	3 (1%)
5	E	236/263 (89%)	0.71	11 (4%)	36	31	46, 77, 119, 152	0
5	S	236/263 (89%)	0.57	11 (4%)	36	31	43, 69, 117, 170	2 (0%)
6	F	244/255 (95%)	0.71	8 (3%)	49	42	47, 78, 120, 141	6 (2%)
6	T	241/255 (94%)	0.62	10 (4%)	41	35	44, 71, 105, 131	2 (0%)
7	G	241/246 (97%)	0.77	10 (4%)	41	35	47, 82, 119, 156	4 (1%)
7	U	242/246 (98%)	0.64	11 (4%)	38	32	39, 73, 108, 140	5 (2%)
8	H	218/273 (79%)	0.52	6 (2%)	55	47	42, 65, 109, 130	2 (0%)
8	V	218/273 (79%)	0.54	8 (3%)	45	38	40, 64, 104, 148	3 (1%)
9	I	204/205 (99%)	0.35	2 (0%)	79	74	34, 59, 84, 126	2 (0%)
9	W	204/205 (99%)	0.49	8 (3%)	43	36	39, 60, 83, 125	0
10	J	196/201 (97%)	0.49	4 (2%)	65	58	38, 63, 91, 125	2 (1%)
10	X	197/201 (98%)	0.45	7 (3%)	46	39	40, 63, 89, 121	4 (2%)
11	K	203/276 (73%)	0.37	7 (3%)	48	41	41, 62, 87, 107	0
11	Y	203/276 (73%)	0.48	4 (1%)	65	58	40, 60, 92, 117	0
12	L	213/241 (88%)	0.42	5 (2%)	61	53	35, 58, 91, 118	2 (0%)
12	Z	213/241 (88%)	0.37	3 (1%)	73	67	32, 57, 86, 117	2 (0%)

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
13	M	214/264 (81%)	0.35	1 (0%) 87 83	40, 63, 85, 122	1 (0%)
13	a	214/264 (81%)	0.39	3 (1%) 73 67	38, 61, 87, 106	0
14	N	199/219 (90%)	0.41	3 (1%) 72 65	42, 61, 89, 103	1 (0%)
14	b	199/219 (90%)	0.40	1 (0%) 87 83	33, 60, 84, 106	2 (1%)
All	All	6207/6854 (90%)	0.57	204 (3%) 49 42	32, 68, 110, 176	89 (1%)

The worst 5 of 204 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
2	B	194	ILE	4.6
12	L	33	PHE	4.5
7	G	159	TYR	4.1
7	G	40	VAL	4.1
5	E	236	LEU	3.9

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.


Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
16	NA	I	301	1/1	0.79	0.14	66,66,66,66	0
16	NA	Z	302	1/1	0.79	0.13	62,62,62,62	0
17	HUJ	K	901	33/34	0.80	0.14	53,77,118,131	2
16	NA	N	902	1/1	0.82	0.11	49,49,49,49	0
17	HUJ	Y	901	33/34	0.82	0.17	56,93,126,129	2
17	HUJ	b	901	33/34	0.83	0.18	40,66,132,147	0
16	NA	L	303	1/1	0.84	0.13	68,68,68,68	0

Continued on next page...

Continued from previous page...




Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
17	HUJ	N	901	33/34	0.86	0.12	42,63,82,83	11
15	SCN	Z	301	3/3	0.87	0.19	65,65,72,81	0
15	SCN	L	301	3/3	0.88	0.12	46,46,64,71	0
16	NA	I	302	1/1	0.89	0.08	58,58,58,58	0
16	NA	L	302	1/1	0.91	0.10	68,68,68,68	0
15	SCN	E	301	3/3	0.91	0.11	67,67,68,74	0
16	NA	G	301	1/1	0.92	0.08	58,58,58,58	0
16	NA	U	301	1/1	0.93	0.08	60,60,60,60	0
16	NA	H	301	1/1	0.95	0.05	62,62,62,62	0
16	NA	b	902	1/1	0.95	0.06	61,61,61,61	0
16	NA	W	301	1/1	0.95	0.07	66,66,66,66	0
16	NA	K	902	1/1	0.98	0.04	48,48,48,48	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.







Electron density around HUJ Y 901:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)

**Electron density around HUJ b 901:**

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)





6.5 Other polymers [i](#)

There are no such residues in this entry.