



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 10, 2026 – 10:03 AM UTC

PDB ID : 6E6J / pdb_00006e6j
Title : BRD2_Bromodomain2 complex with inhibitor 744
Authors : Longenecker, K.L.; Park, C.H.; Bigelow, L.
Deposited on : 2018-07-25
Resolution : 2.44 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Buster-report	:	wwPDB partial adaption of 1.1.7 (2018)
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance ⓘ

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.44 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	2340 (2.46-2.42)
Clashscore	190562	2400 (2.46-2.42)
Ramachandran outliers	187476	2379 (2.46-2.42)
RSRZ outliers	180081	2340 (2.46-2.42)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	112	93% . . .
1	B	112	90% 7% .
1	C	112	92% 5% .
1	D	112	93% . .
1	E	112	96% . .
1	F	112	94% . .

2 Entry composition

There are 3 unique types of molecules in this entry. The entry contains 5950 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Bromodomain-containing protein 2.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	109	Total	C	N	O	S	0	0	0
			906	580	160	158	8			
1	B	109	Total	C	N	O	S	0	0	0
			906	580	160	158	8			
1	C	109	Total	C	N	O	S	0	0	0
			906	580	160	158	8			
1	D	109	Total	C	N	O	S	0	0	0
			906	580	160	158	8			
1	E	109	Total	C	N	O	S	0	0	0
			906	580	160	158	8			
1	F	109	Total	C	N	O	S	0	0	0
			906	580	160	158	8			

There are 24 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	344	GLY	-	expression tag	UNP P25440
A	345	SER	-	expression tag	UNP P25440
A	346	HIS	-	expression tag	UNP P25440
A	347	MET	-	expression tag	UNP P25440
B	344	GLY	-	expression tag	UNP P25440
B	345	SER	-	expression tag	UNP P25440
B	346	HIS	-	expression tag	UNP P25440
B	347	MET	-	expression tag	UNP P25440
C	344	GLY	-	expression tag	UNP P25440
C	345	SER	-	expression tag	UNP P25440
C	346	HIS	-	expression tag	UNP P25440
C	347	MET	-	expression tag	UNP P25440
D	344	GLY	-	expression tag	UNP P25440
D	345	SER	-	expression tag	UNP P25440
D	346	HIS	-	expression tag	UNP P25440
D	347	MET	-	expression tag	UNP P25440
E	344	GLY	-	expression tag	UNP P25440

Continued on next page...

Chain	Residue	Modelled	Actual	Comment	Reference
E	345	SER	-	expression tag	UNP P25440
E	346	HIS	-	expression tag	UNP P25440
E	347	MET	-	expression tag	UNP P25440
F	344	GLY	-	expression tag	UNP P25440
F	345	SER	-	expression tag	UNP P25440
F	346	HIS	-	expression tag	UNP P25440
F	347	MET	-	expression tag	UNP P25440

- # HWV
-
- The chemical structure is a complex molecule with the following features:
- Central Pyrazole Ring:** A five-membered ring with two nitrogen atoms (N1, N2) and three carbon atoms (C3, C4, C5). It is connected to a pyridine ring at C3 and a benzene ring at C5.
 - Pyridine Ring:** A six-membered aromatic ring with one nitrogen atom (N1) and five carbon atoms (C6, C7, C8, C9, C10). It is connected to the central pyrazole ring at C3 and a benzene ring at C5.
 - Benzene Ring (Left):** A six-membered aromatic ring with one fluorine atom (F) and five carbon atoms (C11, C12, C13, C14, C15). It is connected to the central pyrazole ring at C5 and a benzene ring at C11.
 - Benzene Ring (Right):** A six-membered aromatic ring with one fluorine atom (F) and five carbon atoms (C16, C17, C18, C19, C20). It is connected to the central pyrazole ring at C11 and a benzene ring at C16.
 - Hydroxyl Group:** A hydroxyl group (OH) is attached to the pyrazole ring at C4, with the oxygen atom labeled O3.
 - Methyl Group:** A methyl group (CH3) is attached to the pyridine ring at C8, with the carbon atom labeled C1.
 - Other Labels:** Various carbon atoms are labeled with C1 through C27, and nitrogen atoms are labeled with N1 and N2. Oxygen atoms are labeled with O1, O2, and O3.

- Molecule 3 is water.


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	79	Total 79	O 79	0	0
3	B	54	Total 54	O 54	0	0
3	C	63	Total 63	O 63	0	0
3	D	75	Total 75	O 75	0	0
3	E	11	Total 11	O 11	0	0
3	F	16	Total 16	O 16	0	0

3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Bromodomain-containing protein 2

Chain A:  93% . . .



- Molecule 1: Bromodomain-containing protein 2

Chain B:  90% 7% .





- Molecule 1: Bromodomain-containing protein 2

Chain C:  92% 5% .





- Molecule 1: Bromodomain-containing protein 2

Chain D:  93% . .




- Molecule 1: Bromodomain-containing protein 2

Chain E:  96% . .



- Molecule 1: Bromodomain-containing protein 2

Chain F:  94% . .



4 Data and refinement statistics

Property	Value	Source
Space group	P 41	Depositor
Cell constants a, b, c, α , β , γ	107.78Å 107.78Å 89.75Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	34.48 – 2.44 34.48 – 2.44	Depositor EDS
% Data completeness (in resolution range)	100.0 (34.48-2.44) 100.0 (34.48-2.44)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.66 (at 2.42Å)	Xtriage
Refinement program	BUSTER 2.11.7	Depositor
R, R_{free}	0.211 , 0.238 (Not available) , 0.242	Depositor DCC
R_{free} test set	1908 reflections (4.97%)	wwPDB-VP
Wilson B-factor (Å ²)	47.2	Xtriage
Anisotropy	0.242	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.35 , 52.5	EDS
L-test for twinning ²	$\langle L \rangle = 0.46$, $\langle L^2 \rangle = 0.28$	Xtriage
Estimated twinning fraction	0.418 for h,-k,-l	Xtriage
F_o, F_c correlation	0.96	EDS
Total number of atoms	5950	wwPDB-VP
Average B, all atoms (Å ²)	62.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 8.06% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: HWV

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.90	0/932	1.41	4/1255 (0.3%)
1	B	0.86	0/932	1.39	0/1255
1	C	0.88	0/932	1.41	2/1255 (0.2%)
1	D	0.86	0/932	1.39	0/1255
1	E	0.82	0/932	1.42	2/1255 (0.2%)
1	F	0.81	0/932	1.45	0/1255
All	All	0.85	0/5592	1.41	8/7530 (0.1%)

There are no bond length outliers.

All (8) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	C	362	SER	CA-C-N	5.57	128.01	120.38
1	C	362	SER	C-N-CA	5.57	128.01	120.38
1	A	411	ALA	CA-C-N	5.48	127.62	120.28
1	A	411	ALA	C-N-CA	5.48	127.62	120.28
1	A	410	ASP	CA-C-N	5.02	127.51	120.28
1	A	410	ASP	C-N-CA	5.02	127.51	120.28
1	E	411	ALA	CA-C-N	5.02	127.42	120.29
1	E	411	ALA	C-N-CA	5.02	127.42	120.29

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen

atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	906	0	885	4	0
1	B	906	0	885	6	0
1	C	906	0	885	4	0
1	D	906	0	885	4	0
1	E	906	0	885	0	0
1	F	906	0	885	2	0
2	A	36	0	0	0	0
2	B	36	0	0	0	0
2	C	36	0	0	0	0
2	D	36	0	0	0	0
2	E	36	0	0	0	0
2	F	36	0	0	0	0
3	A	79	0	0	0	0
3	B	54	0	0	0	0
3	C	63	0	0	0	0
3	D	75	0	0	0	0
3	E	11	0	0	0	0
3	F	16	0	0	0	0
All	All	5950	0	5310	20	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 2.

All (20) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:359:GLU:HG2	1:B:449:ARG:HD3	1.88	0.55
1:F:349:GLN:HE21	1:F:411:ALA:H	1.54	0.54
1:B:346:HIS:HB3	1:B:349:GLN:HG2	1.93	0.51
1:C:346:HIS:HB3	1:C:349:GLN:HG2	1.95	0.49
1:A:363:LYS:H	1:A:363:LYS:HE3	1.78	0.48
1:D:346:HIS:HB3	1:D:349:GLN:HG2	1.95	0.47
1:D:349:GLN:HE21	1:D:410:ASP:HB2	1.80	0.47
1:A:349:GLN:HE21	1:A:410:ASP:HB2	1.80	0.46
1:A:349:GLN:NE2	1:A:410:ASP:HB2	2.31	0.46
1:B:346:HIS:HD2	1:B:347:MET:HE3	1.81	0.46
1:B:349:GLN:HE21	1:B:410:ASP:HB2	1.81	0.45
1:C:349:GLN:HE21	1:C:410:ASP:HB2	1.82	0.44
1:B:401:ARG:O	1:B:405:ASN:HB2	2.18	0.44

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:349:GLN:NE2	1:B:410:ASP:HB2	2.33	0.44
1:C:349:GLN:NE2	1:C:410:ASP:HB2	2.34	0.43
1:D:453:MET:HA	1:D:454:PRO:HD3	1.92	0.42
1:D:349:GLN:NE2	1:D:410:ASP:HB2	2.34	0.42
1:A:346:HIS:HB3	1:A:349:GLN:HG2	2.03	0.41
1:C:359:GLU:O	1:C:365:HIS:HD2	2.04	0.41
1:F:354:ASN:O	1:F:358:LYS:HG2	2.20	0.41

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	107/112 (96%)	104 (97%)	3 (3%)	0	100	100
1	B	107/112 (96%)	106 (99%)	1 (1%)	0	100	100
1	C	107/112 (96%)	104 (97%)	3 (3%)	0	100	100
1	D	107/112 (96%)	104 (97%)	3 (3%)	0	100	100
1	E	107/112 (96%)	107 (100%)	0	0	100	100
1	F	107/112 (96%)	105 (98%)	2 (2%)	0	100	100
All	All	642/672 (96%)	630 (98%)	12 (2%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [i](#)

There are no protein residues with a non-rotameric sidechain to report in this entry.

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no oligosaccharides in this entry.

5.6 Ligand geometry ⓘ

6 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
2	HWV	A	501	-	39,39,39	0.35	0	55,59,59	0.73	2 (3%)
2	HWV	F	501	-	39,39,39	0.40	0	55,59,59	0.73	3 (5%)
2	HWV	C	501	-	39,39,39	0.41	0	55,59,59	0.77	3 (5%)
2	HWV	E	501	-	39,39,39	0.33	0	55,59,59	0.68	2 (3%)
2	HWV	D	501	-	39,39,39	0.37	0	55,59,59	0.75	3 (5%)
2	HWV	B	501	-	39,39,39	0.39	0	55,59,59	0.77	3 (5%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	HWV	A	501	-	-	0/21/21/21	0/4/4/4
2	HWV	F	501	-	-	0/21/21/21	0/4/4/4
2	HWV	C	501	-	-	0/21/21/21	0/4/4/4

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	HWV	E	501	-	-	0/21/21/21	0/4/4/4
2	HWV	D	501	-	-	0/21/21/21	0/4/4/4
2	HWV	B	501	-	-	0/21/21/21	0/4/4/4

There are no bond length outliers.

All (16) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	C	501	HWV	C4-C3-C2	-2.82	122.40	132.80
2	F	501	HWV	C16-O2-C17	2.75	123.85	117.94
2	B	501	HWV	C4-C3-C2	-2.65	123.03	132.80
2	E	501	HWV	C16-O2-C17	2.62	123.58	117.94
2	B	501	HWV	C16-O2-C17	2.51	123.33	117.94
2	C	501	HWV	C16-O2-C17	2.49	123.29	117.94
2	C	501	HWV	C2-C3-N2	2.38	129.33	120.53
2	D	501	HWV	C16-O2-C17	2.37	123.03	117.94
2	E	501	HWV	C6-C5-C10	-2.35	116.16	118.69
2	D	501	HWV	C6-C5-C10	-2.25	116.27	118.69
2	A	501	HWV	C6-C5-C10	-2.24	116.27	118.69
2	F	501	HWV	C6-C5-C10	-2.21	116.31	118.69
2	B	501	HWV	C2-C3-N2	2.18	128.60	120.53
2	D	501	HWV	C4-C3-C2	-2.13	124.93	132.80
2	F	501	HWV	C4-C3-C2	-2.04	125.28	132.80
2	A	501	HWV	C4-C3-C2	-2.03	125.30	132.80

There are no chirality outliers.


There are no torsion outliers.


There are no ring outliers.


No monomer is involved in short contacts.


The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient


equivalents in the CSD to analyse the geometry.














5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	109/112 (97%)	-1.14	0 100 100	27, 43, 67, 87	0
1	B	109/112 (97%)	-1.07	0 100 100	31, 47, 73, 96	0
1	C	109/112 (97%)	-1.06	0 100 100	32, 46, 75, 93	0
1	D	109/112 (97%)	-1.15	0 100 100	30, 44, 68, 88	0
1	E	109/112 (97%)	-0.60	0 100 100	64, 85, 107, 116	0
1	F	109/112 (97%)	-0.41	0 100 100	64, 94, 124, 134	0
All	All	654/672 (97%)	-0.91	0 100 100	27, 54, 111, 134	0

There are no RSRZ outliers to report.

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
2	HWV	A	501	36/36	0.99	0.05	32,45,57,60	0

Continued on next page...




Continued from previous page...

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
2	HWV	B	501	36/36	0.99	0.03	32,39,43,44	0
2	HWV	C	501	36/36	0.99	0.04	30,42,47,48	0
2	HWV	D	501	36/36	0.99	0.05	35,47,62,64	0
2	HWV	E	501	36/36	0.99	0.07	92,98,102,103	0
2	HWV	F	501	36/36	0.99	0.05	68,73,87,89	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.




Electron density around HWV A 501:

2mF_o-DF_c (at 0.7 rmsd) in gray
mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around HWV B 501:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around HWV C 501:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around HWV D 501:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around HWV E 501:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



Electron density around HWV F 501:

2mF_o-DF_c (at 0.7 rmsd) in gray
mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



6.5 Other polymers [i](#)

There are no such residues in this entry.