



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 8, 2026 – 03:39 AM UTC

PDB ID : 8E7Y / pdb_00008e7y
Title : RsTSPO A138F with two heme bound
Authors : Liu, J.; Hiser, C.; Li, F.; Garavito, R.; Ferguson-Miller, S.
Deposited on : 2022-08-25
Resolution : 2.30 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0
Mogul : 2022.3.0, CSD as543be (2022)
Xtriage (Phenix) : 2.0
EDS : 3.0
Buster-report : wwPDB partial adaption of 1.1.7 (2018)
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4 : 9.0.010 (Gargrove)
Density-Fitness : 1.0.12
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.30 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	6319 (2.30-2.30)
Clashscore	190562	6919 (2.30-2.30)
Ramachandran outliers	187476	6854 (2.30-2.30)
Sidechain outliers	187428	6854 (2.30-2.30)
RSRZ outliers	180081	6325 (2.30-2.30)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	158	8% 88% 11%
1	B	158	8% 92% 8%

2 Entry composition [i](#)

There are 4 unique types of molecules in this entry. The entry contains 2716 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called Tryptophan-rich sensory protein.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	B	157	Total	C	N	O	S	0	0	0
			1258	848	198	201	11			
1	A	157	Total	C	N	O	S	0	0	0
			1254	848	196	199	11			

There are 2 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
B	138	PHE	ALA	engineered mutation	UNP Q9RFC8
A	138	PHE	ALA	engineered mutation	UNP Q9RFC8

- Molecule 2 is (2R)-2,3-dihydroxypropyl (9Z)-octadec-9-enoate (CCD ID: OLC) (formula: C₂₁H₄₀O₄) (labeled as "Ligand of Interest" by depositor).




Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
2	B	1	Total	C	O	0	0
			25	21	4		

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
2	B	1	Total C 16 16	0	0
2	A	1	Total C 18 18	0	0
2	A	1	Total C 9 9	0	0
2	A	1	Total C 17 17	0	0
2	A	1	Total C O 23 20 3	0	0

- Molecule 3 is PROTOPORPHYRIN IX CONTAINING FE (CCD ID: HEM) (formula: $C_{34}H_{32}FeN_4O_4$) (labeled as "Ligand of Interest" by depositor).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	B	1	Total C Fe N O 43 34 1 4 4	0	0
3	A	1	Total C Fe N O 43 34 1 4 4	0	0


- Molecule 4 is water.


Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
4	B	6	Total O 6 6	0	0
4	A	4	Total O 4 4	0	0

3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Tryptophan-rich sensory protein

Chain B: 



- Molecule 1: Tryptophan-rich sensory protein

Chain A: 



4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	40.11Å 95.35Å 105.55Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	46.17 – 2.30 46.17 – 2.30	Depositor EDS
% Data completeness (in resolution range)	99.9 (46.17-2.30) 94.3 (46.17-2.30)	Depositor EDS
R_{merge}	0.12	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.51 (at 2.29Å)	Xtriage
Refinement program	PHENIX 1.17.1_3660	Depositor
R, R_{free}	0.216 , 0.251 0.220 , 0.250	Depositor DCC
R_{free} test set	561 reflections (3.00%)	wwPDB-VP
Wilson B-factor (Å ²)	49.2	Xtriage
Anisotropy	0.171	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.29 , 30.3	EDS
L-test for twinning ²	$\langle L \rangle = 0.50$, $\langle L^2 \rangle = 0.34$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.94	EDS
Total number of atoms	2716	wwPDB-VP
Average B, all atoms (Å ²)	55.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 10.03% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: HEM, OLC

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.17	0/1304	0.34	0/1791
1	B	0.14	0/1308	0.31	0/1797
All	All	0.15	0/2612	0.33	0/3588

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1254	0	1219	16	0
1	B	1258	0	1219	11	0
2	A	67	0	116	9	0
2	B	41	0	69	2	0
3	A	43	0	30	4	0
3	B	43	0	30	4	0
4	A	4	0	0	0	0
4	B	6	0	0	0	0
All	All	2716	0	2683	35	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 6.

All (35) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:27:PRO:HD2	3:B:203:HEM:HBD2	1.78	0.63
1:A:27:PRO:HD2	3:A:205:HEM:HBD2	1.81	0.62
1:B:104:VAL:HG23	3:B:203:HEM:HBB1	1.83	0.59
3:A:205:HEM:HMC2	3:A:205:HEM:HBC2	1.84	0.59
1:B:97:MET:HE1	1:B:150:ASN:HB2	1.84	0.58
3:B:203:HEM:HBC2	3:B:203:HEM:HMC1	1.87	0.56
1:B:102:ALA:HA	1:B:105:MET:HE2	1.92	0.52
1:A:118:PHE:CE1	1:A:129:PHE:CE2	2.98	0.52
2:B:201:OLC:H6A	2:A:201:OLC:H13A	1.96	0.48
1:A:9:LEU:HB3	2:A:203:OLC:H4	1.96	0.48
1:A:91:PHE:O	1:A:95:LYS:HA	2.15	0.47
2:B:202:OLC:H8A	2:B:202:OLC:H11A	1.70	0.46
1:B:94:MET:HE2	1:A:25:LEU:HD11	1.98	0.45
1:A:104:VAL:HG23	3:A:205:HEM:HBB1	1.97	0.45
1:A:124:TRP:O	1:A:127:VAL:HG12	2.16	0.45
1:B:89:PRO:HB3	1:A:89:PRO:HB3	1.97	0.45
1:B:27:PRO:HG2	3:B:203:HEM:HMD3	1.97	0.45
1:A:1:MET:HE1	1:A:122:ASP:OD2	2.17	0.45
2:A:203:OLC:H12A	2:A:203:OLC:H15A	1.36	0.45
1:A:1:MET:HE2	1:A:1:MET:HB2	1.68	0.45
1:B:39:TRP:CZ2	1:B:145:GLU:HG3	2.52	0.44
1:A:1:MET:HE3	1:A:63:ARG:NH2	2.32	0.44
1:B:39:TRP:HZ2	1:B:145:GLU:HG3	1.82	0.44
1:A:1:MET:O	1:A:2:ASN:ND2	2.51	0.43
2:A:203:OLC:H10	2:A:203:OLC:H13	1.59	0.43
1:A:143:ASN:C	1:A:143:ASN:HD22	2.27	0.43
1:A:6:ALA:HB1	2:A:203:OLC:H11A	1.99	0.43
3:A:205:HEM:HBB2	3:A:205:HEM:HMB2	2.01	0.42
1:A:124:TRP:HA	1:A:127:VAL:HG12	2.01	0.41
2:A:202:OLC:H17	2:A:202:OLC:H14	1.85	0.41
2:A:201:OLC:H11	2:A:201:OLC:H8	1.56	0.41
1:B:143:ASN:HD22	1:B:143:ASN:C	2.29	0.41
2:A:203:OLC:H8	2:A:203:OLC:H5A	1.75	0.41
1:A:60:ALA:HB3	1:A:129:PHE:HE2	1.87	0.40
1:B:110:PHE:CZ	2:A:204:OLC:H14A	2.57	0.40

There are no symmetry-related clashes.

5.3 Torsion angles

5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	155/158 (98%)	151 (97%)	4 (3%)	0	100	100
1	B	155/158 (98%)	151 (97%)	4 (3%)	0	100	100
All	All	310/316 (98%)	302 (97%)	8 (3%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	120/124 (97%)	120 (100%)	0	100	100
1	B	121/124 (98%)	121 (100%)	0	100	100
All	All	241/248 (97%)	241 (100%)	0	100	100

There are no protein residues with a non-rotameric sidechain to report.

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (3) such sidechains are listed below:

Mol	Chain	Res	Type
1	B	66	GLN
1	B	143	ASN
1	A	2	ASN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no oligosaccharides in this entry.

5.6 Ligand geometry ⓘ

8 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# $ Z > 2$	Counts	RMSZ	# $ Z > 2$
3	HEM	A	205	-	50,50,50	1.39	9 (18%)	67,82,82	1.12	4 (5%)
2	OLC	A	204	-	22,22,24	0.71	1 (4%)	22,22,25	0.92	1 (4%)
2	OLC	B	201	-	24,24,24	0.68	1 (4%)	25,25,25	0.94	1 (4%)
3	HEM	B	203	-	50,50,50	1.41	9 (18%)	67,82,82	1.09	3 (4%)
2	OLC	B	202	-	15,15,24	0.21	0	14,14,25	0.80	0
2	OLC	A	203	-	16,16,24	0.21	0	15,15,25	0.68	0
2	OLC	A	201	-	17,17,24	0.20	0	16,16,25	0.76	0
2	OLC	A	202	-	8,8,24	0.10	0	7,7,25	0.75	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	HEM	A	205	-	-	4/14/54/54	-

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	OLC	A	204	-	-	9/21/21/24	-
2	OLC	B	201	-	-	13/24/24/24	-
3	HEM	B	203	-	-	4/14/54/54	-
2	OLC	B	202	-	-	4/13/13/24	-
2	OLC	A	203	-	-	8/14/14/24	-
2	OLC	A	201	-	-	6/15/15/24	-
2	OLC	A	202	-	-	1/6/6/24	-

All (20) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	B	203	HEM	FE-NB	3.72	2.06	1.94
3	A	205	HEM	FE-NB	3.31	2.05	1.94
3	A	205	HEM	CAC-C3C	3.15	1.55	1.47
3	A	205	HEM	FE-ND	3.14	2.04	1.94
3	B	203	HEM	CAC-C3C	3.12	1.55	1.47
3	A	205	HEM	FE-NA	3.04	2.05	1.95
3	B	203	HEM	CAB-C3B	2.92	1.55	1.47
3	A	205	HEM	CAB-C3B	2.91	1.55	1.47
3	B	203	HEM	FE-NA	2.90	2.04	1.95
3	B	203	HEM	FE-ND	2.82	2.03	1.94
2	A	204	OLC	O20-C1	2.82	1.41	1.33
2	B	201	OLC	O20-C1	2.64	1.41	1.33
3	B	203	HEM	FE-NC	2.61	2.03	1.95
3	A	205	HEM	FE-NC	2.48	2.03	1.95
3	B	203	HEM	CMD-C2D	2.25	1.55	1.50
3	A	205	HEM	CMD-C2D	2.21	1.55	1.50
3	B	203	HEM	CMB-C2B	2.19	1.55	1.50
3	B	203	HEM	CMC-C2C	2.18	1.55	1.50
3	A	205	HEM	CMC-C2C	2.14	1.55	1.50
3	A	205	HEM	CMB-C2B	2.12	1.55	1.50

All (9) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	B	201	OLC	O20-C1-C2	2.74	120.18	111.83
3	A	205	HEM	C1B-NB-C4B	2.57	108.25	105.21
2	A	204	OLC	O20-C1-C2	2.50	119.45	111.83
3	A	205	HEM	C4D-ND-C1D	2.43	108.08	105.21
3	B	203	HEM	C4D-ND-C1D	2.39	108.04	105.21
3	B	203	HEM	C1B-NB-C4B	2.39	108.03	105.21

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	A	205	HEM	C2A-C1A-NA	-2.12	107.80	110.15
3	B	203	HEM	C2A-C1A-NA	-2.07	107.85	110.15
3	A	205	HEM	C3B-C4B-NB	-2.03	108.01	109.47

There are no chirality outliers.

All (49) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
3	B	203	HEM	C2D-C3D-CAD-CBD
3	B	203	HEM	C4D-C3D-CAD-CBD
3	A	205	HEM	C2D-C3D-CAD-CBD
3	A	205	HEM	C4D-C3D-CAD-CBD
2	B	201	OLC	O19-C1-O20-C21
2	B	201	OLC	C2-C1-O20-C21
2	A	203	OLC	C12-C13-C14-C15
2	B	201	OLC	O20-C21-C22-O23
2	B	201	OLC	C2-C3-C4-C5
2	B	201	OLC	C5-C6-C7-C8
2	A	201	OLC	C11-C12-C13-C14
2	A	204	OLC	O20-C21-C22-O23
2	B	201	OLC	C12-C13-C14-C15
2	A	201	OLC	C5-C6-C7-C8
2	A	204	OLC	C11-C12-C13-C14
2	A	203	OLC	C13-C14-C15-C16
2	B	201	OLC	C13-C14-C15-C16
2	A	203	OLC	C14-C15-C16-C17
2	A	204	OLC	C5-C6-C7-C8
2	A	202	OLC	C12-C13-C14-C15
2	B	201	OLC	C6-C7-C8-C9
2	A	204	OLC	C6-C7-C8-C9
2	A	203	OLC	C11-C12-C13-C14
2	A	201	OLC	C4-C5-C6-C7
2	A	204	OLC	C4-C5-C6-C7
2	A	203	OLC	C2-C3-C4-C5
2	B	202	OLC	C12-C13-C14-C15
2	A	201	OLC	C13-C14-C15-C16
2	A	203	OLC	C4-C5-C6-C7
2	B	201	OLC	C4-C5-C6-C7
2	B	202	OLC	C11-C12-C13-C14
2	A	204	OLC	C14-C15-C16-C17
2	A	201	OLC	C6-C7-C8-C9
2	A	204	OLC	C13-C14-C15-C16

Continued on next page...

Continued from previous page...



Mol	Chain	Res	Type	Atoms
2	B	202	OLC	C4-C5-C6-C7
3	B	203	HEM	CAA-CBA-CGA-O1A
3	A	205	HEM	CAA-CBA-CGA-O1A
3	A	205	HEM	CAA-CBA-CGA-O2A
3	B	203	HEM	CAA-CBA-CGA-O2A
2	A	203	OLC	C10-C11-C12-C13
2	B	201	OLC	O20-C21-C22-C24
2	B	201	OLC	C9-C10-C11-C12
2	A	203	OLC	C9-C10-C11-C12
2	B	201	OLC	C14-C15-C16-C17
2	A	204	OLC	C2-C3-C4-C5
2	A	204	OLC	C9-C10-C11-C12
2	B	201	OLC	C10-C11-C12-C13
2	B	202	OLC	C6-C7-C8-C9
2	A	201	OLC	C2-C3-C4-C5

There are no ring outliers.


8 monomers are involved in 18 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	A	205	HEM	4	0
2	A	204	OLC	1	0
2	B	201	OLC	1	0
3	B	203	HEM	4	0
2	B	202	OLC	1	0
2	A	203	OLC	5	0
2	A	201	OLC	2	0
2	A	202	OLC	1	0


The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.







Ligand OLC B 201



Ligand HEM B 203







5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	157/158 (99%)	0.57	12 (7%) 20 21	43, 53, 66, 90	0
1	B	157/158 (99%)	0.71	13 (8%) 17 19	42, 52, 64, 95	0
All	All	314/316 (99%)	0.64	25 (7%) 18 20	42, 52, 65, 95	0

All (25) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	1	MET	8.8
1	A	1	MET	4.1
1	B	157	ARG	3.9
1	B	2	ASN	3.6
1	A	81	LEU	3.6
1	B	156	ALA	3.5
1	A	132	TYR	3.4
1	B	28	ASP	3.2
1	A	129	PHE	2.8
1	B	27	PRO	2.6
1	B	3	MET	2.6
1	A	78	ALA	2.5
1	A	77	TYR	2.4
1	B	67	LEU	2.4
1	B	55	PHE	2.3
1	A	152	ASN	2.3
1	B	134	ILE	2.2
1	A	80	GLN	2.2
1	A	2	ASN	2.2
1	A	55	PHE	2.1
1	B	68	GLU	2.1
1	A	68	GLU	2.1
1	B	51	THR	2.1
1	B	155	GLU	2.1

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	23	ALA	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)




In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
2	OLC	A	204	23/25	0.74	0.25	61,74,80,87	0
2	OLC	A	201	18/25	0.75	0.25	56,69,75,77	0
2	OLC	A	203	17/25	0.77	0.23	62,71,80,80	0
2	OLC	B	201	25/25	0.78	0.21	64,71,86,101	0
2	OLC	A	202	9/25	0.80	0.24	47,61,71,75	0
2	OLC	B	202	16/25	0.83	0.21	63,71,77,83	0
3	HEM	B	203	43/43	0.92	0.11	50,60,71,76	2
3	HEM	A	205	43/43	0.96	0.09	44,54,69,76	0




The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.

Electron density around OLC A 204:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)




**Electron density around OLC A 201:**

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around OLC A 203:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)




**Electron density around OLC B 201:**

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around OLC A 202:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)




**Electron density around OLC B 202:**


$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



Electron density around HEM B 203:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)





6.5 Other polymers ⓘ

There are no such residues in this entry.