



Full wwPDB X-ray Structure Validation Report ⓘ

Apr 24, 2026 – 09:50 PM EDT

PDB ID : 1E8D / pdb_00001e8d
Title : MECHANISTIC ASPECTS OF CYANOGENESIS FROM ACTIVE SITE
MUTANT SER80ALA OF HYDROXYNITRILE LYASE FROM MANIHOT
ESCULENTA IN COMPLEX WITH ACETONE CYANOHYDRIN
Authors : Lauble, H.; Miehlich, B.; Foerster, S.; Wajant, H.; Effenberger, F.
Deposited on : 2000-09-19
Resolution : 2.20 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.20 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
Clashscore	190562	6851 (2.20-2.20)
Ramachandran outliers	187476	6768 (2.20-2.20)
Sidechain outliers	187428	6769 (2.20-2.20)
RSRZ outliers	180081	6166 (2.20-2.20)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	262	
1	B	262	

2 Entry composition

There are 3 unique types of molecules in this entry. The entry contains 4473 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called HYDROXYNITRILE LYASE.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	262	Total	C	N	O	S	0	0	0
			2103	1359	349	387	8			
1	B	258	Total	C	N	O	S	0	0	0
			2073	1339	344	382	8			

There are 12 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	-4	PRO	-	cloning artifact	UNP P52705
A	-3	ILE	-	cloning artifact	UNP P52705
A	-2	SER	-	cloning artifact	UNP P52705
A	-1	LYS	-	cloning artifact	UNP P52705
A	1	MET	-	cloning artifact	UNP P52705
A	80	ALA	SER	engineered mutation	UNP P52705
B	-4	PRO	-	cloning artifact	UNP P52705
B	-3	ILE	-	cloning artifact	UNP P52705
B	-2	SER	-	cloning artifact	UNP P52705
B	-1	LYS	-	cloning artifact	UNP P52705
B	1	MET	-	cloning artifact	UNP P52705
B	80	ALA	SER	engineered mutation	UNP P52705

- Molecule 2 is 2-HYDROXY-2-METHYLPROPANENITRILE (CCD ID: CNH) (formula: C₄H₇NO).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
2	A	1	Total	C	N	O	0	0
			6	4	1	1		
2	A	1	Total	C	N	O	0	0
			6	4	1	1		
2	A	1	Total	C	N	O	0	0
			6	4	1	1		
2	B	1	Total	C	N	O	0	0
			6	4	1	1		


- Molecule 3 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	124	Total	O	0	0
			124	124		
3	B	149	Total	O	0	0
			149	149		


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: HYDROXYNITRILE LYASE



• Molecule 1: HYDROXYNITRILE LYASE



4 Data and refinement statistics

Property	Value	Source
Space group	P 41 21 2	Depositor
Cell constants a, b, c, α , β , γ	105.90Å 105.90Å 187.80Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	8.00 – 2.20 8.00 – 2.20	Depositor EDS
% Data completeness (in resolution range)	93.6 (8.00-2.20) 94.2 (8.00-2.20)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	0.07	Depositor
$\langle I/\sigma(I) \rangle$ ¹	7.51 (at 2.20Å)	Xtriage
Refinement program	X-PLOR 3.851	Depositor
R, R_{free}	0.186 , 0.238 0.183 , (Not available)	Depositor DCC
R_{free} test set	No test flags present.	wwPDB-VP
Wilson B-factor (Å ²)	23.5	Xtriage
Anisotropy	0.545	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.39 , 74.0	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.33$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.95	EDS
Total number of atoms	4473	wwPDB-VP
Average B, all atoms (Å ²)	25.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.31% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: CNH

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	1.08	11/2155 (0.5%)	1.70	35/2920 (1.2%)
1	B	1.08	9/2124 (0.4%)	1.67	28/2879 (1.0%)
All	All	1.08	20/4279 (0.5%)	1.69	63/5799 (1.1%)

All (20) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	B	10	HIS	CD2-NE2	-7.27	1.29	1.37
1	A	10	HIS	CD2-NE2	-6.89	1.30	1.37
1	A	112	HIS	CD2-NE2	-6.87	1.30	1.37
1	A	20	HIS	CD2-NE2	-6.77	1.30	1.37
1	B	248	HIS	CD2-NE2	-6.76	1.30	1.37
1	B	112	HIS	CD2-NE2	-6.51	1.30	1.37
1	A	5	HIS	CD2-NE2	-6.49	1.30	1.37
1	A	31	HIS	CD2-NE2	-6.45	1.30	1.37
1	A	103	HIS	CD2-NE2	-6.35	1.30	1.37
1	B	236	HIS	CD2-NE2	-6.25	1.30	1.37
1	B	5	HIS	CD2-NE2	-6.23	1.30	1.37
1	B	20	HIS	CD2-NE2	-5.97	1.31	1.37
1	A	14	HIS	CD2-NE2	-5.83	1.31	1.37
1	B	103	HIS	CD2-NE2	-5.59	1.31	1.37
1	A	236	HIS	CD2-NE2	-5.48	1.31	1.37
1	A	248	HIS	CD2-NE2	-5.32	1.32	1.37
1	B	31	HIS	CD2-NE2	-5.29	1.32	1.37
1	A	143	THR	CA-CB	5.11	1.60	1.53
1	B	14	HIS	CD2-NE2	-5.04	1.32	1.37
1	A	112	HIS	CG-ND1	-5.04	1.32	1.38

All (63) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	138	ASN	OD1-CG-ND2	-7.87	114.73	122.60
1	A	138	ASN	CA-CB-CG	7.87	120.47	112.60
1	B	182	ASN	OD1-CG-ND2	-7.65	114.95	122.60
1	A	109	ASP	CA-CB-CG	7.36	119.95	112.60
1	A	206	ASP	CA-CB-CG	7.20	119.80	112.60
1	A	155	ARG	NE-CZ-NH2	-7.00	112.90	119.20
1	B	118	VAL	N-CA-C	-6.87	104.16	110.82
1	B	47	GLN	OE1-CD-NE2	-6.78	115.82	122.60
1	A	186	GLN	OE1-CD-NE2	-6.63	115.97	122.60
1	A	163	THR	N-CA-CB	-6.57	100.87	110.26
1	B	217	ARG	CA-CB-CG	-6.52	101.06	114.10
1	A	125	PHE	CA-CB-CG	-6.43	107.37	113.80
1	A	50	GLN	OE1-CD-NE2	-6.38	116.22	122.60
1	A	19	TRP	CG-CD2-CE3	6.36	140.26	133.90
1	A	163	THR	CA-CB-OG1	-6.27	100.19	109.60
1	A	130	ASP	CA-CB-CG	6.26	118.86	112.60
1	A	103	HIS	CB-CG-CD2	-6.24	123.09	131.20
1	A	236	HIS	CB-CG-CD2	-6.17	123.17	131.20
1	B	208	ASP	CA-CB-CG	6.13	118.73	112.60
1	A	244	GLU	CB-CG-CD	6.12	123.01	112.60
1	B	157	ASN	CA-CB-CG	-6.09	106.51	112.60
1	A	138	ASN	CB-CG-ND2	6.06	125.49	116.40
1	B	112	HIS	CB-CG-CD2	-6.02	123.38	131.20
1	B	182	ASN	CB-CG-ND2	6.02	125.42	116.40
1	A	31	HIS	CB-CG-CD2	-6.01	123.39	131.20
1	B	236	HIS	CB-CG-CD2	-5.97	123.44	131.20
1	A	14	HIS	CB-CG-CD2	-5.89	123.54	131.20
1	B	31	HIS	CB-CG-CD2	-5.83	123.63	131.20
1	B	155	ARG	NE-CZ-NH2	-5.81	113.97	119.20
1	A	19	TRP	CB-CG-CD1	-5.76	118.25	126.90
1	B	52	ASN	OD1-CG-ND2	-5.76	116.84	122.60
1	A	218	TRP	CG-CD2-CE3	5.71	139.61	133.90
1	B	248	HIS	CB-CG-CD2	-5.69	123.80	131.20
1	A	112	HIS	CB-CG-CD2	-5.59	123.94	131.20
1	A	245	GLU	N-CA-C	-5.59	105.11	111.14
1	B	59	GLU	CA-C-O	-5.52	113.55	118.63
1	B	104	ASN	OD1-CG-ND2	-5.51	117.09	122.60
1	B	109	ASP	CA-CB-CG	5.50	118.10	112.60
1	A	-1	LYS	CB-CG-CD	-5.47	98.71	111.30
1	B	182	ASN	CA-CB-CG	5.47	118.07	112.60
1	B	130	ASP	CA-CB-CG	5.46	118.06	112.60
1	A	192	GLU	O-C-N	5.45	128.36	122.15
1	B	236	HIS	CB-CG-ND1	5.42	130.82	122.70

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	103	HIS	CA-CB-CG	5.38	119.18	113.80
1	B	111	VAL	N-CA-C	5.37	117.81	111.09
1	B	218	TRP	CG-CD2-CE3	5.35	139.25	133.90
1	A	218	TRP	CB-CG-CD1	-5.32	118.92	126.90
1	B	19	TRP	CG-CD2-CE3	5.31	139.21	133.90
1	A	173	VAL	N-CA-C	5.30	118.29	113.20
1	B	3	THR	O-C-N	-5.30	117.04	123.03
1	A	179	LEU	N-CA-C	-5.28	106.34	112.89
1	A	142	GLU	CA-C-O	-5.22	115.31	121.16
1	B	244	GLU	CB-CG-CD	5.16	121.38	112.60
1	A	186	GLN	CG-CD-NE2	5.10	124.05	116.40
1	A	163	THR	CA-CB-CG2	5.09	119.16	110.50
1	A	248	HIS	CB-CG-CD2	-5.07	124.61	131.20
1	A	-4	PRO	N-CA-CB	5.07	108.57	103.00
1	A	100	GLY	O-C-N	-5.06	119.14	123.80
1	A	131	THR	CA-CB-OG1	-5.06	102.01	109.60
1	A	217	ARG	CA-CB-CG	-5.05	104.01	114.10
1	B	103	HIS	CB-CG-CD2	-5.04	124.65	131.20
1	B	119	GLU	CA-CB-CG	-5.03	104.03	114.10
1	B	230	GLN	OE1-CD-NE2	-5.01	117.58	122.60

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2103	0	2101	27	0
1	B	2073	0	2065	23	0
2	A	18	0	21	3	0
2	B	6	0	7	0	0
3	A	124	0	0	1	0
3	B	149	0	0	0	0
All	All	4473	0	4194	49	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including

hydrogen atoms). The all-atom clashscore for this structure is 6.

All (49) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:10:HIS:HE1	1:B:38:MET:H	1.20	0.90
1:A:10:HIS:HE1	1:A:38:MET:H	1.19	0.89
1:A:137:THR:H	2:A:1260:CNH:HC31	1.49	0.76
1:B:52:ASN:HD22	1:B:138:ASN:HB2	1.52	0.74
1:A:81:CYS:HA	1:A:106:LEU:HD22	1.72	0.70
1:A:10:HIS:CE1	1:A:38:MET:H	2.09	0.67
1:A:52:ASN:HD22	1:A:138:ASN:HB2	1.60	0.67
1:B:10:HIS:HD2	1:B:11:THR:O	1.82	0.63
1:A:10:HIS:HD2	1:A:11:THR:O	1.82	0.61
1:B:10:HIS:CE1	1:B:38:MET:H	2.11	0.60
1:A:163:THR:HG22	1:A:166:GLU:H	1.66	0.60
1:B:82:ALA:HA	1:B:85:ASN:HD22	1.66	0.60
1:A:138:ASN:ND2	1:A:142:GLU:H	2.00	0.60
1:A:10:HIS:HE1	1:A:38:MET:N	1.96	0.58
1:A:82:ALA:HA	1:A:85:ASN:HD22	1.71	0.55
1:A:2:VAL:HG11	1:A:258:ALA:HA	1.90	0.53
2:A:1260:CNH:HC33	3:A:2054:HOH:O	2.07	0.53
1:B:5:HIS:CD2	1:B:32:LYS:HG2	2.44	0.51
1:B:81:CYS:HA	1:B:106:LEU:HD22	1.91	0.51
1:A:14:HIS:O	1:A:41:SER:HB3	2.10	0.50
1:B:104:ASN:ND2	1:B:238:LEU:HD22	2.26	0.50
1:B:135:THR:HA	1:B:144:ILE:O	2.12	0.49
1:A:52:ASN:ND2	1:A:139:ILE:H	2.11	0.49
1:A:69:PRO:HB2	1:A:72:GLU:HG2	1.94	0.48
1:A:68:LEU:HD13	1:A:72:GLU:HB2	1.96	0.46
1:B:104:ASN:HD21	1:B:238:LEU:HD22	1.79	0.46
1:A:20:HIS:HE1	1:A:166:GLU:OE1	1.98	0.45
1:A:139:ILE:HG23	1:A:140:THR:HG23	1.99	0.45
1:A:236:HIS:NE2	2:A:1259:CNH:N5	2.58	0.45
1:B:2:VAL:CG2	1:B:255:ASP:HA	2.47	0.45
1:B:2:VAL:HG11	1:B:258:ALA:HA	1.98	0.45
1:B:155:ARG:HD3	1:B:155:ARG:HA	1.61	0.45
1:A:39:ALA:HB3	1:A:57:TYR:HA	2.00	0.44
1:B:47:GLN:O	1:B:50:GLN:HG2	2.17	0.44
1:B:1:MET:HE2	1:B:1:MET:HB3	1.92	0.44
1:B:52:ASN:ND2	1:B:139:ILE:H	2.15	0.43
1:A:20:HIS:HD2	1:B:165:GLY:O	2.01	0.43
1:A:224:LYS:HA	1:A:225:PRO:HD3	1.94	0.42

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:80:ALA:HA	1:A:105:SER:HA	2.01	0.42
1:B:20:HIS:HE1	1:B:166:GLU:OE1	2.02	0.42
1:B:119:GLU:HG3	1:B:181:GLN:HE21	1.84	0.42
1:A:11:THR:HB	1:A:80:ALA:HB3	2.02	0.42
1:A:24:PRO:HD3	1:B:168:GLU:HG2	2.01	0.41
1:A:209:LYS:HA	1:A:209:LYS:HD3	1.92	0.41
1:A:68:LEU:HD11	1:A:74:VAL:HG13	2.03	0.41
1:A:120:LYS:HG3	1:A:218:TRP:CH2	2.56	0.41
1:B:80:ALA:HA	1:B:105:SER:HA	2.03	0.41
1:B:47:GLN:H	1:B:50:GLN:HE21	1.69	0.40
1:B:154:LEU:HD11	1:B:174:MET:HE3	2.04	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	260/262 (99%)	251 (96%)	9 (4%)	0	100	100
1	B	256/262 (98%)	244 (95%)	12 (5%)	0	100	100
All	All	516/524 (98%)	495 (96%)	21 (4%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	226/226 (100%)	212 (94%)	14 (6%)	16	20
1	B	222/226 (98%)	210 (95%)	12 (5%)	20	25
All	All	448/452 (99%)	422 (94%)	26 (6%)	18	22

All (26) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	-2	SER
1	A	70	GLN
1	A	92	ARG
1	A	103	HIS
1	A	106	LEU
1	A	129	ARG
1	A	138	ASN
1	A	143	THR
1	A	154	LEU
1	A	155	ARG
1	A	163	THR
1	A	169	LEU
1	A	182	ASN
1	A	244	GLU
1	B	32	LYS
1	B	68	LEU
1	B	106	LEU
1	B	121	LEU
1	B	140	THR
1	B	147	MET
1	B	154	LEU
1	B	155	ARG
1	B	169	LEU
1	B	182	ASN
1	B	224	LYS
1	B	238	LEU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (17) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	10	HIS
1	A	20	HIS
1	A	52	ASN
1	A	85	ASN

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	A	138	ASN
1	A	182	ASN
1	B	10	HIS
1	B	20	HIS
1	B	50	GLN
1	B	52	ASN
1	B	70	GLN
1	B	85	ASN
1	B	104	ASN
1	B	181	GLN
1	B	182	ASN
1	B	216	GLN
1	B	248	HIS

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

4 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# $ Z > 2$	Counts	RMSZ	# $ Z > 2$
2	CNH	A	1261	-	5,5,5	2.56	2 (40%)	4,7,7	0.35	0

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
2	CNH	A	1259	-	5,5,5	3.14	1 (20%)	4,7,7	1.19	0
2	CNH	B	1259	-	5,5,5	2.64	2 (40%)	4,7,7	0.95	0
2	CNH	A	1260	-	5,5,5	2.49	2 (40%)	4,7,7	0.72	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	CNH	A	1261	-	-	0/0/3/3	-
2	CNH	A	1259	-	-	0/0/3/3	-
2	CNH	B	1259	-	-	0/0/3/3	-
2	CNH	A	1260	-	-	0/0/3/3	-

All (7) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	A	1259	CNH	C1-C4	-6.39	1.40	1.48
2	A	1261	CNH	C1-C4	-5.00	1.42	1.48
2	B	1259	CNH	C1-C4	-4.86	1.42	1.48
2	A	1260	CNH	C1-C4	-4.79	1.42	1.48
2	B	1259	CNH	C4-N5	2.64	1.18	1.14
2	A	1260	CNH	C4-N5	2.45	1.17	1.14
2	A	1261	CNH	C4-N5	2.21	1.17	1.14

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

2 monomers are involved in 3 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	A	1259	CNH	1	0
2	A	1260	CNH	2	0

5.7 Other polymers ⓘ

There are no such residues in this entry.

5.8 Polymer linkage issues ⓘ

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	262/262 (100%)	-0.74	4 (1%) 72 69	14, 23, 43, 62	0
1	B	258/262 (98%)	-0.84	1 (0%) 88 87	15, 22, 38, 74	0
All	All	520/524 (99%)	-0.79	5 (0%) 79 77	14, 22, 41, 74	0

All (5) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	258	ALA	4.8
1	B	258	ALA	3.0
1	A	-4	PRO	2.3
1	A	95	ASP	2.2
1	A	-2	SER	2.2

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
2	CNH	A	1259	6/6	0.64	0.15	30,39,44,54	0
2	CNH	B	1259	6/6	0.71	0.20	43,48,54,59	0
2	CNH	A	1260	6/6	0.89	0.07	17,24,27,29	0
2	CNH	A	1261	6/6	0.97	0.04	20,26,27,27	0

6.5 Other polymers [i](#)

There are no such residues in this entry.