



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 14, 2026 – 10:41 AM UTC

PDB ID : 5E8P / pdb_00005e8p
Title : The structure of the TEIPP associated altered peptide ligand Trh4-p5NLE in complex with H-2D(b)
Authors : Hafstrand, I.; Doorduijn, E.; Duru, A.D.; Buratto, J.; Oliveira, C.C.; Sandalova, T.; van Hall, T.; Achour, A.
Deposited on : 2015-10-14
Resolution : 2.00 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:


MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49

1 Overall quality at a glance i

The following experimental techniques were used to determine the structure:
X-RAY DIFFRACTION

The reported resolution of this entry is 2.00 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	10052 (2.00-2.00)
Clashscore	190562	11152 (2.00-2.00)
Ramachandran outliers	187476	11031 (2.00-2.00)
Sidechain outliers	187428	11029 (2.00-2.00)
RSRZ outliers	180081	10067 (2.00-2.00)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	276	15% 87% 11% ..
1	D	276	13% 84% 13% .
2	B	99	4% 86% 14%
2	E	99	3% 93% 7%
3	C	9	89% 11%

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
3	F	9	 89% 11%

2 Entry composition

There are 5 unique types of molecules in this entry. The entry contains 6507 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called H-2 class I histocompatibility antigen, D-B alpha chain.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	272	Total	C	N	O	S	0	1	0
			2235	1410	395	421	9			
1	D	271	Total	C	N	O	S	0	1	0
			2222	1405	388	419	10			

- Molecule 2 is a protein called Beta-2-microglobulin.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	B	99	Total	C	N	O	S	0	0	0
			821	524	138	152	7			
2	E	99	Total	C	N	O	S	0	2	0
			833	532	139	154	8			

- Molecule 3 is a protein called Ceramide synthase 5.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
3	C	9	Total	C	N	O	S	0	1	0
			77	48	15	11	3			
3	F	9	Total	C	N	O	S	0	0	0
			69	43	12	11	3			

There are 2 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
C	5	NLE	MET	engineered mutation	UNP Q9D6K9
F	5	NLE	MET	engineered mutation	UNP Q9D6K9

- Molecule 4 is SULFATE ION (CCD ID: SO4) (formula: O₄S).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
4	B	1	Total	O	S	0	0
			5	4	1		


- Molecule 5 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	67	Total	O	0	0
			67	67		
5	B	46	Total	O	0	0
			46	46		
5	C	3	Total	O	0	0
			3	3		
5	D	78	Total	O	0	0
			78	78		
5	E	42	Total	O	0	0
			42	42		
5	F	9	Total	O	0	0
			9	9		


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: H-2 class I histocompatibility antigen, D-B alpha chain




- Molecule 1: H-2 class I histocompatibility antigen, D-B alpha chain



- Molecule 2: Beta-2-microglobulin



- Molecule 2: Beta-2-microglobulin



- Molecule 3: Ceramide synthase 5

Chain C:  89% 11%



● Molecule 3: Ceramide synthase 5

Chain F:  89% 11%



4 Data and refinement statistics

Property	Value	Source
Space group	I 1 2 1	Depositor
Cell constants a, b, c, α , β , γ	90.59Å 123.77Å 97.68Å 90.00° 104.31° 90.00°	Depositor
Resolution (Å)	31.80 – 2.00 31.80 – 2.00	Depositor EDS
% Data completeness (in resolution range)	97.2 (31.80-2.00) 97.4 (31.80-2.00)	Depositor EDS
R_{merge}	0.08	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.79 (at 2.00Å)	Xtriage
Refinement program	PHENIX (1.10_2155: ???)	Depositor
R, R_{free}	0.214 , 0.258 0.215 , 0.255	Depositor DCC
R_{free} test set	3404 reflections (4.84%)	wwPDB-VP
Wilson B-factor (Å ²)	20.1	Xtriage
Anisotropy	0.633	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.33 , 34.4	EDS
L-test for twinning ²	$\langle L \rangle = 0.47$, $\langle L^2 \rangle = 0.30$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.92	EDS
Total number of atoms	6507	wwPDB-VP
Average B, all atoms (Å ²)	32.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 5.23% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: NLE, SO4

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.35	1/2304 (0.0%)	0.61	3/3127 (0.1%)
1	D	0.38	0/2291	0.60	1/3109 (0.0%)
2	B	0.38	0/847	0.58	0/1148
2	E	0.37	0/859	0.60	0/1165
3	C	1.21	0/69	0.71	0/85
3	F	0.44	0/59	0.57	0/74
All	All	0.39	1/6429 (0.0%)	0.60	4/8708 (0.0%)

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	220	ASN	CA-C	-5.23	1.48	1.53

All (4) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	D	196	LYS	N-CA-C	6.00	123.57	110.80
1	A	222	GLU	N-CA-C	5.86	119.47	107.69
1	A	221	GLY	N-CA-C	-5.44	100.28	113.18
1	A	220	ASN	N-CA-C	-5.28	100.31	110.56

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within

the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2235	0	2092	23	0
1	D	2222	0	2082	33	0
2	B	821	0	796	12	0
2	E	833	0	805	5	0
3	C	77	0	93	2	0
3	F	69	0	80	1	0
4	B	5	0	0	0	0
5	A	67	0	0	1	0
5	B	46	0	0	0	0
5	C	3	0	0	0	0
5	D	78	0	0	2	0
5	E	42	0	0	1	0
5	F	9	0	0	0	0
All	All	6507	0	5948	69	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 6.

All (69) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:D:93:HIS:HD2	1:D:119:GLU:OE1	1.74	0.71
1:D:195:SER:O	1:D:197:GLY:N	2.20	0.70
1:A:220:ASN:H	1:A:222:GLU:HG2	1.58	0.68
1:D:196:LYS:HE2	1:D:251:LEU:CD1	2.23	0.68
1:D:196:LYS:HE2	1:D:251:LEU:HD12	1.74	0.68
2:B:39:MET:H	2:B:45:LYS:HZ2	1.45	0.65
1:A:252:GLY:N	1:A:254:GLU:OE2	2.34	0.60
2:B:38:GLN:HA	2:B:45:LYS:HZ3	1.65	0.60
2:B:39:MET:H	2:B:45:LYS:NZ	2.01	0.57
2:B:50:GLU:HB2	2:B:67:HIS:CE1	2.40	0.56
1:A:141:GLN:OE1	1:A:144[A]:ARG:NH1	2.38	0.56
2:B:48:LYS:HZ1	2:B:69:GLU:H	1.54	0.56
1:D:191:HIS:NE2	1:D:199:VAL:HG21	2.20	0.56
1:A:226:GLN:O	1:A:227:ASP:HB3	2.05	0.56
1:A:169:HIS:O	1:A:173:LYS:HG2	2.06	0.55
1:A:193:PRO:HA	1:A:199:VAL:HA	1.88	0.55
1:A:255:GLN:HA	1:A:255:GLN:OE1	2.06	0.55
2:B:45:LYS:HE3	2:B:46:ILE:O	2.07	0.54
1:D:218:GLN:OE1	1:D:260:ARG:NH1	2.41	0.53

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:D:195:SER:C	1:D:197:GLY:H	2.15	0.51
2:B:29:GLN:HA	2:B:61:SER:HB2	1.93	0.51
1:A:203:CYS:SG	1:A:272:LEU:HD21	2.51	0.51
1:D:260:ARG:HH11	1:D:260:ARG:HG3	1.76	0.51
2:B:48:LYS:HD2	2:B:48:LYS:O	2.12	0.50
1:D:51:TRP:O	1:D:54:GLN:HG3	2.11	0.49
1:D:94:THR:HG23	5:D:326:HOH:O	2.12	0.49
1:D:117:ALA:HB2	2:E:60:TRP:CE2	2.48	0.49
1:D:219:LEU:HD13	1:D:257:TYR:CZ	2.47	0.49
1:D:251:LEU:HD23	1:D:252:GLY:H	1.77	0.48
1:A:81:LEU:HD13	1:A:118:TYR:CD1	2.48	0.48
1:A:33:PHE:C	1:A:48:ARG:HB2	2.39	0.47
1:D:251:LEU:HA	1:D:254:GLU:OE2	2.14	0.47
1:D:45:TYR:CE2	1:D:67:ALA:HB2	2.50	0.47
1:D:255:GLN:NE2	1:D:274:TRP:O	2.44	0.46
1:D:192:HIS:HB2	1:D:200:THR:HB	1.98	0.46
1:D:266:LEU:HD13	1:D:270:LEU:HG	1.99	0.45
1:D:194:ARG:HH11	1:D:248:VAL:HG11	1.81	0.45
1:D:201:LEU:HD12	1:D:249:VAL:HG21	1.97	0.45
1:A:117:ALA:HB2	2:B:60:TRP:CE2	2.51	0.45
2:B:7:ILE:HD12	2:B:91:LYS:HE3	1.98	0.44
2:B:48:LYS:NZ	2:B:69:GLU:H	2.13	0.44
1:D:191:HIS:CE1	1:D:199:VAL:HG21	2.53	0.44
1:A:225:THR:OG1	1:A:226:GLN:N	2.51	0.44
2:E:21:ASN:HB3	2:E:70:PHE:CE1	2.53	0.44
1:D:43:PRO:C	1:D:44:ARG:HG2	2.42	0.43
2:E:1[B]:ILE:HG22	2:E:3:LYS:NZ	2.34	0.43
1:D:98[B]:MET:SD	2:E:58:LYS:HA	2.59	0.43
1:A:220:ASN:HB3	1:A:222:GLU:OE1	2.18	0.43
1:A:66:LYS:HZ1	3:C:1:MET:HG2	1.83	0.43
1:D:218:GLN:HE21	1:D:223:GLU:HG2	1.83	0.43
2:B:38:GLN:HB3	2:B:45:LYS:HD3	2.01	0.43
1:D:73:TRP:CD1	3:F:8:VAL:HG12	2.54	0.43
1:A:44:ARG:HD3	1:A:46:GLU:OE2	2.19	0.42
1:A:108:ARG:HD3	1:D:262:TYR:CG	2.54	0.42
1:D:218:GLN:CD	1:D:260:ARG:NH1	2.77	0.42
1:A:219:LEU:HB3	1:A:222:GLU:CG	2.48	0.42
1:D:218:GLN:HG2	1:D:222:GLU:C	2.44	0.42
1:A:203:CYS:HB2	1:A:217:TRP:CZ2	2.54	0.42
1:D:196:LYS:HA	1:D:196:LYS:HD3	1.77	0.42
1:A:214:THR:HB	1:A:262:TYR:HB2	2.01	0.41

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:66:LYS:NZ	3:C:1:MET:HE2	2.34	0.41
1:A:12:VAL:HG13	5:A:347:HOH:O	2.20	0.41
1:A:32:GLU:OE2	1:A:48:ARG:HD2	2.20	0.41
1:D:190:THR:OG1	1:D:192:HIS:HE1	2.02	0.41
1:A:46:GLU:OE1	1:A:47:PRO:HD2	2.20	0.41
1:D:266:LEU:HB3	1:D:268:GLU:O	2.21	0.40
2:E:1[B]:ILE:HG12	5:E:121:HOH:O	2.20	0.40
1:D:94:THR:HG22	5:D:368:HOH:O	2.21	0.40
1:D:194:ARG:O	1:D:196:LYS:N	2.54	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	269/276 (98%)	260 (97%)	8 (3%)	1 (0%)	30	27
1	D	268/276 (97%)	260 (97%)	6 (2%)	2 (1%)	18	14
2	B	97/99 (98%)	93 (96%)	4 (4%)	0	100	100
2	E	98/99 (99%)	97 (99%)	1 (1%)	0	100	100
3	C	7/9 (78%)	6 (86%)	1 (14%)	0	100	100
3	F	6/9 (67%)	4 (67%)	2 (33%)	0	100	100
All	All	745/768 (97%)	720 (97%)	22 (3%)	3 (0%)	30	27

All (3) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	D	196	LYS
1	D	195	SER
1	A	219	LEU

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	230/234 (98%)	227 (99%)	3 (1%)	61	68
1	D	229/234 (98%)	229 (100%)	0	100	100
2	B	94/94 (100%)	93 (99%)	1 (1%)	65	73
2	E	95/94 (101%)	94 (99%)	1 (1%)	65	73
3	C	8/7 (114%)	8 (100%)	0	100	100
3	F	7/7 (100%)	7 (100%)	0	100	100
All	All	663/670 (99%)	658 (99%)	5 (1%)	73	80

All (5) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	219	LEU
1	A	220	ASN
1	A	222	GLU
2	B	64	ILE
2	E	57	SER

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (15) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	42	ASN
1	A	96	GLN
1	A	220	ASN
2	B	13	HIS
2	B	29	GLN
1	D	80	ASN
1	D	87	GLN
1	D	93	HIS
1	D	115	GLN
1	D	192	HIS
1	D	220	ASN
1	D	256	ASN

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
2	E	2	GLN
2	E	34	HIS
2	E	38	GLN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

2 non-standard protein/DNA/RNA residues are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
3	NLE	F	5	3	6,7,8	0.52	0	2,7,9	0.35	0
3	NLE	C	5	3	6,7,8	0.88	0	2,7,9	0.43	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	NLE	F	5	3	-	0/5/6/8	-
3	NLE	C	5	3	-	0/5/6/8	-

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

1 ligand is modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
4	SO4	B	101	-	4,4,4	0.26	0	6,6,6	0.09	0

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	272/276 (98%)	0.67	41 (15%) 5 5	14, 30, 84, 113	1 (0%)
1	D	271/276 (98%)	0.56	36 (13%) 7 6	10, 29, 78, 105	1 (0%)
2	B	99/99 (100%)	0.15	4 (4%) 42 41	15, 24, 40, 65	0
2	E	99/99 (100%)	-0.03	3 (3%) 52 51	10, 20, 37, 62	2 (2%)
3	C	8/9 (88%)	-0.01	0 100 100	15, 19, 27, 33	1 (12%)
3	F	8/9 (88%)	-0.11	0 100 100	15, 18, 27, 31	0
All	All	757/768 (98%)	0.46	84 (11%) 10 9	10, 27, 77, 113	5 (0%)

All (84) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
2	B	1	ILE	6.1
1	A	16	GLY	6.0
1	A	225	THR	5.4
1	D	224	LEU	5.0
1	A	224	LEU	4.8
1	A	197	GLY	4.6
1	A	17	LEU	4.6
2	B	48	LYS	4.5
1	A	193	PRO	4.5
1	D	274	TRP	4.4
1	D	221	GLY	4.3
1	D	227	ASP	4.2
1	A	251	LEU	4.2
1	A	274	TRP	4.1
1	D	225	THR	4.1
1	D	219	LEU	3.9
1	D	195	SER	3.9
1	A	196	LYS	3.8
1	D	251	LEU	3.8

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	195	SER	3.7
1	A	257	TYR	3.7
1	D	197	GLY	3.7
2	E	54[A]	MET	3.6
1	A	219	LEU	3.6
1	A	250	PRO	3.5
1	A	276	PRO	3.5
1	D	90	GLY	3.5
1	D	175	GLY	3.4
1	D	196	LYS	3.4
1	D	220	ASN	3.3
1	A	176	ASN	3.3
1	A	221	GLY	3.3
1	D	193	PRO	3.2
1	D	249	VAL	3.2
1	A	252	GLY	3.2
1	D	89	ALA	3.1
1	D	248	VAL	3.1
1	A	256	ASN	3.1
1	A	253	LYS	3.1
1	A	220	ASN	3.1
1	D	250	PRO	3.0
1	A	199	VAL	3.0
1	A	258	THR	3.0
1	A	248	VAL	3.0
1	A	249	VAL	3.0
1	D	16	GLY	2.9
2	E	48	LYS	2.9
1	D	17	LEU	2.8
1	A	194	ARG	2.8
1	D	200	THR	2.8
1	D	194	ARG	2.8
1	A	42	ASN	2.7
1	D	253	LYS	2.7
2	E	1[A]	ILE	2.7
1	D	181	ARG	2.7
1	A	201	LEU	2.7
1	A	15	PRO	2.7
1	D	276	PRO	2.6
1	A	191	HIS	2.6
1	A	246	SER	2.6
1	A	272	LEU	2.6

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	D	252	GLY	2.5
1	A	254	GLU	2.3
1	A	200	THR	2.3
1	D	41	GLU	2.3
1	D	199	VAL	2.3
1	A	175	GLY	2.3
1	A	192	HIS	2.3
1	A	181	ARG	2.3
1	D	217	TRP	2.3
1	A	227	ASP	2.2
1	D	223	GLU	2.2
1	D	275	GLU	2.2
1	D	31	LYS	2.2
2	B	45	LYS	2.2
1	A	1	GLY	2.2
1	D	15	PRO	2.2
2	B	47	PRO	2.2
1	A	222	GLU	2.1
1	A	228	MET	2.1
1	D	222	GLU	2.1
1	A	255	GLN	2.1
1	D	257	TYR	2.1
1	D	50	PRO	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
3	NLE	C	5	8/9	0.94	0.09	16,21,26,26	0
3	NLE	F	5	8/9	0.94	0.07	13,18,19,19	0

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
4	SO4	B	101	5/5	0.70	0.15	77,81,84,87	0

6.5 Other polymers

There are no such residues in this entry.