



# wwPDB X-ray Structure Validation Summary Report ⓘ

Mar 8, 2026 – 05:51 PM UTC

PDB ID : 5E93 / pdb\_00005e93  
Title : Crystal Structure of Trypanosoma cruzi Dihydroorotate Dehydrogenase in Complex with Neq0071  
Authors : Rocha, J.R.; Inaoka, D.K.; Cheleski, J.; Shiba, T.; Harada, S.; Montanari, C.A.; Kita, K.  
Deposited on : 2015-10-14  
Resolution : 1.41 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0  
Mogul : 2022.3.0, CSD as543be (2022)  
Xtrriage (Phenix) : 2.0  
EDS : 3.0  
Buster-report : wwPDB partial adaption of 1.1.7 (2018)  
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)  
CCP4 : 9.0.010 (Gargrove)  
Density-Fitness : 1.0.12  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.49


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*



The reported resolution of this entry is 1.41 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	180053	4041 (1.44-1.40)
Clashscore	190562	4154 (1.44-1.40)
Ramachandran outliers	187476	4083 (1.44-1.40)
Sidechain outliers	187428	4082 (1.44-1.40)
RSRZ outliers	180081	4039 (1.44-1.40)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	314	
1	B	314	

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
2	5LL	A	401	-	X	-	-
3	GOL	A	405	-	X	-	-
6	EDO	A	417	-	-	X	-
6	EDO	A	418	-	-	X	-
6	EDO	B	414	-	-	X	-
6	EDO	B	416	-	-	X	-
6	EDO	B	421	-	-	X	-
7	CAC	A	421	-	X	-	-
7	CAC	A	422	-	X	-	-
8	PEG	B	423	-	X	X	-

## 2 Entry composition [i](#)

There are 9 unique types of molecules in this entry. The entry contains 5936 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called Dihydroorotate dehydrogenase (fumarate).

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	313	2498	1607	407	463	21	0	19	0
1	B	313	2468	1586	406	456	20	0	14	0

There are 2 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	-1	SER	-	expression tag	UNP Q4D3W2
B	-1	SER	-	expression tag	UNP Q4D3W2

- Molecule 2 is 5-[(E)-3-(furan-2-yl)prop-2-enylidene]-1,3-diazinane-2,4,6-trione (CCD ID: 5LL) (formula: C<sub>11</sub>H<sub>8</sub>N<sub>2</sub>O<sub>4</sub>).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
			Total	C	N	O		
2	A	1	17	11	2	4	0	0

*Continued on next page...*

Continued from previous page...

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
			Total	C	N	O		
2	B	1	17	11	2	4	0	0

- Molecule 3 is GLYCEROL (CCD ID: GOL) (formula: C<sub>3</sub>H<sub>8</sub>O<sub>3</sub>).




Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
			Total	C	O		
3	A	1	6	3	3	0	0
3	A	1	6	3	3	0	0
3	A	1	6	3	3	0	0
3	A	1	6	3	3	0	0
3	A	1	6	3	3	0	0
3	A	1	6	3	3	0	0
3	B	1	6	3	3	0	0
3	B	1	6	3	3	0	0
3	B	1	6	3	3	0	0
3	B	1	6	3	3	0	0

Continued on next page...

Continued from previous page...


Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	B	1	Total	C	O	0	0
			6	3	3		
3	B	1	Total	C	O	0	1
			12	6	6		

- Molecule 4 is FLAVIN MONONUCLEOTIDE (CCD ID: FMN) (formula: C<sub>17</sub>H<sub>21</sub>N<sub>4</sub>O<sub>9</sub>P).




Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
4	A	1	Total	C	N	O	P	0	0
			31	17	4	9	1		
4	B	1	Total	C	N	O	P	0	0
			31	17	4	9	1		

- Molecule 5 is COBALT HEXAMMINE(III) (CCD ID: NCO) (formula: CoH<sub>18</sub>N<sub>6</sub>).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
5	A	1	Total	Co	N	0	0
			7	1	6		
5	B	1	Total	Co	N	0	0
			7	1	6		
5	B	1	Total	Co	N	0	1
			14	2	12		

- Molecule 6 is 1,2-ETHANEDIOL (CCD ID: EDO) (formula: C<sub>2</sub>H<sub>6</sub>O<sub>2</sub>).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
6	A	1	Total	C	O	0	0
			4	2	2		

*Continued on next page...*

*Continued from previous page...*


Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
6	A	1	Total 4	C 2	O 2	0	0
6	A	1	Total 4	C 2	O 2	0	0
6	A	1	Total 4	C 2	O 2	0	0
6	A	1	Total 8	C 4	O 4	0	1
6	A	1	Total 4	C 2	O 2	0	0
6	A	1	Total 4	C 2	O 2	0	0
6	A	1	Total 4	C 2	O 2	0	0
6	A	1	Total 4	C 2	O 2	0	0
6	A	1	Total 4	C 2	O 2	0	0
6	A	1	Total 4	C 2	O 2	0	0
6	A	1	Total 4	C 2	O 2	0	0
6	B	1	Total 4	C 2	O 2	0	0
6	B	1	Total 4	C 2	O 2	0	0
6	B	1	Total 4	C 2	O 2	0	0
6	B	1	Total 4	C 2	O 2	0	0
6	B	1	Total 4	C 2	O 2	0	0
6	B	1	Total 4	C 2	O 2	0	0
6	B	1	Total 4	C 2	O 2	0	0
6	B	1	Total 4	C 2	O 2	0	0
6	B	1	Total 4	C 2	O 2	0	0
6	B	1	Total 4	C 2	O 2	0	0
6	B	1	Total 4	C 2	O 2	0	0

*Continued on next page...*

Continued from previous page...


Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
			Total	C	O		
6	B	1	4	2	2	0	0

- Molecule 7 is CACODYLATE ION (CCD ID: CAC) (formula:  $C_2H_6AsO_2$ ).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
			Total	As	C	O		
7	A	1	5	1	2	2	0	0
7	A	1	Total	As	C	O	0	0
			5	1	2	2		

- Molecule 8 is DI(HYDROXYETHYL)ETHER (CCD ID: PEG) (formula:  $C_4H_{10}O_3$ ).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
8	B	1	Total C O 7 4 3	0	0


- Molecule 9 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
9	A	330	Total O 330 330	0	0
9	B	325	Total O 325 325	0	0


### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Dihydroorotate dehydrogenase (fumarate)



- Molecule 1: Dihydroorotate dehydrogenase (fumarate)



## 4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	68.33Å 71.79Å 124.34Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	50.00 – 1.41 50.00 – 1.41	Depositor EDS
% Data completeness (in resolution range)	97.9 (50.00-1.41) 98.0 (50.00-1.41)	Depositor EDS
$R_{merge}$	0.08	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	5.04 (at 1.41Å)	Xtrriage
Refinement program	REFMAC 5.8.0103	Depositor
R, $R_{free}$	0.145 , 0.168 0.145 , 0.169	Depositor DCC
$R_{free}$ test set	5946 reflections (5.02%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	13.7	Xtrriage
Anisotropy	0.039	Xtrriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.38 , 41.4	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.50$ , $\langle L^2 \rangle = 0.34$	Xtrriage
Estimated twinning fraction	0.000 for k,h,-l	Xtrriage
$F_o, F_c$ correlation	0.97	EDS
Total number of atoms	5936	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	18.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The analyses of the Patterson function reveals a significant off-origin peak that is 59.13 % of the origin peak, indicating pseudo-translational symmetry. The chance of finding a peak of this or larger height randomly in a structure without pseudo-translational symmetry is equal to 1.8480e-05. The detected translational NCS is most likely also responsible for the elevated intensity ratio.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: PEG, CAC, 5LL, FMN, GOL, NCO, EDO

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	A	1.59	17/2599 (0.7%)	1.33	12/3515 (0.3%)
1	B	1.69	27/2559 (1.1%)	1.46	20/3458 (0.6%)
All	All	1.64	44/5158 (0.9%)	1.39	32/6973 (0.5%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	1
1	B	0	2
All	All	0	3

The worst 5 of 44 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	71	LEU	CA-C	12.15	1.68	1.52
1	B	71	LEU	CA-C	9.52	1.64	1.52
1	A	71	LEU	C-N	9.32	1.42	1.33
1	A	266	SER	C-O	-9.26	1.13	1.24
1	B	114	ALA	C-O	-8.83	1.16	1.24

The worst 5 of 32 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	71	LEU	CA-C-N	-11.81	105.08	119.84
1	B	71	LEU	C-N-CA	-11.81	105.08	119.84
1	A	4	LEU	O-C-N	11.43	138.05	123.19
1	A	71	LEU	CA-C-N	-9.86	105.96	120.46

*Continued on next page...*

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	71	LEU	C-N-CA	-9.86	105.96	120.46

There are no chirality outliers.

All (3) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	71	LEU	Peptide
1	B	50	ARG	Mainchain
1	B	71	LEU	Peptide

## 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2498	0	2536	47	0
1	B	2468	0	2514	37	0
2	A	17	0	0	1	0
2	B	17	0	0	1	0
3	A	36	0	46	6	0
3	B	42	0	56	0	0
4	A	31	0	19	1	0
4	B	31	0	19	0	0
5	A	7	0	0	0	0
5	B	21	0	0	3	0
6	A	48	0	69	25	0
6	B	48	0	70	21	0
7	A	10	0	0	3	0
8	B	7	0	10	5	0
9	A	330	0	0	9	0
9	B	325	0	0	11	0
All	All	5936	0	5339	105	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 10.

The worst 5 of 105 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:3[B]:LYS:CE	1:B:3[B]:LYS:NZ	1.80	1.42
1:A:50:ARG:HD3	1:A:71:LEU:HD12	1.10	1.06
1:A:50:ARG:HD3	1:A:71:LEU:CD1	1.86	1.05
1:A:138:GLN:HE21	1:A:138:GLN:H	1.12	0.96
6:B:414:EDO:H22	6:B:421:EDO:C1	1.96	0.94

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	331/314 (105%)	321 (97%)	9 (3%)	1 (0%)	36 17
1	B	326/314 (104%)	316 (97%)	9 (3%)	1 (0%)	36 17
All	All	657/628 (105%)	637 (97%)	18 (3%)	2 (0%)	36 17

All (2) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	B	71	LEU
1	A	71	LEU

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	278/259 (107%)	274 (99%)	4 (1%)	59 27
1	B	273/259 (105%)	268 (98%)	5 (2%)	51 18
All	All	551/518 (106%)	542 (98%)	9 (2%)	55 22

5 of 9 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	B	138	GLN
1	B	312	GLU
1	A	214[C]	LYS
1	B	0	MET
1	B	71	LEU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 12 such sidechains are listed below:

Mol	Chain	Res	Type
1	B	9	HIS
1	B	87	HIS
1	B	275	GLN
1	B	138	GLN
1	A	118	GLN

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

### 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

## 5.6 Ligand geometry [i](#)

48 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
6	EDO	B	420	-	3,3,3	1.86	1 (33%)	2,2,2	0.74	0
2	5LL	B	401	1	18,18,18	2.46	7 (38%)	24,24,24	2.94	16 (66%)
6	EDO	B	415	-	3,3,3	0.54	0	2,2,2	0.66	0
3	GOL	B	406	-	5,5,5	0.97	0	5,5,5	0.79	0
6	EDO	B	418	-	3,3,3	0.42	0	2,2,2	0.43	0
3	GOL	B	403	-	5,5,5	0.51	0	5,5,5	0.94	0
2	5LL	A	401	1	18,18,18	2.77	8 (44%)	24,24,24	3.57	16 (66%)
6	EDO	A	416	-	3,3,3	1.49	1 (33%)	2,2,2	1.59	0
6	EDO	B	419	-	3,3,3	0.22	0	2,2,2	0.68	0
8	PEG	B	423	-	6,6,6	1.59	2 (33%)	5,5,5	6.23	5 (100%)
5	NCO	B	410[A]	-	6,6,6	0.92	0	-	-	-
3	GOL	A	403	-	5,5,5	0.40	0	5,5,5	0.60	0
3	GOL	A	405	-	5,5,5	1.24	1 (20%)	5,5,5	3.49	4 (80%)
6	EDO	B	412	-	3,3,3	0.45	0	2,2,2	0.72	0
5	NCO	B	410[B]	-	6,6,6	1.61	1 (16%)	-	-	-
3	GOL	A	402	-	5,5,5	0.85	0	5,5,5	1.24	1 (20%)
3	GOL	B	407[A]	-	5,5,5	0.66	0	5,5,5	1.03	0
6	EDO	B	411	-	3,3,3	0.56	0	2,2,2	0.40	0
6	EDO	A	420	-	3,3,3	1.72	1 (33%)	2,2,2	1.35	0
3	GOL	B	407[B]	-	5,5,5	0.67	0	5,5,5	0.99	0
3	GOL	A	406	-	5,5,5	1.12	1 (20%)	5,5,5	0.95	0
4	FMN	A	408	-	33,33,33	1.14	2 (6%)	48,50,50	1.28	4 (8%)
6	EDO	A	410	-	3,3,3	1.05	0	2,2,2	0.79	0
6	EDO	B	421	-	3,3,3	0.95	0	2,2,2	0.78	0
7	CAC	A	421	-	2,4,4	2.19	1 (50%)	4,6,6	4.28	3 (75%)
4	FMN	B	408	-	33,33,33	1.47	5 (15%)	48,50,50	1.04	3 (6%)
7	CAC	A	422	-	2,4,4	3.16	1 (50%)	4,6,6	5.43	3 (75%)
6	EDO	B	414	-	3,3,3	0.72	0	2,2,2	1.79	1 (50%)

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
5	NCO	B	409	-	6,6,6	4.22	6 (100%)	-		
6	EDO	B	417	-	3,3,3	0.32	0	2,2,2	1.02	0
6	EDO	A	411	-	3,3,3	0.63	0	2,2,2	0.04	0
6	EDO	A	414[A]	-	3,3,3	0.84	0	2,2,2	0.21	0
3	GOL	B	404	-	5,5,5	0.61	0	5,5,5	0.88	0
3	GOL	B	405	-	5,5,5	0.95	0	5,5,5	2.52	3 (60%)
6	EDO	A	417	-	3,3,3	1.69	1 (33%)	2,2,2	0.58	0
5	NCO	A	409	-	6,6,6	1.68	3 (50%)	-		
6	EDO	A	414[B]	-	3,3,3	0.84	0	2,2,2	0.33	0
6	EDO	B	413	-	3,3,3	0.92	0	2,2,2	1.92	1 (50%)
6	EDO	B	416	-	3,3,3	1.03	0	2,2,2	0.63	0
3	GOL	A	404	-	5,5,5	0.86	0	5,5,5	0.80	0
3	GOL	B	402	-	5,5,5	0.54	0	5,5,5	0.78	0
3	GOL	A	407	-	5,5,5	1.44	1 (20%)	5,5,5	1.37	1 (20%)
6	EDO	A	415	-	3,3,3	0.41	0	2,2,2	1.26	0
6	EDO	A	413	-	3,3,3	0.41	0	2,2,2	0.27	0
6	EDO	A	418	-	3,3,3	1.10	0	2,2,2	1.79	1 (50%)
6	EDO	A	412	-	3,3,3	0.65	0	2,2,2	0.96	0
6	EDO	A	419	-	3,3,3	0.59	0	2,2,2	0.35	0
6	EDO	B	422	-	3,3,3	0.63	0	2,2,2	0.56	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
6	EDO	B	420	-	-	0/1/1/1	-
2	5LL	B	401	1	-	3/6/22/22	0/2/2/2
6	EDO	B	415	-	-	1/1/1/1	-
3	GOL	B	406	-	-	0/4/4/4	-
6	EDO	B	418	-	-	1/1/1/1	-
3	GOL	B	403	-	-	0/4/4/4	-
2	5LL	A	401	1	-	3/6/22/22	0/2/2/2
6	EDO	A	416	-	-	1/1/1/1	-
6	EDO	B	419	-	-	0/1/1/1	-
8	PEG	B	423	-	-	4/4/4/4	-
3	GOL	A	403	-	-	2/4/4/4	-
3	GOL	A	405	-	-	2/4/4/4	-
6	EDO	B	412	-	-	0/1/1/1	-

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	GOL	A	402	-	-	2/4/4/4	-
3	GOL	B	407[A]	-	-	0/4/4/4	-
6	EDO	B	411	-	-	1/1/1/1	-
6	EDO	A	420	-	-	1/1/1/1	-
3	GOL	B	407[B]	-	-	0/4/4/4	-
3	GOL	A	406	-	-	0/4/4/4	-
4	FMN	A	408	-	-	1/18/18/18	0/3/3/3
6	EDO	A	410	-	-	1/1/1/1	-
6	EDO	B	421	-	-	1/1/1/1	-
4	FMN	B	408	-	-	1/18/18/18	0/3/3/3
6	EDO	B	414	-	-	1/1/1/1	-
6	EDO	B	417	-	-	1/1/1/1	-
6	EDO	A	411	-	-	0/1/1/1	-
6	EDO	A	414[A]	-	-	1/1/1/1	-
3	GOL	B	404	-	-	0/4/4/4	-
3	GOL	B	405	-	-	0/4/4/4	-
6	EDO	A	417	-	-	1/1/1/1	-
6	EDO	A	414[B]	-	-	0/1/1/1	-
6	EDO	B	413	-	-	1/1/1/1	-
6	EDO	B	416	-	-	1/1/1/1	-
3	GOL	A	404	-	-	0/4/4/4	-
3	GOL	B	402	-	-	0/4/4/4	-
3	GOL	A	407	-	-	0/4/4/4	-
6	EDO	A	415	-	-	1/1/1/1	-
6	EDO	A	413	-	-	1/1/1/1	-
6	EDO	A	418	-	-	1/1/1/1	-
6	EDO	A	412	-	-	1/1/1/1	-
6	EDO	A	419	-	-	1/1/1/1	-
6	EDO	B	422	-	-	0/1/1/1	-

The worst 5 of 43 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
5	B	409	NCO	CO-N5	6.49	2.17	1.96
2	B	401	5LL	C2-N1	-6.05	1.27	1.37
2	A	401	5LL	OAD-CAE	-5.42	1.28	1.37
5	B	409	NCO	CO-N3	5.34	2.13	1.96
2	A	401	5LL	CAH-C5	5.19	1.51	1.37

The worst 5 of 62 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
8	B	423	PEG	O2-C2-C1	8.12	145.91	110.11
8	B	423	PEG	O2-C3-C4	7.31	142.31	110.11
8	B	423	PEG	C3-O2-C2	7.19	144.71	113.26
2	B	401	5LL	CAF-CAE-CAA	-7.05	117.54	132.30
2	A	401	5LL	CAH-CAG-CAF	-6.90	103.20	123.20

There are no chirality outliers.

5 of 36 torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	A	401	5LL	C4-C5-CAH-CAG
2	A	401	5LL	C6-C5-CAH-CAG
2	A	401	5LL	CAF-CAG-CAH-C5
2	B	401	5LL	C4-C5-CAH-CAG
2	B	401	5LL	C6-C5-CAH-CAG

There are no ring outliers.

25 monomers are involved in 61 short contacts:


Mol	Chain	Res	Type	Clashes	Symm-Clashes
6	B	420	EDO	1	0
2	B	401	5LL	1	0
2	A	401	5LL	1	0
6	A	416	EDO	3	0
8	B	423	PEG	5	0
3	A	403	GOL	1	0
3	A	405	GOL	3	0
6	B	412	EDO	3	0
5	B	410[B]	NCO	1	0
6	A	420	EDO	3	0
3	A	406	GOL	2	0
4	A	408	FMN	1	0
6	A	410	EDO	2	0
6	B	421	EDO	6	0
7	A	421	CAC	3	0
6	B	414	EDO	10	0
5	B	409	NCO	2	0
6	A	414[A]	EDO	1	0
6	A	417	EDO	4	0
6	B	413	EDO	2	0
6	B	416	EDO	4	0
6	A	415	EDO	1	0


*Continued on next page...*

*Continued from previous page...*

<b>Mol</b>	<b>Chain</b>	<b>Res</b>	<b>Type</b>	<b>Clashes</b>	<b>Symm-Clashes</b>
6	A	418	EDO	5	0
6	A	412	EDO	3	0
6	A	419	EDO	3	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.





## 5.7 Other polymers [\(i\)](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [\(i\)](#)

There are no chain breaks in this entry.

## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	313/314 (99%)	-0.07	14 (4%) 38 40	6, 13, 28, 46	19 (6%)
1	B	313/314 (99%)	0.01	13 (4%) 40 43	7, 14, 31, 48	14 (4%)
All	All	626/628 (99%)	-0.03	27 (4%) 40 42	6, 14, 30, 48	33 (5%)

The worst 5 of 27 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	71	LEU	6.1
1	B	71	LEU	5.2
1	B	52	GLY	4.9
1	A	54	PRO	4.7
1	B	133	VAL	4.3

### 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

### 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.


Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors( $\text{\AA}^2$ )	Q<0.9
3	GOL	A	403	6/6	0.69	0.18	38,41,44,46	0
6	EDO	B	420	4/4	0.73	0.19	26,26,30,33	0
6	EDO	A	418	4/4	0.74	0.15	26,26,33,35	0
6	EDO	A	414[A]	4/4	0.75	0.13	25,28,30,34	4
6	EDO	A	414[B]	4/4	0.75	0.13	25,27,29,32	4
7	CAC	A	421	5/5	0.75	0.26	15,22,33,33	0
7	CAC	A	422	5/5	0.75	0.28	21,22,35,35	0
3	GOL	B	403	6/6	0.77	0.15	36,40,47,48	0
6	EDO	B	421	4/4	0.77	0.15	32,35,37,37	0
5	NCO	A	409	7/7	0.78	0.27	34,38,39,43	0
3	GOL	A	406	6/6	0.78	0.13	28,29,31,36	0
6	EDO	B	415	4/4	0.81	0.15	35,41,43,53	0
6	EDO	A	410	4/4	0.82	0.13	32,32,35,44	0
6	EDO	A	413	4/4	0.84	0.13	39,45,46,51	0
6	EDO	A	420	4/4	0.85	0.14	18,27,29,36	0
3	GOL	A	402	6/6	0.86	0.11	27,30,32,39	0
6	EDO	B	419	4/4	0.86	0.13	29,36,42,43	0
6	EDO	B	412	4/4	0.86	0.11	32,37,39,40	0
6	EDO	B	414	4/4	0.87	0.11	20,26,28,43	0
6	EDO	B	411	4/4	0.87	0.14	32,37,47,52	0
6	EDO	B	417	4/4	0.87	0.15	25,40,41,63	0
3	GOL	A	407	6/6	0.87	0.11	28,32,34,41	0
6	EDO	A	415	4/4	0.88	0.11	25,30,32,35	0
6	EDO	B	422	4/4	0.88	0.08	27,27,28,28	0
6	EDO	B	416	4/4	0.89	0.11	23,26,26,28	0
6	EDO	A	411	4/4	0.89	0.09	37,38,38,47	0
3	GOL	B	406	6/6	0.90	0.10	23,24,24,25	0
6	EDO	A	419	4/4	0.90	0.10	28,28,30,30	0
3	GOL	A	404	6/6	0.90	0.10	20,25,27,28	0
6	EDO	A	412	4/4	0.91	0.11	25,26,27,29	0
2	5LL	B	401	17/17	0.91	0.11	12,22,37,39	0
3	GOL	A	405	6/6	0.92	0.08	21,26,32,32	0
2	5LL	A	401	17/17	0.92	0.12	13,21,37,38	0
6	EDO	A	417	4/4	0.93	0.09	22,26,29,29	0
6	EDO	B	418	4/4	0.93	0.08	36,36,39,40	0
6	EDO	B	413	4/4	0.94	0.08	25,27,29,45	0
5	NCO	B	410[A]	7/7	0.94	0.17	21,23,24,25	7
5	NCO	B	410[B]	7/7	0.94	0.17	20,22,25,26	7
3	GOL	B	405	6/6	0.94	0.11	14,31,34,36	0
3	GOL	B	402	6/6	0.94	0.07	18,23,25,26	0
3	GOL	B	404	6/6	0.94	0.08	17,23,26,28	0
8	PEG	B	423	7/7	0.94	0.10	15,18,26,26	0
6	EDO	A	416	4/4	0.96	0.12	18,21,29,32	0


*Continued on next page...*

Continued from previous page...

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å <sup>2</sup> )	Q<0.9
3	GOL	B	407[A]	6/6	0.97	0.05	12,17,17,17	6
3	GOL	B	407[B]	6/6	0.97	0.05	14,17,18,18	6
5	NCO	B	409	7/7	0.99	0.06	12,13,14,14	0
4	FMN	B	408	31/31	0.99	0.04	8,10,12,13	0
4	FMN	A	408	31/31	0.99	0.04	8,9,11,13	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.





## 6.5 Other polymers [i](#)

There are no such residues in this entry.