



wwPDB X-ray Structure Validation Summary Report ⓘ

Mar 5, 2026 – 06:40 PM UTC

PDB ID : 5E9G / pdb_00005e9g
Title : Structural insights of isocitrate lyases from Magnaporthe oryzae
Authors : Park, Y.; Cho, Y.; Lee, Y.-H.; Lee, Y.-W.; Rhee, S.
Deposited on : 2015-10-15
Resolution : 2.10 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.10 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	6658 (2.10-2.10)
Clashscore	190562	7164 (2.10-2.10)
Ramachandran outliers	187476	7099 (2.10-2.10)
Sidechain outliers	187428	7100 (2.10-2.10)
RSRZ outliers	180081	6662 (2.10-2.10)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	560	4% 69% 24% 6%
1	B	560	2% 68% 24% 6%
1	C	560	8% 65% 22% 11%
1	D	560	7% 63% 21% 13%

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard

residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
2	GLV	A	601	-	X	-	-
2	GLV	B	601	-	X	X	-
2	GLV	D	601	-	-	X	-

2 Entry composition

There are 5 unique types of molecules in this entry. The entry contains 16409 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Isocitrate lyase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	524	Total	C	N	O	S	0	0	0
			4130	2606	720	783	21			
1	B	525	Total	C	N	O	S	0	0	0
			4134	2608	721	784	21			
1	C	499	Total	C	N	O	S	0	0	0
			3949	2501	686	741	21			
1	D	486	Total	C	N	O	S	0	0	0
			3850	2437	672	720	21			

There are 56 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
A	219	GLY	ALA	engineered mutation	UNP P0CT06
A	548	ALA	-	expression tag	UNP P0CT06
A	549	ALA	-	expression tag	UNP P0CT06
A	550	ALA	-	expression tag	UNP P0CT06
A	551	LEU	-	expression tag	UNP P0CT06
A	552	GLU	-	expression tag	UNP P0CT06
A	553	HIS	-	expression tag	UNP P0CT06
A	554	HIS	-	expression tag	UNP P0CT06
A	555	HIS	-	expression tag	UNP P0CT06
A	556	HIS	-	expression tag	UNP P0CT06
A	557	HIS	-	expression tag	UNP P0CT06
A	558	HIS	-	expression tag	UNP P0CT06
A	559	HIS	-	expression tag	UNP P0CT06
A	560	HIS	-	expression tag	UNP P0CT06
B	219	GLY	ALA	engineered mutation	UNP P0CT06
B	548	ALA	-	expression tag	UNP P0CT06
B	549	ALA	-	expression tag	UNP P0CT06
B	550	ALA	-	expression tag	UNP P0CT06
B	551	LEU	-	expression tag	UNP P0CT06
B	552	GLU	-	expression tag	UNP P0CT06
B	553	HIS	-	expression tag	UNP P0CT06

Continued on next page...

Continued from previous page...


Chain	Residue	Modelled	Actual	Comment	Reference
B	554	HIS	-	expression tag	UNP P0CT06
B	555	HIS	-	expression tag	UNP P0CT06
B	556	HIS	-	expression tag	UNP P0CT06
B	557	HIS	-	expression tag	UNP P0CT06
B	558	HIS	-	expression tag	UNP P0CT06
B	559	HIS	-	expression tag	UNP P0CT06
B	560	HIS	-	expression tag	UNP P0CT06
C	219	GLY	ALA	engineered mutation	UNP P0CT06
C	548	ALA	-	expression tag	UNP P0CT06
C	549	ALA	-	expression tag	UNP P0CT06
C	550	ALA	-	expression tag	UNP P0CT06
C	551	LEU	-	expression tag	UNP P0CT06
C	552	GLU	-	expression tag	UNP P0CT06
C	553	HIS	-	expression tag	UNP P0CT06
C	554	HIS	-	expression tag	UNP P0CT06
C	555	HIS	-	expression tag	UNP P0CT06
C	556	HIS	-	expression tag	UNP P0CT06
C	557	HIS	-	expression tag	UNP P0CT06
C	558	HIS	-	expression tag	UNP P0CT06
C	559	HIS	-	expression tag	UNP P0CT06
C	560	HIS	-	expression tag	UNP P0CT06
D	219	GLY	ALA	engineered mutation	UNP P0CT06
D	548	ALA	-	expression tag	UNP P0CT06
D	549	ALA	-	expression tag	UNP P0CT06
D	550	ALA	-	expression tag	UNP P0CT06
D	551	LEU	-	expression tag	UNP P0CT06
D	552	GLU	-	expression tag	UNP P0CT06
D	553	HIS	-	expression tag	UNP P0CT06
D	554	HIS	-	expression tag	UNP P0CT06
D	555	HIS	-	expression tag	UNP P0CT06
D	556	HIS	-	expression tag	UNP P0CT06
D	557	HIS	-	expression tag	UNP P0CT06
D	558	HIS	-	expression tag	UNP P0CT06
D	559	HIS	-	expression tag	UNP P0CT06
D	560	HIS	-	expression tag	UNP P0CT06

- Molecule 2 is GLYOXYLIC ACID (CCD ID: GLV) (formula: $C_2H_2O_3$).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
2	A	1	Total	C	O	0	0
			5	2	3		
2	B	1	Total	C	O	0	0
			5	2	3		
2	C	1	Total	C	O	0	0
			5	2	3		
2	D	1	Total	C	O	0	0
			5	2	3		

- Molecule 3 is GLYCEROL (CCD ID: GOL) (formula: $C_3H_8O_3$).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	A	1	Total	C	O	0	0
			6	3	3		
3	B	1	Total	C	O	0	0
			6	3	3		
3	D	1	Total	C	O	0	0
			6	3	3		

- Molecule 4 is MAGNESIUM ION (CCD ID: MG) (formula: Mg).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	1	Total	Mg	0	0
			1	1		
4	B	1	Total	Mg	0	0
			1	1		
4	C	1	Total	Mg	0	0
			1	1		
4	D	1	Total	Mg	0	0
			1	1		


- Molecule 5 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	89	Total	O	0	0
			89	89		
5	B	82	Total	O	0	0
			82	82		
5	C	73	Total	O	0	0
			73	73		
5	D	60	Total	O	0	0
			60	60		


3 Residue-property plots [i](#)


These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: Isocitrate lyase




• Molecule 1: Isocitrate lyase






● Molecule 1: Isocitrate lyase



● Molecule 1: Isocitrate lyase



4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	121.12Å 135.08Å 158.70Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	50.00 – 2.10 50.00 – 2.10	Depositor EDS
% Data completeness (in resolution range)	90.8 (50.00-2.10) 90.8 (50.00-2.10)	Depositor EDS
R_{merge}	0.08	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	3.63 (at 2.10Å)	Xtriage
Refinement program	CNS 1.3	Depositor
R, R_{free}	0.224 , 0.259 0.224 , 0.259	Depositor DCC
R_{free} test set	14129 reflections (9.93%)	wwPDB-VP
Wilson B-factor (Å ²)	29.9	Xtriage
Anisotropy	0.616	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.34 , 30.8	EDS
L-test for twinning ²	$\langle L \rangle = 0.48$, $\langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.94	EDS
Total number of atoms	16409	wwPDB-VP
Average B, all atoms (Å ²)	36.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 2.38% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: GLV, GOL, MG

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# $ Z > 5$	RMSZ	# $ Z > 5$
1	A	0.42	0/4224	0.92	9/5723 (0.2%)
1	B	0.41	0/4228	0.92	15/5728 (0.3%)
1	C	0.41	0/4040	0.90	12/5472 (0.2%)
1	D	0.42	1/3938 (0.0%)	0.91	13/5327 (0.2%)
All	All	0.41	1/16430 (0.0%)	0.91	49/22250 (0.2%)

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	D	6	MET	SD-CE	-7.30	1.61	1.79

The worst 5 of 49 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	316	ARG	N-CA-C	-8.31	99.56	110.53
1	B	435	SER	N-CA-C	8.24	119.71	109.57
1	C	39	THR	N-CA-C	7.98	121.74	108.73
1	A	316	ARG	N-CA-C	-7.77	98.98	110.48
1	A	435	SER	N-CA-C	7.56	120.37	109.48

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	4130	0	4067	121	0
1	B	4134	0	4070	107	0
1	C	3949	0	3889	116	0
1	D	3850	0	3784	144	0
2	A	5	0	1	0	0
2	B	5	0	1	2	0
2	C	5	0	1	1	0
2	D	5	0	1	2	0
3	A	6	0	8	0	0
3	B	6	0	8	2	0
3	D	6	0	8	2	0
4	A	1	0	0	0	0
4	B	1	0	0	0	0
4	C	1	0	0	0	0
4	D	1	0	0	0	0
5	A	89	0	0	4	0
5	B	82	0	0	1	0
5	C	73	0	0	1	0
5	D	60	0	0	2	0
All	All	16409	0	15838	435	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 14.

The worst 5 of 435 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:D:601:GLV:H1	3:D:602:GOL:H31	1.45	0.95
1:A:100:LEU:HA	1:D:6:MET:HE2	1.48	0.94
1:A:350:ASN:HD21	1:A:354:ARG:HE	1.16	0.91
1:A:101:ASP:H	1:D:6:MET:CE	1.87	0.87
1:C:150:GLN:HE22	1:C:168:ILE:H	1.22	0.86

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	522/560 (93%)	505 (97%)	14 (3%)	3 (1%)	21	18
1	B	523/560 (93%)	502 (96%)	21 (4%)	0	100	100
1	C	493/560 (88%)	464 (94%)	27 (6%)	2 (0%)	30	28
1	D	476/560 (85%)	451 (95%)	24 (5%)	1 (0%)	43	44
All	All	2014/2240 (90%)	1922 (95%)	86 (4%)	6 (0%)	36	36

5 of 6 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	123	ASP
1	C	216	GLY
1	D	219	GLY
1	A	293	GLN
1	A	294	SER

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	436/462 (94%)	427 (98%)	9 (2%)	47	54
1	B	436/462 (94%)	424 (97%)	12 (3%)	38	43
1	C	415/462 (90%)	406 (98%)	9 (2%)	45	53
1	D	403/462 (87%)	389 (96%)	14 (4%)	32	35
All	All	1690/1848 (92%)	1646 (97%)	44 (3%)	40	46

5 of 44 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	C	502	MET
1	D	185	LEU
1	C	524	GLN

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	D	83	TYR
1	D	268	ARG

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 71 such sidechains are listed below:

Mol	Chain	Res	Type
1	D	150	GLN
1	D	208	GLN
1	D	422	HIS
1	B	350	ASN
1	B	313	ASN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 11 ligands modelled in this entry, 4 are monoatomic - leaving 7 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
2	GLV	B	601	4	4,4,4	2.62	2 (50%)	3,4,4	2.44	2 (66%)

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
2	GLV	C	601	4	4,4,4	1.27	1 (25%)	3,4,4	2.13	2 (66%)
2	GLV	A	601	4	4,4,4	2.60	2 (50%)	3,4,4	2.40	2 (66%)
3	GOL	A	602	-	5,5,5	0.64	0	5,5,5	0.27	0
3	GOL	D	602	-	5,5,5	0.58	0	5,5,5	0.36	0
3	GOL	B	602	-	5,5,5	0.49	0	5,5,5	0.48	0
2	GLV	D	601	4	4,4,4	1.12	0	3,4,4	2.28	2 (66%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	GLV	B	601	4	-	0/0/2/2	-
2	GLV	C	601	4	-	0/0/2/2	-
2	GLV	A	601	4	-	0/0/2/2	-
3	GOL	A	602	-	-	0/4/4/4	-
3	GOL	D	602	-	-	2/4/4/4	-
3	GOL	B	602	-	-	2/4/4/4	-
2	GLV	D	601	4	-	0/0/2/2	-

All (5) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	B	601	GLV	O2-C2	3.97	1.32	1.23
2	A	601	GLV	O2-C2	3.80	1.32	1.23
2	A	601	GLV	O3-C2	-3.22	1.22	1.30
2	B	601	GLV	O3-C2	-3.03	1.22	1.30
2	C	601	GLV	C1-C2	2.12	1.48	1.44

The worst 5 of 8 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	601	GLV	O3-C2-C1	3.18	122.80	113.42
2	B	601	GLV	O3-C2-C1	3.09	122.56	113.42
2	D	601	GLV	O3-C2-C1	2.97	122.20	113.42
2	C	601	GLV	O3-C2-C1	2.74	121.50	113.42
2	B	601	GLV	O3-C2-O2	-2.54	117.52	122.70

There are no chirality outliers.

All (4) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
3	B	602	GOL	O1-C1-C2-C3
3	B	602	GOL	O1-C1-C2-O2
3	D	602	GOL	C1-C2-C3-O3
3	D	602	GOL	O2-C2-C3-O3

There are no ring outliers.

5 monomers are involved in 5 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	B	601	GLV	2	0
2	C	601	GLV	1	0
3	D	602	GOL	2	0
3	B	602	GOL	2	0
2	D	601	GLV	2	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	524/560 (93%)	0.25	25 (4%) 35 38	22, 32, 55, 90	0
1	B	525/560 (93%)	0.27	13 (2%) 58 61	22, 34, 52, 66	0
1	C	499/560 (89%)	0.44	44 (8%) 15 16	22, 34, 68, 78	0
1	D	486/560 (86%)	0.49	40 (8%) 17 18	23, 35, 68, 84	0
All	All	2034/2240 (90%)	0.36	122 (5%) 27 29	22, 34, 63, 90	0

The worst 5 of 122 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	C	339	ALA	7.5
1	C	291	ALA	6.1
1	D	339	ALA	5.9
1	A	301	LEU	5.2
1	D	287	LEU	5.2

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
3	GOL	D	602	6/6	0.76	0.17	34,34,36,36	0
3	GOL	B	602	6/6	0.78	0.16	29,32,35,37	0
3	GOL	A	602	6/6	0.78	0.16	32,33,34,35	0
2	GLV	C	601	5/5	0.82	0.16	47,49,50,52	0
4	MG	D	603	1/1	0.89	0.14	46,46,46,46	0
2	GLV	A	601	5/5	0.93	0.09	32,32,33,33	0
2	GLV	D	601	5/5	0.93	0.10	34,35,36,37	0
2	GLV	B	601	5/5	0.93	0.09	30,31,32,35	0
4	MG	C	602	1/1	0.94	0.13	41,41,41,41	0
4	MG	B	603	1/1	0.95	0.15	43,43,43,43	0
4	MG	A	603	1/1	0.96	0.15	34,34,34,34	0

6.5 Other polymers [i](#)

There are no such residues in this entry.