



Full wwPDB X-ray Structure Validation Report ⓘ

Apr 25, 2026 – 12:47 PM EDT

PDB ID : 6E9W / pdb_00006e9w
Title : Crystal structure of Rock1 with a pyridinylbenzamide based inhibitor
Authors : Judge, R.A.; Hobson, A.D.
Deposited on : 2018-08-01
Resolution : 2.96 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0
Mogul : 2022.3.0, CSD as543be (2022)
Xtriage (Phenix) : 2.0
EDS : 3.0
Buster-report : wwPDB partial adaption of 1.1.7 (2018)
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4 : 9.0.010 (Gargrove)
Density-Fitness : 1.0.12
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.96 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	1130 (2.98-2.94)
Clashscore	190562	1157 (2.98-2.94)
Ramachandran outliers	187476	1101 (2.98-2.94)
Sidechain outliers	187428	1101 (2.98-2.94)
RSRZ outliers	180081	1130 (2.98-2.94)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	443	66% 17% 5% 14%
1	B	443	62% 19% 5% 14%

2 Entry composition [i](#)

There are 4 unique types of molecules in this entry. The entry contains 6461 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Rho-associated protein kinase 1.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	383	Total	C	N	O	S	0	0	0
			3129	2005	523	580	21			
1	B	380	Total	C	N	O	S	0	0	0
			3102	1992	510	579	21			

There are 66 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	-27	MET	-	expression tag	UNP Q13464
A	-26	SER	-	expression tag	UNP Q13464
A	-25	TYR	-	expression tag	UNP Q13464
A	-24	TYR	-	expression tag	UNP Q13464
A	-23	HIS	-	expression tag	UNP Q13464
A	-22	HIS	-	expression tag	UNP Q13464
A	-21	HIS	-	expression tag	UNP Q13464
A	-20	HIS	-	expression tag	UNP Q13464
A	-19	HIS	-	expression tag	UNP Q13464
A	-18	HIS	-	expression tag	UNP Q13464
A	-17	ASP	-	expression tag	UNP Q13464
A	-16	TYR	-	expression tag	UNP Q13464
A	-15	ASP	-	expression tag	UNP Q13464
A	-14	ILE	-	expression tag	UNP Q13464
A	-13	PRO	-	expression tag	UNP Q13464
A	-12	THR	-	expression tag	UNP Q13464
A	-11	THR	-	expression tag	UNP Q13464
A	-10	GLU	-	expression tag	UNP Q13464
A	-9	ASN	-	expression tag	UNP Q13464
A	-8	LEU	-	expression tag	UNP Q13464
A	-7	TYR	-	expression tag	UNP Q13464
A	-6	PHE	-	expression tag	UNP Q13464
A	-5	GLN	-	expression tag	UNP Q13464
A	-4	GLY	-	expression tag	UNP Q13464
A	-3	ALA	-	expression tag	UNP Q13464


Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
A	-2	MET	-	expression tag	UNP Q13464
A	-1	ASP	-	expression tag	UNP Q13464
A	0	PRO	-	expression tag	UNP Q13464
A	1	LEU	-	expression tag	UNP Q13464
A	2	VAL	-	expression tag	UNP Q13464
A	3	PRO	-	expression tag	UNP Q13464
A	4	ARG	-	expression tag	UNP Q13464
A	5	GLY	-	expression tag	UNP Q13464
B	-27	MET	-	expression tag	UNP Q13464
B	-26	SER	-	expression tag	UNP Q13464
B	-25	TYR	-	expression tag	UNP Q13464
B	-24	TYR	-	expression tag	UNP Q13464
B	-23	HIS	-	expression tag	UNP Q13464
B	-22	HIS	-	expression tag	UNP Q13464
B	-21	HIS	-	expression tag	UNP Q13464
B	-20	HIS	-	expression tag	UNP Q13464
B	-19	HIS	-	expression tag	UNP Q13464
B	-18	HIS	-	expression tag	UNP Q13464
B	-17	ASP	-	expression tag	UNP Q13464
B	-16	TYR	-	expression tag	UNP Q13464
B	-15	ASP	-	expression tag	UNP Q13464
B	-14	ILE	-	expression tag	UNP Q13464
B	-13	PRO	-	expression tag	UNP Q13464
B	-12	THR	-	expression tag	UNP Q13464
B	-11	THR	-	expression tag	UNP Q13464
B	-10	GLU	-	expression tag	UNP Q13464
B	-9	ASN	-	expression tag	UNP Q13464
B	-8	LEU	-	expression tag	UNP Q13464
B	-7	TYR	-	expression tag	UNP Q13464
B	-6	PHE	-	expression tag	UNP Q13464
B	-5	GLN	-	expression tag	UNP Q13464
B	-4	GLY	-	expression tag	UNP Q13464
B	-3	ALA	-	expression tag	UNP Q13464
B	-2	MET	-	expression tag	UNP Q13464
B	-1	ASP	-	expression tag	UNP Q13464
B	0	PRO	-	expression tag	UNP Q13464
B	1	LEU	-	expression tag	UNP Q13464
B	2	VAL	-	expression tag	UNP Q13464
B	3	PRO	-	expression tag	UNP Q13464
B	4	ARG	-	expression tag	UNP Q13464
B	5	GLY	-	expression tag	UNP Q13464


- Molecule 2 is N-[(2,3-dihydro-1,4-benzodioxin-5-yl)methyl]-4-(pyridin-4-yl)benzamide (CCD

ID: J0P) (formula: C₂₁H₁₈N₂O₃).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
2	A	1	Total	C	N	O	0	0
			26	21	2	3		
2	B	1	Total	C	N	O	0	0
			26	21	2	3		

- Molecule 3 is SULFATE ION (CCD ID: SO4) (formula: O₄S).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	A	1	Total	O	S	0	0
			5	4	1		


- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	86	Total 86	O 86	0	0
4	B	87	Total 87	O 87	0	0


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: Rho-associated protein kinase 1



• Molecule 1: Rho-associated protein kinase 1



4 Data and refinement statistics

Property	Value	Source
Space group	P 31 2 1	Depositor
Cell constants a, b, c, α , β , γ	180.63Å 180.63Å 90.29Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	24.73 – 2.96 24.73 – 2.96	Depositor EDS
% Data completeness (in resolution range)	99.9 (24.73-2.96) 99.7 (24.73-2.96)	Depositor EDS
R_{merge}	0.08	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.31 (at 2.94Å)	Xtriage
Refinement program	BUSTER 2.11.7	Depositor
R, R_{free}	0.194 , 0.255 0.203 , 0.264	Depositor DCC
R_{free} test set	1802 reflections (5.09%)	wwPDB-VP
Wilson B-factor (Å ²)	81.8	Xtriage
Anisotropy	0.135	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.34 , 73.0	EDS
L-test for twinning ²	$\langle L \rangle = 0.47$, $\langle L^2 \rangle = 0.30$	Xtriage
Estimated twinning fraction	0.031 for -h,-k,l	Xtriage
F_o, F_c correlation	0.94	EDS
Total number of atoms	6461	wwPDB-VP
Average B, all atoms (Å ²)	81.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 2.80% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: J0P, SO4

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.86	0/3203	1.39	21/4326 (0.5%)
1	B	0.88	0/3176	1.40	24/4290 (0.6%)
All	All	0.87	0/6379	1.39	45/8616 (0.5%)

There are no bond length outliers.

All (45) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	35	ASP	CA-CB-CG	8.71	121.31	112.60
1	B	216	ASP	CA-CB-CG	7.78	120.38	112.60
1	A	65	LYS	CA-C-N	7.12	129.53	120.56
1	A	65	LYS	C-N-CA	7.12	129.53	120.56
1	A	316	PHE	CA-CB-CG	6.95	120.75	113.80
1	A	169	ASP	CA-C-N	-6.93	117.60	122.59
1	A	169	ASP	C-N-CA	-6.93	117.60	122.59
1	B	182	VAL	N-CA-CB	6.92	120.96	110.58
1	B	362	ASP	CA-CB-CG	6.81	119.41	112.60
1	A	207	ASP	CA-CB-CG	6.26	118.86	112.60
1	B	215	ALA	CA-C-N	-6.20	113.97	123.14
1	B	215	ALA	C-N-CA	-6.20	113.97	123.14
1	B	224	ASN	CA-C-N	6.08	128.75	120.54
1	B	224	ASN	C-N-CA	6.08	128.75	120.54
1	A	11	PHE	N-CA-C	-5.96	106.01	113.28
1	A	319	ASP	CA-C-N	5.91	129.69	120.82
1	A	319	ASP	C-N-CA	5.91	129.69	120.82
1	B	307	LYS	N-CA-C	-5.89	104.94	111.36
1	B	316	PHE	CA-CB-CG	5.88	119.69	113.80
1	B	307	LYS	CA-C-N	5.87	128.07	120.44
1	B	307	LYS	C-N-CA	5.87	128.07	120.44
1	B	75	ASP	N-CA-C	-5.81	105.85	113.17

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	26	ASN	CA-CB-CG	5.78	118.38	112.60
1	A	145	ASP	CA-CB-CG	5.78	118.38	112.60
1	A	215	ALA	CA-C-N	-5.76	114.93	123.11
1	A	215	ALA	C-N-CA	-5.76	114.93	123.11
1	A	231	CYS	CA-C-N	5.73	130.16	120.88
1	A	231	CYS	C-N-CA	5.73	130.16	120.88
1	A	224	ASN	CA-CB-CG	5.64	118.24	112.60
1	A	117	ASP	CA-CB-CG	5.64	118.24	112.60
1	B	162	VAL	N-CA-C	-5.58	105.28	110.53
1	B	145	ASP	CA-CB-CG	5.55	118.15	112.60
1	B	202	ASP	CA-CB-CG	5.41	118.00	112.60
1	B	126	ASP	CA-CB-CG	5.39	117.99	112.60
1	B	266	VAL	CA-C-N	5.36	127.47	120.28
1	B	266	VAL	C-N-CA	5.36	127.47	120.28
1	A	146	ASP	CA-CB-CG	5.36	117.96	112.60
1	A	148	TYR	N-CA-C	5.32	118.25	109.85
1	B	12	GLU	CB-CG-CD	5.28	121.57	112.60
1	B	331	GLU	N-CA-C	-5.20	105.69	111.36
1	A	142	ALA	N-CA-C	-5.17	100.61	109.24
1	B	57	SER	CA-C-N	5.15	127.60	120.29
1	B	57	SER	C-N-CA	5.15	127.60	120.29
1	A	356	VAL	N-CA-C	-5.07	102.53	107.55
1	A	216	ASP	CA-CB-CG	5.02	117.62	112.60

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3129	0	3058	33	0
1	B	3102	0	3012	36	0
2	A	26	0	0	0	0
2	B	26	0	0	0	0
3	A	5	0	0	0	0
4	A	86	0	0	1	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
4	B	87	0	0	2	0
All	All	6461	0	6070	67	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 5.

All (67) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:257:ARG:NH1	1:B:328:GLY:HA2	2.00	0.76
1:A:236:GLY:HA3	1:A:241:ILE:HD11	1.71	0.71
1:A:72:LYS:HG3	1:A:74:GLU:HG2	1.74	0.68
1:A:95:HIS:HB3	1:A:98:THR:HG22	1.77	0.66
1:B:204:MET:HB3	1:B:212:LEU:HD11	1.78	0.65
1:B:35:ASP:HB2	4:B:647:HOH:O	1.97	0.63
1:B:225:LYS:HA	4:B:627:HOH:O	2.00	0.62
1:A:223:MET:HB3	1:A:227:GLY:HA2	1.79	0.62
1:B:31:LEU:HD22	1:B:66:ILE:HG21	1.82	0.62
1:A:246:LEU:HD22	1:A:287:TYR:CE1	2.35	0.61
1:B:257:ARG:HH11	1:B:328:GLY:HA2	1.65	0.61
1:A:38:VAL:HG11	1:A:63:ILE:HG21	1.81	0.61
1:A:195:ILE:HG22	1:A:197:ARG:HG3	1.86	0.58
1:B:290:ILE:O	1:B:293:HIS:HB3	2.05	0.56
1:A:171:PRO:HG3	1:B:171:PRO:HB3	1.87	0.56
1:A:400:TYR:O	1:A:403:ARG:HD2	2.04	0.56
1:A:316:PHE:O	1:A:323:ARG:HD2	2.06	0.56
1:A:241:ILE:HG22	1:A:246:LEU:HG	1.88	0.55
1:A:242:SER:HB2	1:A:244:GLU:OE2	2.06	0.55
1:A:68:ASP:HA	1:A:72:LYS:HE2	1.88	0.55
1:A:246:LEU:HD22	1:A:287:TYR:HE1	1.73	0.54
1:A:141:TYR:CE1	1:A:403:ARG:NH2	2.76	0.54
1:B:128:MET:HB3	1:B:139:LEU:HB2	1.90	0.54
1:B:246:LEU:HD23	1:B:290:ILE:HG21	1.91	0.52
1:A:113:ILE:HG13	1:A:381:PHE:HZ	1.74	0.51
1:A:197:ARG:HD3	1:A:219:THR:O	2.10	0.51
1:B:127:ILE:HG21	1:B:189:ILE:HD11	1.94	0.50
1:A:49:ASN:HB3	1:A:52:ILE:HB	1.95	0.49
1:A:244:GLU:CD	1:A:323:ARG:HH22	2.19	0.49
1:B:199:VAL:HG13	1:B:263:SER:CB	2.43	0.49
1:B:232:ASP:HA	1:B:249:GLN:C	2.38	0.49
1:A:127:ILE:O	1:A:131:ALA:HB2	2.13	0.48

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:285:GLY:O	1:B:289:LYS:HG3	2.13	0.48
1:B:280:ALA:HB2	1:B:289:LYS:HD2	1.96	0.48
1:B:84:ARG:HH22	1:B:372:GLU:HB2	1.78	0.48
1:B:333:LYS:HA	1:B:345:TRP:CZ2	2.49	0.47
1:A:10:ARG:HH11	1:A:13:LYS:HE3	1.78	0.47
1:A:266:VAL:HG13	1:A:277:PRO:HD2	1.96	0.47
1:B:223:MET:HB3	1:B:227:GLY:HA2	1.96	0.47
1:A:237:THR:O	1:A:241:ILE:HG12	2.15	0.46
1:B:195:ILE:HD13	1:B:256:GLY:HA2	1.96	0.46
1:A:321:GLU:H	1:A:321:GLU:CD	2.24	0.46
1:B:43:PHE:HB2	1:B:46:LEU:HD12	1.97	0.46
1:B:139:LEU:HD11	1:B:142:ALA:HB2	1.97	0.46
1:B:196:HIS:O	1:B:197:ARG:HB2	2.16	0.45
1:A:346:GLU:H	1:A:346:GLU:CD	2.24	0.45
1:B:225:LYS:H	1:B:225:LYS:HD2	1.82	0.45
1:B:238:PRO:HA	1:B:241:ILE:HD12	1.98	0.45
1:B:20:ASP:HB3	1:B:23:SER:HB3	1.98	0.44
1:B:128:MET:CB	1:B:139:LEU:HB2	2.48	0.44
1:A:98:THR:HG23	1:A:100:LYS:HG3	2.00	0.43
1:B:59:TYR:HB3	1:B:63:ILE:HD12	2.00	0.43
1:A:154:GLU:HG3	1:A:156:MET:HE3	1.99	0.43
1:B:312:LEU:HD23	1:B:335:HIS:CD2	2.54	0.42
1:B:123:GLU:HB3	1:B:194:PHE:CE2	2.54	0.42
1:B:70:ARG:HG2	1:B:400:TYR:OH	2.19	0.42
1:A:63:ILE:O	1:A:67:ARG:HG3	2.20	0.41
1:B:127:ILE:O	1:B:131:ALA:HB2	2.20	0.41
1:A:35:ASP:OD1	1:A:67:ARG:HD3	2.20	0.41
1:B:316:PHE:O	1:B:323:ARG:HD2	2.20	0.41
1:A:404:ARG:HD2	4:A:620:HOH:O	2.20	0.41
1:B:343:TRP:HB3	1:B:351:THR:HG21	2.02	0.41
1:A:139:LEU:HD21	1:A:142:ALA:HB2	2.02	0.41
1:B:241:ILE:HG21	1:B:246:LEU:HD13	2.03	0.41
1:B:315:ALA:HB1	1:B:324:LEU:HB2	2.03	0.41
1:A:173:LYS:HD2	1:B:169:ASP:OD2	2.21	0.40
1:A:204:MET:HB3	1:A:212:LEU:HD11	2.02	0.40

There are no symmetry-related clashes.

5.3 Torsion angles

5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	377/443 (85%)	347 (92%)	30 (8%)	0	100	100
1	B	372/443 (84%)	343 (92%)	24 (6%)	5 (1%)	9	26
All	All	749/886 (84%)	690 (92%)	54 (7%)	5 (1%)	18	40

All (5) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	B	257	ARG
1	B	258	GLU
1	B	197	ARG
1	B	74	GLU
1	B	345	TRP

5.3.2 Protein sidechains

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	343/395 (87%)	300 (88%)	43 (12%)	4	13
1	B	339/395 (86%)	294 (87%)	45 (13%)	4	12
All	All	682/790 (86%)	594 (87%)	88 (13%)	4	13

All (88) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	9	THR

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	A	14	MET
1	A	31	LEU
1	A	47	ARG
1	A	50	LYS
1	A	60	LYS
1	A	67	ARG
1	A	68	ASP
1	A	72	LYS
1	A	74	GLU
1	A	78	VAL
1	A	82	ILE
1	A	98	THR
1	A	111	GLU
1	A	117	ASP
1	A	162	VAL
1	A	191	SER
1	A	197	ARG
1	A	221	MET
1	A	225	LYS
1	A	232	ASP
1	A	244	GLU
1	A	247	LYS
1	A	268	LEU
1	A	283	LEU
1	A	294	LYS
1	A	297	LEU
1	A	308	GLU
1	A	311	ASN
1	A	312	LEU
1	A	317	LEU
1	A	320	ARG
1	A	321	GLU
1	A	323	ARG
1	A	342	GLN
1	A	346	GLU
1	A	347	THR
1	A	352	VAL
1	A	358	ASP
1	A	359	LEU
1	A	383	ILE
1	A	403	ARG
1	A	404	ARG

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	B	8	GLU
1	B	23	SER
1	B	24	GLU
1	B	48	LYS
1	B	56	LEU
1	B	61	ASP
1	B	62	THR
1	B	68	ASP
1	B	70	ARG
1	B	89	GLU
1	B	98	THR
1	B	139	LEU
1	B	147	ARG
1	B	161	LEU
1	B	182	VAL
1	B	189	ILE
1	B	199	VAL
1	B	212	LEU
1	B	214	LEU
1	B	219	THR
1	B	221	MET
1	B	225	LYS
1	B	228	MET
1	B	246	LEU
1	B	249	GLN
1	B	258	GLU
1	B	268	LEU
1	B	281	ASP
1	B	291	MET
1	B	297	LEU
1	B	301	ASP
1	B	304	ASP
1	B	308	GLU
1	B	312	LEU
1	B	320	ARG
1	B	321	GLU
1	B	323	ARG
1	B	324	LEU
1	B	336	LEU
1	B	349	ARG
1	B	351	THR
1	B	358	ASP

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	B	366	SER
1	B	383	ILE
1	B	401	SER

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (8) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	167	ASN
1	A	196	HIS
1	A	327	ASN
1	B	138	GLN
1	B	196	HIS
1	B	203	ASN
1	B	293	HIS
1	B	342	GLN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

3 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
2	J0P	B	501	-	29,29,29	1.23	2 (6%)	39,39,39	0.75	0
2	J0P	A	501	-	29,29,29	1.16	2 (6%)	39,39,39	0.97	1 (2%)
3	SO4	A	502	-	4,4,4	0.33	0	6,6,6	0.31	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	J0P	B	501	-	-	0/13/20/20	0/4/4/4
2	J0P	A	501	-	-	0/13/20/20	0/4/4/4

All (4) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	A	501	J0P	C3-C2	3.72	1.45	1.39
2	B	501	J0P	C-N	3.20	1.40	1.33
2	B	501	J0P	C3-C2	2.70	1.44	1.39
2	A	501	J0P	C-N	2.34	1.38	1.33

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	501	J0P	O2-C6-C7	-2.79	119.67	121.78

There are no chirality outliers.


There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier.

The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.



5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues ⓘ

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	383/443 (86%)	-0.09	4 (1%) 79 74	53, 77, 110, 146	0
1	B	380/443 (85%)	0.01	2 (0%) 87 83	57, 79, 114, 135	0
All	All	763/886 (86%)	-0.04	6 (0%) 82 78	53, 78, 111, 146	0

All (6) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	119	ALA	3.1
1	A	9	THR	3.1
1	A	11	PHE	3.0
1	A	287	TYR	2.7
1	B	39	TYR	2.2
1	A	254	TYR	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)




In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
3	SO4	A	502	5/5	0.94	0.10	112,113,113,114	0
2	J0P	B	501	26/26	0.96	0.09	49,75,79,80	0
2	J0P	A	501	26/26	0.96	0.08	50,60,67,67	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.




Electron density around J0P B 501:

2mF_o-DF_c (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
 and green (positive)



Electron density around J0P A 501:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



6.5 Other polymers [i](#)

There are no such residues in this entry.