



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 5, 2026 – 02:42 PM UTC

PDB ID : 3EA4 / pdb_00003ea4
Title : Arabidopsis thaliana acetoxyacid synthase in complex with
monosulfuron-ester
Authors : Guddat, L.W.; Wang, J.-G.; Li, Z.-M.
Deposited on : 2008-08-24
Resolution : 2.80 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0
Mogul : 2022.3.0, CSD as543be (2022)
Xtrriage (Phenix) : 2.0
EDS : 3.0
Buster-report : wwPDB partial adaption of 1.1.7 (2018)
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4 : 9.0.010 (Gargrove)
Density-Fitness : 1.0.12
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.80 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	3866 (2.80-2.80)
Clashscore	190562	4276 (2.80-2.80)
Ramachandran outliers	187476	4196 (2.80-2.80)
Sidechain outliers	187428	4198 (2.80-2.80)
RSRZ outliers	180081	3869 (2.80-2.80)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	584	

2 Entry composition [i](#)

There are 7 unique types of molecules in this entry. The entry contains 4707 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called Acetolactate synthase, chloroplastic.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	582	4479	2843	769	842	25	0	3	0

There is a discrepancy between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
A	330	THR	SER	conflict	UNP P17597

- Molecule 2 is methyl 2-[[[(4-methylpyrimidin-2-yl)carbamoyl]sulfamoyl]benzoate (CCD ID: 2SM) (formula: C₁₄H₁₄N₄O₅S).




Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
			Total	C	N	O	S		
2	A	1	24	14	4	5	1	0	0

- Molecule 3 is FLAVIN-ADENINE DINUCLEOTIDE-N5-ISOBUTYL KETONE (CCD ID: FAB) (formula: C₃₁H₃₉N₉O₁₆P₂).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
			Total	C	N	O	P		
3	A	1	53	27	9	15	2	0	0


- Molecule 4 is 2-[N-CYCLOHEXYLAMINO]ETHANE SULFONIC ACID (CCD ID: NHE) (formula: C₈H₁₇NO₃S).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
			Total	C	N	O	S		
4	A	1	13	8	1	3	1	0	0

- Molecule 5 is 2-[(2E)-3-[(4-AMINO-2-METHYLPYRIMIDIN-5-YL)METHYL]-2-(1-H YDROXYETHYLIDENE)-4-METHYL-2,3-DIHYDRO-1,3-THIAZOL-5-YL]ETHYL

TRIHYDROGEN DIPHOSPHATE (CCD ID: TDM) (formula: C₁₄H₂₂N₄O₈P₂S).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	
			Total	C	N	O	P			S
5	A	1	29	14	4	8	2	1	0	0

- Molecule 6 is MAGNESIUM ION (CCD ID: MG) (formula: Mg).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
			Total	Mg		
6	A	1	1	1	0	0


- Molecule 7 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
			Total	O		
7	A	108	108	108	0	0

3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Acetolactate synthase, chloroplactic



4 Data and refinement statistics

Property	Value	Source
Space group	P 64 2 2	Depositor
Cell constants a, b, c, α , β , γ	178.47Å 178.47Å 184.95Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	58.42 – 2.80 58.42 – 2.80	Depositor EDS
% Data completeness (in resolution range)	98.5 (58.42-2.80) 98.6 (58.42-2.80)	Depositor EDS
R_{merge}	0.07	Depositor
R_{sym}	0.07	Depositor
$\langle I/\sigma(I) \rangle$ ¹	3.51 (at 2.81Å)	Xtrriage
Refinement program	REFMAC 5.2.0019	Depositor
R, R_{free}	0.199 , 0.213 0.192 , 0.206	Depositor DCC
R_{free} test set	2153 reflections (4.96%)	wwPDB-VP
Wilson B-factor (Å ²)	58.1	Xtrriage
Anisotropy	0.003	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.35 , 35.0	EDS
L-test for twinning ²	$\langle L \rangle = 0.52$, $\langle L^2 \rangle = 0.36$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
F_o, F_c correlation	0.93	EDS
Total number of atoms	4707	wwPDB-VP
Average B, all atoms (Å ²)	52.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 2.05% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: TDM, 2SM, CSD, FAB, MG, NHE

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.47	0/4570	0.79	0/6204

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	4479	0	4471	40	0
2	A	24	0	14	1	0
3	A	53	0	31	0	0
4	A	13	0	16	1	0
5	A	29	0	19	4	0
6	A	1	0	0	0	0
7	A	108	0	0	2	0
All	All	4707	0	4551	44	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 5.

All (44) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:644:GLN:HG3	1:A:667:ARG:HG3	1.57	0.84
1:A:145:GLN:HE21	1:A:544:ASN:HD21	1.30	0.80
1:A:117:ALA:H	1:A:138:ASN:HD21	1.30	0.80
1:A:332:LEU:HA	1:A:348:MET:HE2	1.64	0.78
1:A:566:GLN:HE22	1:A:598:PHE:H	1.33	0.76
1:A:143:HIS:HD2	1:A:145:GLN:H	1.33	0.72
1:A:137:ARG:HD2	7:A:702:HOH:O	1.97	0.63
1:A:394:ILE:HG12	1:A:411:VAL:HB	1.81	0.62
1:A:312:ASN:HD21	1:A:502:ARG:HH21	1.47	0.61
2:A:695:2SM:N8	2:A:695:2SM:O25	2.34	0.60
1:A:460:GLU:HB3	1:A:617:LYS:HE2	1.84	0.60
1:A:143:HIS:CD2	1:A:145:GLN:H	2.17	0.59
1:A:614:ARG:HD2	1:A:640:ILE:HD11	1.86	0.57
1:A:266:ASN:HB3	1:A:269:GLN:HE21	1.69	0.57
1:A:117:ALA:N	1:A:138:ASN:HD21	2.01	0.55
1:A:656:THR:HG22	1:A:657:PHE:H	1.74	0.53
1:A:90:ARG:NH2	7:A:779:HOH:O	2.43	0.51
1:A:656:THR:HG22	1:A:657:PHE:N	2.25	0.51
5:A:698:TDM:C27	5:A:698:TDM:H4'1	2.24	0.50
1:A:562:LEU:HD21	1:A:601:MET:HG3	1.93	0.50
1:A:92:ALA:HB3	1:A:95:GLN:HB2	1.93	0.49
1:A:370:PHE:HB3	1:A:415:VAL:HG21	1.94	0.49
1:A:350:GLY:HA2	1:A:378:VAL:HA	1.95	0.48
1:A:139:VAL:HG13	1:A:554:GLU:HG3	1.95	0.48
5:A:698:TDM:H7'1	5:A:698:TDM:O28	2.14	0.48
1:A:144:GLU:HG2	1:A:174:ASN:HB2	1.96	0.48
1:A:568:LEU:HD22	5:A:698:TDM:HM43	1.97	0.47
1:A:277:MET:HA	1:A:280:MET:HG3	1.97	0.47
1:A:485:VAL:HG21	1:A:511:GLY:C	2.40	0.46
1:A:570:MET:HB3	5:A:698:TDM:S1	2.55	0.46
1:A:144:GLU:HG2	1:A:174:ASN:CB	2.47	0.45
1:A:552:ARG:HD3	1:A:552:ARG:O	2.16	0.45
1:A:114:THR:HG21	1:A:525:VAL:HG11	1.99	0.44
4:A:697:NHE:H6'2	4:A:697:NHE:HC11	1.63	0.43
1:A:482:SER:HA	1:A:505:LEU:O	2.19	0.42
1:A:369:ALA:HB1	1:A:372:VAL:CG2	2.50	0.42
1:A:463:PRO:HA	1:A:464:PRO:HD2	1.92	0.41
1:A:332:LEU:CA	1:A:348:MET:HE2	2.45	0.41
1:A:198:ARG:HD3	1:A:258:ILE:HD11	2.03	0.41
1:A:351[B]:MET:HE1	1:A:380:GLY:HA2	2.03	0.41
1:A:351[B]:MET:HE3	1:A:653:SER:HA	2.03	0.41
1:A:272:ARG:C	1:A:274:PRO:HD3	2.46	0.41

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:401:ILE:HG21	1:A:410:SER:HB3	2.02	0.40
1:A:501:PRO:O	1:A:502:ARG:HB2	2.22	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	582/584 (100%)	575 (99%)	7 (1%)	0	100 100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	480/480 (100%)	468 (98%)	12 (2%)	42 76

All (12) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	130	LEU
1	A	137	ARG
1	A	144	GLU
1	A	148	VAL

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	A	344	LEU
1	A	375	ASP
1	A	377	ARG
1	A	378	VAL
1	A	390	LYS
1	A	456	LYS
1	A	473	GLU
1	A	614	ARG

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (10) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	138	ASN
1	A	143	HIS
1	A	145	GLN
1	A	174	ASN
1	A	261	GLN
1	A	269	GLN
1	A	312	ASN
1	A	420	GLN
1	A	494	GLN
1	A	566	GLN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

1 non-standard protein/DNA/RNA residue is modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
1	CSD	A	340	1	4,7,8	2.23	1 (25%)	1,8,10	1.84	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
1	CSD	A	340	1	-	1/2/6/8	-

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	340	CSD	OD1-SG	-4.00	1.44	1.47

There are no bond angle outliers.

There are no chirality outliers.

All (1) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
1	A	340	CSD	CA-CB-SG-OD1

There are no ring outliers.

No monomer is involved in short contacts.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 5 ligands modelled in this entry, 1 is monoatomic - leaving 4 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
5	TDM	A	698	6	28,30,30	2.84	7 (25%)	37,45,45	2.16	9 (24%)
2	2SM	A	695	-	25,25,25	2.38	7 (28%)	34,35,35	3.38	11 (32%)
4	NHE	A	697	-	13,13,13	1.57	1 (7%)	16,17,17	1.02	1 (6%)
3	FAB	A	696	-	58,58,63	2.00	9 (15%)	85,89,97	1.91	14 (16%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
5	TDM	A	698	6	-	3/17/21/21	0/2/2/2
2	2SM	A	695	-	-	9/21/21/21	0/2/2/2
4	NHE	A	697	-	-	6/7/15/15	0/1/1/1
3	FAB	A	696	-	-	8/34/50/58	0/6/6/6

All (24) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
5	A	698	TDM	O28-C27	9.43	1.56	1.31
5	A	698	TDM	C2-N3	8.39	1.54	1.39
3	A	696	FAB	C2A-N3A	8.07	1.48	1.33
3	A	696	FAB	C2A-N1A	7.98	1.48	1.33
2	A	695	2SM	C14-C13	6.52	1.54	1.39
2	A	695	2SM	O17-S7	5.55	1.50	1.43
3	A	696	FAB	C4X-N5	4.93	1.41	1.30
2	A	695	2SM	C11-N16	4.34	1.40	1.34
4	A	697	NHE	C2-S	4.08	1.83	1.77
5	A	698	TDM	PB-O1B	3.82	1.62	1.50
3	A	696	FAB	C10-N1	3.33	1.40	1.33
2	A	695	2SM	C11-N10	-3.27	1.34	1.38
5	A	698	TDM	C5-C4	3.21	1.41	1.35
2	A	695	2SM	C15-N16	2.96	1.40	1.34
3	A	696	FAB	C5A-N7A	-2.94	1.33	1.39
5	A	698	TDM	CM4-C4	2.91	1.54	1.49
2	A	695	2SM	C13-N12	2.86	1.40	1.34
5	A	698	TDM	C7'-C5'	2.63	1.56	1.51
3	A	696	FAB	C4A-N3A	2.61	1.39	1.34
3	A	696	FAB	P-O3P	2.48	1.62	1.59
3	A	696	FAB	C4A-N9A	-2.36	1.32	1.37

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
5	A	698	TDM	C29-C27	2.19	1.53	1.48
2	A	695	2SM	C9-N8	-2.11	1.34	1.39
3	A	696	FAB	C8A-N9A	-2.10	1.34	1.37

All (35) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	A	696	FAB	N3A-C2A-N1A	-10.87	112.13	128.58
5	A	698	TDM	C5-S1-C2	8.89	95.93	90.80
2	A	695	2SM	O17-S7-O18	-8.60	109.07	119.52
2	A	695	2SM	C15-N16-C11	7.94	122.06	115.42
2	A	695	2SM	N16-C11-N12	-7.66	119.01	126.42
2	A	695	2SM	C5-S7-N8	7.27	115.00	105.96
2	A	695	2SM	C15-C14-C13	-5.77	113.15	117.60
2	A	695	2SM	O23-C22-C4	5.65	121.69	112.31
2	A	695	2SM	C24-O23-C22	-4.38	107.39	115.81
3	A	696	FAB	C2A-N3A-C4A	4.25	122.22	111.83
2	A	695	2SM	O23-C22-O25	-3.89	115.89	123.46
3	A	696	FAB	C5A-C4A-N3A	-3.62	121.73	126.72
2	A	695	2SM	C11-N10-C9	-3.42	126.82	130.34
3	A	696	FAB	C4-N3-C2	-3.37	119.65	125.64
5	A	698	TDM	C7'-C5'-C4'	3.37	125.97	122.56
3	A	696	FAB	N9A-C8A-N7A	-3.23	109.35	113.94
5	A	698	TDM	CM2-C2'-N1'	3.11	120.51	117.20
3	A	696	FAB	C4A-C5A-N7A	-3.08	107.06	110.58
3	A	696	FAB	C10-C4X-N5	-3.06	118.55	124.81
5	A	698	TDM	N1'-C2'-N3'	-3.04	120.47	125.53
5	A	698	TDM	C5'-C7'-N3	-2.96	109.56	113.14
3	A	696	FAB	C9A-C5X-N5	-2.86	119.42	122.45
5	A	698	TDM	C29-C27-C2	-2.80	120.85	124.73
3	A	696	FAB	C5A-N7A-C8A	2.78	107.81	103.45
5	A	698	TDM	C6'-N1'-C2'	2.76	120.61	116.07
3	A	696	FAB	C4X-C10-N10	2.72	120.37	116.48
5	A	698	TDM	C7'-C5'-C6'	-2.65	117.51	121.30
3	A	696	FAB	C4X-C4-N3	2.63	119.95	113.25
3	A	696	FAB	C2A-N1A-C6A	2.61	123.02	118.73
4	A	697	NHE	O2-S-C2	2.56	110.59	106.73
3	A	696	FAB	C4-C4X-N5	2.49	121.65	118.21
5	A	698	TDM	O28-C27-C29	2.44	119.44	114.60
2	A	695	2SM	C11-N12-C13	2.34	120.52	115.98
2	A	695	2SM	N10-C11-N16	2.30	122.56	116.48
3	A	696	FAB	C4A-N9A-C8A	2.05	107.89	105.74

There are no chirality outliers.

All (26) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	A	695	2SM	C4-C22-O23-C24
3	A	696	FAB	C3D-C4D-C5D-O5'
3	A	696	FAB	O4'-C4D-C5D-O5'
4	A	697	NHE	C6'-C1'-N-C1
4	A	697	NHE	C1-C2-S-O2
2	A	695	2SM	O25-C22-O23-C24
2	A	695	2SM	C4-C5-S7-O18
4	A	697	NHE	C1-C2-S-O3
2	A	695	2SM	C6-C5-S7-O18
4	A	697	NHE	C2-C1-N-C1'
4	A	697	NHE	C2'-C1'-N-C1
2	A	695	2SM	C4-C5-S7-N8
4	A	697	NHE	C1-C2-S-O1
5	A	698	TDM	C5-C6-C7-O7
2	A	695	2SM	C6-C5-S7-N8
5	A	698	TDM	C7-O7-PA-O3A
3	A	696	FAB	C2B-C1B-N9A-C8A
3	A	696	FAB	O2'-C2D-C3D-C4D
3	A	696	FAB	C1'-C2D-C3D-C4D
5	A	698	TDM	PA-O3A-PB-O2B
3	A	696	FAB	C2D-C3D-C4D-O4'
3	A	696	FAB	O3D-C3D-C4D-C5D
2	A	695	2SM	N8-C9-N10-C11
3	A	696	FAB	N10-C1'-C2D-O2'
2	A	695	2SM	C9-N8-S7-O17
2	A	695	2SM	O20-C9-N10-C11


There are no ring outliers.


3 monomers are involved in 6 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
5	A	698	TDM	4	0
2	A	695	2SM	1	0
4	A	697	NHE	1	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier.

Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.





5.7 Other polymers [\(i\)](#)

There are no such residues in this entry.

5.8 Polymer linkage issues

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	581/584 (99%)	0.59	37 (6%) 25 18	29, 50, 66, 87	3 (0%)

All (37) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	668	ILE	11.7
1	A	291[A]	GLN	8.1
1	A	595	ASP	5.8
1	A	594	GLU	5.3
1	A	408	HIS	3.6
1	A	431	GLU	3.6
1	A	430	ALA	3.5
1	A	656	THR	3.5
1	A	268	GLU	3.5
1	A	351[A]	MET	3.4
1	A	424	LYS	3.3
1	A	617	LYS	3.0
1	A	383[A]	GLU	2.8
1	A	658	ASN	2.8
1	A	287	SER	2.8
1	A	644	GLN	2.7
1	A	388	ARG	2.6
1	A	428	ASN	2.5
1	A	330	THR	2.5
1	A	623	GLU	2.4
1	A	555	ASN	2.4
1	A	653	SER	2.3
1	A	643	HIS	2.3
1	A	229	GLU	2.3
1	A	279	ARG	2.3
1	A	542	ILE	2.2
1	A	364	SER	2.2

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	343	GLU	2.2
1	A	94	ASP	2.1
1	A	580	LYS	2.1
1	A	645	GLU	2.1
1	A	269	GLN	2.1
1	A	663	GLU	2.1
1	A	432	GLU	2.1
1	A	214	VAL	2.0
1	A	88	ILE	2.0
1	A	596	GLU	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
1	CSD	A	340	8/9	0.89	0.16	64,65,67,67	0

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.


6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
5	TDM	A	698	29/29	0.91	0.19	52,69,70,70	0
2	2SM	A	695	24/24	0.92	0.16	62,62,63,64	0
4	NHE	A	697	13/13	0.93	0.14	56,57,58,58	0
6	MG	A	699	1/1	0.95	0.07	55,55,55,55	0
3	FAB	A	696	53/58	0.97	0.09	34,39,43,43	0




The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers

as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.






Electron density around 2SM A 695:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)

**Electron density around FAB A 696:**

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



6.5 Other polymers [i](#)

There are no such residues in this entry.