



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 7, 2026 – 05:20 AM UTC

PDB ID : 5EAB / pdb_00005eab
Title : Saccharomyces cerevisiae CYP51 complexed with the plant pathogen inhibitor S-tebuconazole
Authors : Tyndall, J.D.A.; Sabherwal, M.; Sagatova, A.A.; Keniya, M.V.; Wilson, R.K.; Woods, M.V.; Monk, B.C.
Deposited on : 2015-10-16
Resolution : 2.60 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Buster-report	:	wwPDB partial adaption of 1.1.7 (2018)
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.60 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	4008 (2.60-2.60)
Clashscore	190562	4347 (2.60-2.60)
Ramachandran outliers	187476	4277 (2.60-2.60)
Sidechain outliers	187428	4277 (2.60-2.60)
RSRZ outliers	180081	4008 (2.60-2.60)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	539	

2 Entry composition

There are 3 unique types of molecules in this entry. The entry contains 4306 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Lanosterol 14-alpha demethylase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	525	4242	2748	714	759	21	0	0	0

There are 9 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
A	531	GLY	-	expression tag	UNP A6ZSR0
A	532	GLY	-	expression tag	UNP A6ZSR0
A	533	ARG	-	expression tag	UNP A6ZSR0
A	534	HIS	-	expression tag	UNP A6ZSR0
A	535	HIS	-	expression tag	UNP A6ZSR0
A	536	HIS	-	expression tag	UNP A6ZSR0
A	537	HIS	-	expression tag	UNP A6ZSR0
A	538	HIS	-	expression tag	UNP A6ZSR0
A	539	HIS	-	expression tag	UNP A6ZSR0

- Molecule 2 is PROTOPORPHYRIN IX CONTAINING FE (CCD ID: HEM) (formula: $C_{34}H_{32}FeN_4O_4$).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
2	A	1	Total	C	Fe	N	O	0	0
			43	34	1	4	4		

- Molecule 3 is (3 {S})-1-(4-chlorophenyl)-4,4-dimethyl-3-(1,2,4-triazol-1-ylmethyl)pentan-3-ol (CCD ID: TBX) (formula: $C_{16}H_{22}ClN_3O$).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
3	A	1	Total	C	Cl	N	O	0	0
			21	16	1	3	1		

- Molecule 1: Lanosterol 14- α demethylase



4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	80.70Å 67.23Å 81.63Å 90.00° 100.50° 90.00°	Depositor
Resolution (Å)	29.94 – 2.60 29.94 – 2.60	Depositor EDS
% Data completeness (in resolution range)	99.2 (29.94-2.60) 99.3 (29.94-2.60)	Depositor EDS
R_{merge}	0.08	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.40 (at 2.61Å)	Xtriage
Refinement program	PHENIX 1.9_1692	Depositor
R, R_{free}	0.208 , 0.263 0.220 , 0.267	Depositor DCC
R_{free} test set	1337 reflections (2.06%)	wwPDB-VP
Wilson B-factor (Å ²)	60.0	Xtriage
Anisotropy	0.622	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.33 , 56.0	EDS
L-test for twinning ²	$\langle L \rangle = 0.50$, $\langle L^2 \rangle = 0.33$	Xtriage
Estimated twinning fraction	0.017 for l,-k,h	Xtriage
F_o, F_c correlation	0.95	EDS
Total number of atoms	4306	wwPDB-VP
Average B, all atoms (Å ²)	81.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 5.44% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality ⓘ

5.1 Standard geometry ⓘ

Bond lengths and bond angles in the following residue types are not validated in this section: HEM, TBX

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.56	1/4359 (0.0%)	0.90	10/5906 (0.2%)

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	167	VAL	CA-CB	6.71	1.57	1.54

All (10) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	9	GLY	N-CA-C	-9.60	104.11	114.67
1	A	167	VAL	CA-C-N	-6.41	112.40	119.19
1	A	167	VAL	C-N-CA	-6.41	112.40	119.19
1	A	278	ILE	N-CA-C	5.87	112.50	106.21
1	A	378	HIS	CA-C-N	5.79	125.64	119.28
1	A	378	HIS	C-N-CA	5.79	125.64	119.28
1	A	105	GLY	CA-C-N	5.32	125.14	119.28
1	A	105	GLY	C-N-CA	5.32	125.14	119.28
1	A	525	ASN	CA-C-N	5.21	125.26	119.32
1	A	525	ASN	C-N-CA	5.21	125.26	119.32

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within

the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	4242	0	4228	56	0
2	A	43	0	30	4	0
3	A	21	0	22	2	0
All	All	4306	0	4280	59	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 7.

All (59) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:371:LYS:HE2	1:A:431:TRP:CE3	2.28	0.68
1:A:33:LEU:HA	1:A:36:ILE:HG22	1.75	0.68
1:A:183:ASN:HB3	1:A:194:ILE:HG22	1.78	0.64
1:A:9:GLY:HA2	1:A:12:LEU:HD13	1.81	0.63
1:A:121:SER:OG	1:A:123:GLU:HG3	2.02	0.60
1:A:184:PHE:O	1:A:189:ARG:HD3	2.01	0.59
1:A:13:GLU:O	1:A:17:ILE:HG13	2.03	0.57
3:A:602:TBX:N26	3:A:602:TBX:H6	2.21	0.56
1:A:12:LEU:HA	1:A:15:VAL:HG12	1.90	0.54
1:A:368:GLN:HG2	1:A:431:TRP:HB2	1.90	0.53
1:A:465:GLY:HA2	1:A:469:ARG:HB2	1.90	0.52
1:A:31:ILE:HD13	1:A:34:ILE:HD12	1.92	0.52
1:A:52:ARG:NH2	1:A:54:ASP:OD2	2.42	0.52
1:A:30:ARG:O	1:A:34:ILE:HG13	2.08	0.52
1:A:268:LEU:HD23	1:A:284:ILE:HD13	1.91	0.51
1:A:25:LEU:HD13	1:A:29:GLN:HB2	1.92	0.51
2:A:601:HEM:HBB2	2:A:601:HEM:HMB2	1.93	0.51
1:A:273:ARG:NH2	1:A:301:ASP:OD2	2.44	0.50
1:A:317:HIS:HB3	1:A:510:VAL:HG11	1.93	0.50
1:A:496:PRO:HD2	1:A:499:LYS:HB2	1.94	0.49
1:A:198:VAL:O	1:A:202:GLU:HG3	2.12	0.49
1:A:163:PHE:HB3	1:A:357:TYR:CD1	2.47	0.49
1:A:229:TYR:OH	1:A:265:TYR:OH	2.29	0.49
1:A:8:VAL:HG12	1:A:10:GLU:HB3	1.95	0.48
1:A:233:ASP:OD1	1:A:317:HIS:NE2	2.45	0.48
1:A:437:SER:HB2	1:A:473:GLU:OE1	2.13	0.48
1:A:322:THR:O	1:A:326:ILE:HG13	2.13	0.47
1:A:331:ALA:O	1:A:494:HIS:HA	2.14	0.47
1:A:243:PHE:O	1:A:251:TYR:OH	2.25	0.47

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:37:ILE:HB	1:A:38:PRO:HD3	1.97	0.47
1:A:229:TYR:HH	1:A:265:TYR:HH	1.56	0.47
1:A:470:CYS:HA	2:A:601:HEM:C4D	2.50	0.46
1:A:10:GLU:O	1:A:14:TYR:HD2	1.98	0.46
1:A:248:LEU:O	1:A:252:ARG:NH2	2.49	0.46
1:A:255:ASP:O	1:A:259:LYS:HG2	2.16	0.46
1:A:61:TYR:CD2	1:A:67:GLY:HA2	2.51	0.45
1:A:371:LYS:NZ	1:A:436:ALA:O	2.49	0.45
1:A:35:ILE:O	1:A:38:PRO:HD2	2.16	0.45
1:A:210:ARG:HA	1:A:218:ARG:HB2	1.98	0.45
1:A:199:THR:O	1:A:203:MET:HG3	2.17	0.45
1:A:386:LYS:HG3	1:A:405:HIS:CE1	2.52	0.45
1:A:314:GLY:HA3	3:A:602:TBX:H3	2.00	0.44
1:A:146:ARG:HD2	1:A:293:TYR:CE1	2.52	0.44
1:A:27:LEU:HA	1:A:30:ARG:HG2	1.98	0.44
1:A:243:PHE:HB3	1:A:246:LEU:HD23	1.99	0.43
1:A:235:GLY:O	1:A:240:ASN:ND2	2.51	0.43
1:A:22:PHE:O	1:A:25:LEU:HG	2.19	0.42
1:A:485:SER:O	1:A:489:ARG:HD3	2.20	0.42
1:A:213:LEU:HD21	1:A:312:LEU:HD13	2.02	0.41
2:A:601:HEM:HHA	2:A:601:HEM:HBD2	2.02	0.41
1:A:416:ARG:HA	1:A:416:ARG:HD2	1.90	0.41
1:A:417:ASP:HA	1:A:420:PHE:O	2.20	0.41
1:A:470:CYS:HB2	2:A:601:HEM:NA	2.35	0.41
1:A:221:LEU:HD23	1:A:221:LEU:HA	1.91	0.41
1:A:40:ILE:O	1:A:44:VAL:HG23	2.21	0.41
1:A:174:VAL:HG13	1:A:203:MET:HE1	2.02	0.41
1:A:523:LYS:NZ	1:A:528:GLN:O	2.50	0.40
1:A:378:HIS:HA	1:A:379:PRO:HD3	1.93	0.40
1:A:379:PRO:O	1:A:510:VAL:HA	2.21	0.40

There are no symmetry-related clashes.

5.3 Torsion angles

5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	523/539 (97%)	491 (94%)	29 (6%)	3 (1%)	21	42

All (3) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	139	ILE
1	A	434	ASP
1	A	441	VAL

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	460/473 (97%)	435 (95%)	25 (5%)	20	42

All (25) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	8	VAL
1	A	29	GLN
1	A	42	ASN
1	A	98	ARG
1	A	123	GLU
1	A	136	LYS
1	A	139	ILE
1	A	182	LYS
1	A	242	VAL
1	A	248	LEU
1	A	249	GLU
1	A	259	LYS
1	A	271	GLU
1	A	274	LYS
1	A	280	ASP
1	A	312	LEU

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	A	352	LYS
1	A	354	GLU
1	A	418	GLU
1	A	437	SER
1	A	441	VAL
1	A	457	SER
1	A	509	MET
1	A	510	VAL
1	A	512	LEU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (6) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	144	ASN
1	A	200	GLN
1	A	250	HIS
1	A	368	GLN
1	A	405	HIS
1	A	494	HIS

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

2 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The

Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
3	TBX	A	602	2	22,22,22	1.48	2 (9%)	32,32,32	1.77	3 (9%)
2	HEM	A	601	1,3	50,50,50	1.81	9 (18%)	67,82,82	1.26	6 (8%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	TBX	A	602	2	-	4/20/20/20	0/2/2/2
2	HEM	A	601	1,3	-	4/14/54/54	-

All (11) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	A	601	HEM	C3D-C2D	7.78	1.53	1.36
3	A	602	TBX	C19-C18	-5.29	1.50	1.53
3	A	602	TBX	C18-C31	-4.19	1.51	1.57
2	A	601	HEM	FE-NC	3.87	2.07	1.95
2	A	601	HEM	FE-NB	3.51	2.05	1.94
2	A	601	HEM	CAB-C3B	3.31	1.56	1.47
2	A	601	HEM	FE-ND	3.11	2.04	1.94
2	A	601	HEM	CAC-C3C	2.49	1.54	1.47
2	A	601	HEM	CMD-C2D	2.32	1.55	1.50
2	A	601	HEM	CMB-C2B	2.24	1.55	1.50
2	A	601	HEM	CMC-C2C	2.20	1.55	1.50

All (9) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	A	602	TBX	C15-C18-C19	8.61	117.47	111.44
2	A	601	HEM	C4D-ND-C1D	4.35	110.36	105.21
3	A	602	TBX	O29-C18-C15	-3.27	100.31	107.80
2	A	601	HEM	CHC-C4B-NB	3.22	127.89	124.42
2	A	601	HEM	CAD-C3D-C4D	3.20	130.28	124.70
2	A	601	HEM	CHD-C1D-ND	2.46	127.07	124.42

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	601	HEM	CAD-CBD-CGD	-2.23	107.75	113.67
3	A	602	TBX	C19-C18-C31	-2.11	107.27	108.80
2	A	601	HEM	CHA-C1A-NA	2.09	127.64	123.86

There are no chirality outliers.

All (8) torsion outliers are listed below:



Mol	Chain	Res	Type	Atoms
3	A	602	TBX	C12-C15-C18-C19
3	A	602	TBX	C12-C15-C18-O29
2	A	601	HEM	C4D-C3D-CAD-CBD
2	A	601	HEM	C2D-C3D-CAD-CBD
3	A	602	TBX	C15-C12-C3-C4
3	A	602	TBX	C15-C12-C3-C2
2	A	601	HEM	CAA-CBA-CGA-O2A
2	A	601	HEM	CAA-CBA-CGA-O1A

There are no ring outliers.

2 monomers are involved in 6 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	A	602	TBX	2	0
2	A	601	HEM	4	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.



5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	525/539 (97%)	0.60	38 (7%) 21 17	54, 79, 113, 134	0

All (38) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	248	LEU	4.8
1	A	441	VAL	4.4
1	A	9	GLY	4.1
1	A	241	PHE	3.6
1	A	247	PRO	3.5
1	A	12	LEU	3.3
1	A	439	TYR	3.2
1	A	8	VAL	3.2
1	A	41	TYR	2.9
1	A	298	LYS	2.9
1	A	532	GLY	2.8
1	A	422	ASN	2.7
1	A	246	LEU	2.6
1	A	236	PHE	2.5
1	A	250	HIS	2.5
1	A	243	PHE	2.5
1	A	352	LYS	2.4
1	A	135	GLY	2.4
1	A	28	ALA	2.4
1	A	425	GLN	2.4
1	A	128	HIS	2.3
1	A	284	ILE	2.2
1	A	527	GLU	2.2
1	A	275	ASN	2.2
1	A	290	ASN	2.2
1	A	45	TRP	2.1
1	A	103	TYR	2.1

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	25	LEU	2.1
1	A	66	VAL	2.1
1	A	24	ALA	2.1
1	A	132	PRO	2.1
1	A	225	PHE	2.1
1	A	15	VAL	2.0
1	A	517	ALA	2.0
1	A	531	GLY	2.0
1	A	138	VAL	2.0
1	A	252	ARG	2.0
1	A	447	TYR	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)




In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.


Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
3	TBX	A	602	21/21	0.91	0.14	57,65,76,81	0
2	HEM	A	601	43/43	0.97	0.12	52,62,67,68	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.

Electron density around TBX A 602:

$2mF_o - DF_c$ (at 0.7 rmsd) in gray
 $mF_o - DF_c$ (at 3 rmsd) in purple (negative)
and green (positive)





6.5 Other polymers [i](#)

There are no such residues in this entry.