



# wwPDB X-ray Structure Validation Summary Report ⓘ

Mar 8, 2026 – 01:56 PM UTC

PDB ID : 4EBK / pdb\_00004ebk  
Title : Crystal structure of aminoglycoside 4'-O-adenylyltransferase ANT(4')-IIb, tobramycin-bound  
Authors : Stogios, P.J.; Dong, A.; Minasov, G.; Evdokimova, E.; Egorova, O.; Yim, V.; Kudritska, M.; Courvalin, P.; Savchenko, A.; Anderson, W.F.; Center for Structural Genomics of Infectious Diseases (CSGID)  
Deposited on : 2012-03-23  
Resolution : 2.15 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

---

The following versions of software and data (see [references](#) ①) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0  
Mogul : 2022.3.0, CSD as543be (2022)  
Xtrriage (Phenix) : 2.0  
EDS : 3.0  
Buster-report : wwPDB partial adaption of 1.1.7 (2018)  
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)  
CCP4 : 9.0.010 (Gargrove)  
Density-Fitness : 1.0.12  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.49


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*



The reported resolution of this entry is 2.15 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	180053	2057 (2.16-2.16)
Clashscore	190562	2159 (2.16-2.16)
Ramachandran outliers	187476	2134 (2.16-2.16)
Sidechain outliers	187428	2133 (2.16-2.16)
RSRZ outliers	180081	2059 (2.16-2.16)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	272	 82% 12% 6%
1	B	272	 85% 9% 6%

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
4	IOD	A	307	-	-	X	-
4	IOD	A	308	-	-	X	-
4	IOD	B	305	-	-	X	-
5	CL	A	310	-	-	X	-
5	CL	A	319	-	-	X	-

## 2 Entry composition [i](#)

There are 6 unique types of molecules in this entry. The entry contains 4728 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Aminoglycoside nucleotidyltransferase.

Mol	Chain	Residues	Atoms							ZeroOcc	AltConf	Trace
			Total	C	I	N	O	S	Se			
1	A	257	2074	1308	1	366	393	1	5	0	6	0
1	B	256	2066	1302	1	368	390	1	4	0	5	0

There are 48 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
A	-20	MSE	-	initiating methionine	UNP Q83V96
A	-19	GLY	-	expression tag	UNP Q83V96
A	-18	SER	-	expression tag	UNP Q83V96
A	-17	SER	-	expression tag	UNP Q83V96
A	-16	HIS	-	expression tag	UNP Q83V96
A	-15	HIS	-	expression tag	UNP Q83V96
A	-14	HIS	-	expression tag	UNP Q83V96
A	-13	HIS	-	expression tag	UNP Q83V96
A	-12	HIS	-	expression tag	UNP Q83V96
A	-11	HIS	-	expression tag	UNP Q83V96
A	-10	SER	-	expression tag	UNP Q83V96
A	-9	SER	-	expression tag	UNP Q83V96
A	-8	GLY	-	expression tag	UNP Q83V96
A	-7	ARG	-	expression tag	UNP Q83V96
A	-6	GLU	-	expression tag	UNP Q83V96
A	-5	ASN	-	expression tag	UNP Q83V96
A	-4	LEU	-	expression tag	UNP Q83V96
A	-3	TYR	-	expression tag	UNP Q83V96
A	-2	PHE	-	expression tag	UNP Q83V96
A	-1	GLN	-	expression tag	UNP Q83V96
A	0	GLY	-	expression tag	UNP Q83V96
A	1	VAL	-	expression tag	UNP Q83V96
A	111	LEU	MET	SEE REMARK 999	UNP Q83V96
A	127	ASP	ASN	SEE REMARK 999	UNP Q83V96
B	-20	MSE	-	initiating methionine	UNP Q83V96

*Continued on next page...*



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
2	A	1	Total	C	N	O	0	0
			32	18	5	9		
2	B	1	Total	C	N	O	0	0
			32	18	5	9		

- Molecule 3 is 1,2-ETHANEDIOL (CCD ID: EDO) (formula: C<sub>2</sub>H<sub>6</sub>O<sub>2</sub>).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	A	1	Total	C	O	0	0
			4	2	2		
3	B	1	Total	C	O	0	0
			4	2	2		

- Molecule 4 is IODIDE ION (CCD ID: IOD) (formula: I).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	6	Total	I	0	0
			6	6		
4	B	6	Total	I	0	0
			6	6		

- Molecule 5 is CHLORIDE ION (CCD ID: CL) (formula: Cl).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	11	Total	Cl	0	0
			11	11		

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	B	6	Total	Cl	0	0
			6	6		


- Molecule 6 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
6	A	241	Total	O	0	11
			250	250		
6	B	234	Total	O	0	4
			237	237		


### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Aminoglycoside nucleotidyltransferase



- Molecule 1: Aminoglycoside nucleotidyltransferase



## 4 Data and refinement statistics i

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	81.91Å 84.45Å 98.76Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	23.89 – 2.15 23.89 – 2.15	Depositor EDS
% Data completeness (in resolution range)	99.5 (23.89-2.15) 97.4 (23.89-2.15)	Depositor EDS
$R_{merge}$	(Not available)	Depositor
$R_{sym}$	0.06	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	1.51 (at 2.15Å)	Xtrriage
Refinement program	PHENIX (phenix.refine: 1.7.3_928)	Depositor
R, $R_{free}$	0.172 , 0.221 0.160 , 0.208	Depositor DCC
$R_{free}$ test set	2012 reflections (5.31%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	30.5	Xtrriage
Anisotropy	0.325	Xtrriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.35 , 61.3	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.49$ , $\langle L^2 \rangle = 0.32$	Xtrriage
Estimated twinning fraction	0.021 for k,h,-l	Xtrriage
$F_o, F_c$ correlation	0.96	EDS
Total number of atoms	4728	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	38.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.53% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: EDO, CL, TOY, IOD, IYR

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# $ Z  > 5$	RMSZ	# $ Z  > 5$
1	A	0.50	1/2111 (0.0%)	0.78	3/2871 (0.1%)
1	B	0.49	0/2097	0.77	0/2851
All	All	0.49	1/4208 (0.0%)	0.77	3/5722 (0.1%)

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	163	ILE	CA-CB	5.21	1.56	1.54

All (3) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	187	VAL	CA-C-N	-7.07	113.52	120.52
1	A	187	VAL	C-N-CA	-7.07	113.52	120.52
1	A	-4	LEU	N-CA-C	-5.03	107.14	113.28

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2074	0	2053	23	0
1	B	2066	0	2043	19	0

*Continued on next page...*

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
2	A	32	0	37	0	0
2	B	32	0	37	0	0
3	A	4	0	6	1	0
3	B	4	0	6	0	0
4	A	6	0	0	5	0
4	B	6	0	0	3	0
5	A	11	0	0	4	0
5	B	6	0	0	1	0
6	A	250	0	0	2	0
6	B	237	0	0	2	0
All	All	4728	0	4182	43	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 5.

The worst 5 of 43 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:156:LYS:NZ	5:A:310:CL:CL	2.24	1.07
1:A:191:PHE:HD1	1:A:217[B]:MSE:HE1	1.39	0.88
1:B:217:MSE:HG2	4:B:305:IOD:I	2.53	0.79
1:A:217[B]:MSE:SE	4:A:308:IOD:I	3.22	0.77
5:A:313:CL:CL	6:A:633:HOH:O	2.48	0.69

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	260/272 (96%)	258 (99%)	2 (1%)	0	100	100
1	B	258/272 (95%)	255 (99%)	3 (1%)	0	100	100

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
All	All	518/544 (95%)	513 (99%)	5 (1%)	0	100	100

There are no Ramachandran outliers to report.

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	218/220 (99%)	215 (99%)	3 (1%)	59	66
1	B	216/220 (98%)	215 (100%)	1 (0%)	81	87
All	All	434/440 (99%)	430 (99%)	4 (1%)	70	77

All (4) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	48	GLN
1	A	79	ARG
1	A	203	GLU
1	B	40	ASP

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (1) such sidechains are listed below:

Mol	Chain	Res	Type
1	B	241	ASN

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

2 non-standard protein/DNA/RNA residues are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z  > 2$	Counts	RMSZ	$\# Z  > 2$
1	IYR	A	16	1	12,13,14	1.16	0	12,17,19	1.48	2 (16%)
1	IYR	B	16	1	12,13,14	1.32	2 (16%)	12,17,19	1.05	1 (8%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
1	IYR	A	16	1	-	1/5/6/8	0/1/1/1
1	IYR	B	16	1	-	1/5/6/8	0/1/1/1

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	B	16	IYR	O-C	2.23	1.28	1.20
1	B	16	IYR	OF-CF	2.17	1.40	1.36

All (3) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	16	IYR	CF-CE-IE	-3.81	116.08	119.76
1	B	16	IYR	CF-CE-IE	-2.73	117.13	119.76
1	A	16	IYR	CD-CE-IE	2.24	122.66	118.46

There are no chirality outliers.

All (2) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
1	A	16	IYR	O-C-CA-CB
1	B	16	IYR	O-C-CA-CB

There are no ring outliers.

1 monomer is involved in 2 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
1	B	16	IYR	2	0

## 5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

## 5.6 Ligand geometry [i](#)

Of 33 ligands modelled in this entry, 29 are monoatomic - leaving 4 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# $ Z  > 2$	Counts	RMSZ	# $ Z  > 2$
3	EDO	B	303	-	3,3,3	0.44	0	2,2,2	0.17	0
2	TOY	B	302	-	33,34,34	0.92	1 (3%)	43,50,50	1.26	5 (11%)
3	EDO	A	302	-	3,3,3	0.44	0	2,2,2	0.22	0
2	TOY	A	301	-	33,34,34	0.74	1 (3%)	43,50,50	1.39	7 (16%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	EDO	B	303	-	-	0/1/1/1	-
2	TOY	B	302	-	-	3/12/68/68	0/3/3/3
3	EDO	A	302	-	-	0/1/1/1	-
2	TOY	A	301	-	-	3/12/68/68	0/3/3/3

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	A	301	TOY	C23-C33	-2.89	1.49	1.53

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	B	302	TOY	C23-C33	-2.14	1.50	1.53

The worst 5 of 12 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	301	TOY	O51-C51-C61	-3.46	99.42	106.07
2	A	301	TOY	C11-O11-C42	3.10	125.33	117.98
2	A	301	TOY	C53-C43-C33	-3.08	107.27	110.67
2	A	301	TOY	C61-C51-C41	-2.90	105.98	112.75
2	B	302	TOY	C11-O11-C42	2.65	124.27	117.98

There are no chirality outliers.

5 of 6 torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
2	B	302	TOY	C52-C42-O11-C11
2	A	301	TOY	C52-C42-O11-C11
2	A	301	TOY	C32-C42-O11-C11
2	B	302	TOY	C32-C42-O11-C11
2	A	301	TOY	O51-C11-O11-C42


There are no ring outliers.

1 monomer is involved in 1 short contact:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	A	302	EDO	1	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.





## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	252/272 (92%)	-0.41	2 (0%) 82 84	14, 30, 53, 99	5 (1%)
1	B	251/272 (92%)	-0.22	1 (0%) 88 90	15, 34, 71, 86	5 (1%)
All	All	503/544 (92%)	-0.32	3 (0%) 85 87	14, 32, 64, 99	10 (1%)

All (3) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	-4	LEU	4.0
1	B	-4	LEU	3.1
1	A	-2	PHE	2.5

### 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q < 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å <sup>2</sup> )	Q<0.9
1	IYR	A	16	13/14	0.92	0.09	34,39,42,45	1
1	IYR	B	16	13/14	0.92	0.11	47,50,53,65	1

### 6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.


## 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å <sup>2</sup> )	Q<0.9
5	CL	A	312	1/1	0.77	0.17	67,67,67,67	0
3	EDO	B	303	4/4	0.83	0.16	55,55,58,62	0
3	EDO	A	302	4/4	0.85	0.14	55,57,60,63	0
5	CL	A	310	1/1	0.89	0.18	66,66,66,66	0
5	CL	A	315	1/1	0.89	0.12	66,66,66,66	0
4	IOD	A	307	1/1	0.90	0.09	79,79,79,79	1
4	IOD	B	304	1/1	0.91	0.10	59,59,59,59	1
5	CL	A	319	1/1	0.91	0.11	56,56,56,56	0
5	CL	A	311	1/1	0.92	0.10	62,62,62,62	0
5	CL	A	309	1/1	0.92	0.07	47,47,47,47	0
5	CL	A	314	1/1	0.93	0.10	65,65,65,65	0
5	CL	A	317	1/1	0.93	0.19	64,64,64,64	0
4	IOD	A	306	1/1	0.94	0.07	42,42,42,42	1
2	TOY	A	301	32/32	0.94	0.07	27,39,45,50	0
2	TOY	B	302	32/32	0.94	0.08	24,36,50,55	0
4	IOD	B	305	1/1	0.94	0.07	45,45,45,45	1
5	CL	B	310	1/1	0.95	0.09	54,54,54,54	0
4	IOD	A	303	1/1	0.96	0.06	63,63,63,63	1
5	CL	A	313	1/1	0.96	0.07	52,52,52,52	0
5	CL	A	318	1/1	0.96	0.11	60,60,60,60	0
4	IOD	B	309	1/1	0.96	0.08	35,35,35,35	1
5	CL	B	311	1/1	0.96	0.14	56,56,56,56	0
5	CL	B	312	1/1	0.96	0.18	67,67,67,67	0
5	CL	B	313	1/1	0.96	0.05	42,42,42,42	0
4	IOD	B	307	1/1	0.97	0.07	56,56,56,56	1
4	IOD	B	306	1/1	0.97	0.05	52,52,52,52	1
5	CL	A	316	1/1	0.97	0.05	38,38,38,38	0
4	IOD	A	308	1/1	0.98	0.03	34,34,34,34	1
5	CL	B	301	1/1	0.98	0.06	45,45,45,45	0
5	CL	B	314	1/1	0.98	0.04	56,56,56,56	0
4	IOD	A	305	1/1	0.99	0.04	35,35,35,35	1
4	IOD	B	308	1/1	0.99	0.03	38,38,38,38	1
4	IOD	A	304	1/1	1.00	0.01	30,30,30,30	1

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different

orientation to approximate a three-dimensional view.



## 6.5 Other polymers [i](#)

There are no such residues in this entry.