



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 9, 2026 – 03:33 PM UTC

PDB ID : 5EBW / pdb_00005ebw
Title : KcsA with G77ester mutation
Authors : Matulef, K.; Valiyaveetil, F.I.
Deposited on : 2015-10-19
Resolution : 2.30 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0
Mogul : 2022.3.0, CSD as543be (2022)
Xtriage (Phenix) : 2.0
EDS : 3.0
Buster-report : wwPDB partial adaption of 1.1.7 (2018)
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4 : 9.0.010 (Gargrove)
Density-Fitness : 1.0.12
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.30 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	6319 (2.30-2.30)
Clashscore	190562	6919 (2.30-2.30)
Ramachandran outliers	187476	6854 (2.30-2.30)
Sidechain outliers	187428	6854 (2.30-2.30)
RSRZ outliers	180081	6325 (2.30-2.30)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	218	24% 91% 9%
2	B	210	22% 94% 6%
3	C	123	4% 77% 6% 17%

2 Entry composition

There are 7 unique types of molecules in this entry. The entry contains 4215 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Antibody Fab Fragment Light Chain.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	218	Total	C	N	O	S	0	0	0
			1630	1032	272	320	6			

- Molecule 2 is a protein called Antibody Fab Fragment Light Chain.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	B	210	Total	C	N	O	S	0	0	0
			1622	1009	275	333	5			


- Molecule 3 is a protein called pH-gated potassium channel KcsA.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
3	C	102	Total	C	N	O	S	0	0	0
			755	497	127	129	2			

There are 4 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
C	2	ALA	PRO	engineered mutation	UNP P0A334
C	69	ALA	SER	engineered mutation	UNP P0A334
C	70	CYS	VAL	engineered mutation	UNP P0A334
C	77	GOA	GLY	engineered mutation	UNP P0A334

- Molecule 4 is NONAN-1-OL (CCD ID: F09) (formula: C₉H₂₀O).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
4	A	1	Total	C	O	0	0
			10	9	1		

- Molecule 5 is DIACYL GLYCEROL (CCD ID: DGA) (formula: $C_{39}H_{76}O_5$).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
5	C	1	Total	C	O	0	0
			31	26	5		

- Molecule 6 is POTASSIUM ION (CCD ID: K) (formula: K).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
6	C	6	Total K 6 6	0	0


- Molecule 7 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
7	A	48	Total O 48 48	0	0
7	B	85	Total O 85 85	0	0
7	C	28	Total O 28 28	0	0


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


• Molecule 1: Antibody Fab Fragment Light Chain



• Molecule 2: Antibody Fab Fragment Light Chain



• Molecule 3: pH-gated potassium channel KcsA



4 Data and refinement statistics

Property	Value	Source
Space group	I 4	Depositor
Cell constants a, b, c, α , β , γ	156.91Å 156.91Å 75.84Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	39.23 – 2.30 39.23 – 2.30	Depositor EDS
% Data completeness (in resolution range)	93.7 (39.23-2.30) 93.1 (39.23-2.30)	Depositor EDS
R_{merge}	0.12	Depositor
R_{sym}	0.12	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.11 (at 2.12Å)	Xtriage
Refinement program	PHENIX 1.8_1069	Depositor
R, R_{free}	0.222 , 0.249 0.221 , 0.246	Depositor DCC
R_{free} test set	5048 reflections (9.63%)	wwPDB-VP
Wilson B-factor (Å ²)	32.2	Xtriage
Anisotropy	0.421	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.30 , 39.9	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	0.028 for -k,-h,-l	Xtriage
F_o, F_c correlation	0.93	EDS
Total number of atoms	4215	wwPDB-VP
Average B, all atoms (Å ²)	56.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.77% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: GOA, K, DGA, F09

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.28	0/1674	0.72	2/2291 (0.1%)
2	B	0.27	0/1659	0.69	0/2253
3	C	0.35	0/770	0.70	0/1059
All	All	0.29	0/4103	0.70	2/5603 (0.0%)

There are no bond length outliers.

All (2) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	204	HIS	CA-C-N	5.38	125.72	119.47
1	A	204	HIS	C-N-CA	5.38	125.72	119.47

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1630	0	1584	13	0
2	B	1622	0	1539	7	0
3	C	755	0	760	10	0
4	A	10	0	20	0	0
5	C	31	0	44	5	0
6	C	6	0	0	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
7	A	48	0	0	1	0
7	B	85	0	0	0	0
7	C	28	0	0	1	0
All	All	4215	0	3947	27	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 3.

All (27) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:83:LEU:HB3	1:A:86:LEU:HD21	1.74	0.69
3:C:89:ARG:HH11	5:C:1001:DGA:CA2	2.13	0.61
2:B:37:GLN:HB2	2:B:47:LEU:HD11	1.84	0.59
1:A:1:GLN:N	7:A:2101:HOH:O	2.37	0.58
1:A:1:GLN:N	1:A:1:GLN:OE1	2.30	0.57
1:A:38:LYS:HB2	1:A:48:ILE:HD11	1.86	0.55
1:A:69:THR:HB	1:A:82:GLN:HB3	1.90	0.53
3:C:89:ARG:HH11	5:C:1001:DGA:HA21	1.75	0.51
3:C:23:ALA:HB1	3:C:25:HIS:CE1	2.46	0.51
3:C:86:LEU:HD13	5:C:1001:DGA:HA31	1.93	0.51
3:C:89:ARG:HH11	5:C:1001:DGA:HA22	1.76	0.50
2:B:19:VAL:HG12	2:B:75:ILE:HB	1.93	0.49
1:A:31:SER:HB2	3:C:62:TYR:CE1	2.50	0.47
1:A:173:ALA:HB2	1:A:182:LEU:HD23	1.96	0.46
2:B:93:ARG:NH1	3:C:56:GLY:O	2.48	0.46
2:B:149:LYS:HB2	2:B:193:THR:HB	1.97	0.46
3:C:23:ALA:HB3	3:C:26:TRP:HD1	1.84	0.43
1:A:124:PRO:HB3	1:A:150:TYR:HB3	2.01	0.43
2:B:35:TRP:CE2	2:B:73:LEU:HB2	2.54	0.42
1:A:169:HIS:HE1	2:B:138:ASN:ND2	2.18	0.42
2:B:207:LYS:HA	2:B:207:LYS:HD3	1.90	0.42
1:A:102:ASP:OD1	1:A:102:ASP:N	2.51	0.41
1:A:23:LYS:HB2	1:A:23:LYS:HE3	1.87	0.41
1:A:213:LYS:HA	1:A:213:LYS:HD3	1.86	0.41
3:C:25:HIS:ND1	7:C:1102:HOH:O	2.37	0.41
1:A:151:PHE:HA	1:A:152:PRO:HA	1.79	0.41
3:C:89:ARG:NH1	5:C:1001:DGA:HA21	2.35	0.41

There are no symmetry-related clashes.

5.3 Torsion angles

5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	216/218 (99%)	209 (97%)	7 (3%)	0	100	100
2	B	208/210 (99%)	201 (97%)	7 (3%)	0	100	100
3	C	98/123 (80%)	97 (99%)	1 (1%)	0	100	100
All	All	522/551 (95%)	507 (97%)	15 (3%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	180/184 (98%)	180 (100%)	0	100	100
2	B	185/188 (98%)	183 (99%)	2 (1%)	65	81
3	C	71/89 (80%)	71 (100%)	0	100	100
All	All	436/461 (95%)	434 (100%)	2 (0%)	81	90

All (2) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
2	B	181	LEU
2	B	184	ASP

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (5) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	169	HIS
2	B	137	ASN
2	B	138	ASN
3	C	20	HIS
3	C	25	HIS

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 8 ligands modelled in this entry, 6 are monoatomic - leaving 2 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
5	DGA	C	1001	-	30,30,43	1.05	2 (6%)	32,32,45	1.41	3 (9%)
4	F09	A	2001	-	9,9,9	0.22	0	8,8,8	0.49	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
5	DGA	C	1001	-	-	17/32/32/45	-
4	F09	A	2001	-	-	4/7/7/7	-

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
5	C	1001	DGA	OG2-CB1	3.95	1.45	1.34
5	C	1001	DGA	OG1-CA1	3.25	1.42	1.33

All (3) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
5	C	1001	DGA	OG2-CB1-CB2	4.32	120.83	111.48
5	C	1001	DGA	OG1-CA1-OA1	-3.54	114.78	123.63
5	C	1001	DGA	OG1-CA1-CA2	3.09	121.27	111.83

There are no chirality outliers.

All (21) torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
5	C	1001	DGA	CB2-CB1-OG2-CG2
5	C	1001	DGA	CG1-CG2-CG3-OXT
5	C	1001	DGA	OG2-CG2-CG3-OXT
5	C	1001	DGA	OB1-CB1-OG2-CG2
5	C	1001	DGA	OG1-CG1-CG2-OG2
5	C	1001	DGA	CB1-CB2-CB3-CB4
5	C	1001	DGA	CB6-CB7-CB8-CB9
4	A	2001	F09	C5-C6-C7-C8
5	C	1001	DGA	CB5-CB6-CB7-CB8
5	C	1001	DGA	CA2-CA3-CA4-CA5
5	C	1001	DGA	CA5-CA6-CA7-CA8
4	A	2001	F09	C7-C8-C9-OXT
4	A	2001	F09	C4-C5-C6-C7
5	C	1001	DGA	CA3-CA4-CA5-CA6
5	C	1001	DGA	OG1-CG1-CG2-CG3
4	A	2001	F09	C6-C7-C8-C9
5	C	1001	DGA	CA6-CA7-CA8-CA9
5	C	1001	DGA	CB9-CAB-CBB-CCB
5	C	1001	DGA	CB4-CB5-CB6-CB7
5	C	1001	DGA	OG1-CA1-CA2-CA3
5	C	1001	DGA	CB3-CB4-CB5-CB6

There are no ring outliers.

1 monomer is involved in 5 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
5	C	1001	DGA	5	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.



5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	218/218 (100%)	1.45	53 (24%) 2 2	33, 65, 82, 93	0
2	B	210/210 (100%)	1.08	46 (21%) 2 2	28, 59, 89, 103	0
3	C	101/123 (82%)	0.22	5 (4%) 34 35	24, 33, 59, 86	0
All	All	529/551 (96%)	1.07	104 (19%) 3 3	24, 58, 84, 103	0

All (104) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	42	GLY	4.9
1	A	24	ALA	4.5
3	C	23	ALA	4.5
3	C	21	GLY	4.2
1	A	196	GLU	4.2
1	A	137	THR	4.0
2	B	208	SER	4.0
2	B	190	ASN	3.9
1	A	163	SER	3.9
1	A	55	TYR	3.8
3	C	20	HIS	3.7
1	A	120	LYS	3.7
2	B	125	LEU	3.6
1	A	197	THR	3.6
1	A	2	VAL	3.4
2	B	189	HIS	3.3
2	B	188	ARG	3.3
2	B	135	PHE	3.2
1	A	213	LYS	3.2
2	B	179	LEU	3.2
2	B	186	TYR	3.1
2	B	148	TRP	3.1
1	A	27	TYR	3.0

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	25	SER	3.0
2	B	165	ASP	3.0
1	A	134	ALA	3.0
1	A	216	VAL	2.9
1	A	119	ALA	2.9
2	B	210	ASN	2.9
1	A	43	HIS	2.8
2	B	118	PHE	2.8
2	B	195	GLU	2.8
1	A	124	PRO	2.8
1	A	174	VAL	2.8
1	A	199	THR	2.8
2	B	75	ILE	2.8
1	A	165	SER	2.8
3	C	22	SER	2.7
1	A	164	LEU	2.7
2	B	156	GLN	2.7
1	A	167	GLY	2.7
1	A	195	SER	2.7
2	B	153	SER	2.7
2	B	209	PHE	2.7
1	A	30	THR	2.7
1	A	85	SER	2.7
1	A	146	LEU	2.7
1	A	84	SER	2.6
1	A	177	SER	2.6
1	A	183	SER	2.6
2	B	145	ASN	2.6
2	B	205	ILE	2.6
2	B	160	LEU	2.6
2	B	163	TRP	2.6
1	A	136	GLN	2.6
1	A	1	GLN	2.5
2	B	130	ALA	2.5
1	A	185	SER	2.5
1	A	214	LYS	2.5
3	C	120	GLU	2.5
1	A	3	GLN	2.5
2	B	117	ILE	2.4
1	A	169	HIS	2.4
1	A	5	GLN	2.4
2	B	115	VAL	2.4

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
2	B	136	LEU	2.4
1	A	62	GLU	2.4
2	B	193	THR	2.3
1	A	151	PHE	2.3
1	A	159	TRP	2.3
2	B	126	THR	2.3
2	B	180	THR	2.3
2	B	127	SER	2.3
2	B	146	VAL	2.3
2	B	128	GLY	2.3
2	B	192	TYR	2.3
1	A	208	SER	2.3
1	A	179	LEU	2.2
1	A	194	PRO	2.2
1	A	121	THR	2.2
2	B	182	THR	2.2
1	A	193	TRP	2.2
1	A	28	THR	2.2
2	B	181	LEU	2.2
1	A	127	TYR	2.1
1	A	143	LEU	2.1
2	B	175	MET	2.1
1	A	198	VAL	2.1
2	B	120	PRO	2.1
2	B	60	SER	2.1
1	A	83	LEU	2.1
1	A	122	THR	2.1
2	B	77	SER	2.1
2	B	168	SER	2.1
2	B	202	THR	2.1
1	A	82	GLN	2.1
1	A	132	GLY	2.1
2	B	183	LYS	2.0
2	B	187	GLU	2.0
2	B	194	CYS	2.0
2	B	121	SER	2.0
2	B	73	LEU	2.0
1	A	14	PRO	2.0
2	B	150	ILE	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)




In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
5	DGA	C	1001	31/44	0.72	0.29	66,68,72,75	0
6	K	C	1007	1/1	0.76	0.19	65,65,65,65	1
4	F09	A	2001	10/10	0.81	0.23	61,61,62,62	0
6	K	C	1003	1/1	0.94	0.15	49,49,49,49	1
6	K	C	1004	1/1	0.97	0.05	30,30,30,30	1
6	K	C	1005	1/1	0.99	0.04	34,34,34,34	1
6	K	C	1006	1/1	0.99	0.06	45,45,45,45	1
6	K	C	1002	1/1	0.99	0.04	34,34,34,34	1

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.

Electron density around DGA C 1001:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



6.5 Other polymers [i](#)

There are no such residues in this entry.