



wwPDB X-ray Structure Validation Summary Report

Mar 18, 2026 – 04:18 AM UTC

PDB ID : 6EB8 / pdb_00006eb8
Title : Crystal Structure of the Nipah Virus Phosphoprotein Multimerization Domain G519N
Authors : Bruhn, J.F.; Sapphire, E.O.
Deposited on : 2018-08-06
Resolution : 2.50 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the  symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references](#) ) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0
Xtrriage (Phenix) : 2.0
EDS : 3.0
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4 : 9.0.010 (Gargrove)
Density-Fitness : 1.0.12
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.50 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.






Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	5829 (2.50-2.50)
Clashscore	190562	6492 (2.50-2.50)
Ramachandran outliers	187476	6378 (2.50-2.50)
Sidechain outliers	187428	6380 (2.50-2.50)
RSRZ outliers	180081	5833 (2.50-2.50)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	110	
1	B	110	
1	C	110	
1	D	110	
1	E	110	

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
1	F	110	
1	G	110	
1	H	110	

2 Entry composition

There are 2 unique types of molecules in this entry. The entry contains 6673 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Phosphoprotein.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	104	840	530	138	165	7	0	0	0
1	E	103	838	529	138	164	7	0	1	0
1	D	101	816	515	135	160	6	0	0	0
1	C	100	807	509	133	159	6	0	0	0
1	H	101	816	515	135	160	6	0	0	0
1	F	101	816	515	135	160	6	0	0	0
1	B	100	808	509	134	159	6	0	0	0
1	G	101	816	515	135	160	6	0	0	0

There are 16 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	469	SER	-	expression tag	UNP Q9IK91
A	519	ASN	GLY	engineered mutation	UNP Q9IK91
E	469	SER	-	expression tag	UNP Q9IK91
E	519	ASN	GLY	engineered mutation	UNP Q9IK91
D	469	SER	-	expression tag	UNP Q9IK91
D	519	ASN	GLY	engineered mutation	UNP Q9IK91
C	469	SER	-	expression tag	UNP Q9IK91
C	519	ASN	GLY	engineered mutation	UNP Q9IK91
H	469	SER	-	expression tag	UNP Q9IK91
H	519	ASN	GLY	engineered mutation	UNP Q9IK91
F	469	SER	-	expression tag	UNP Q9IK91
F	519	ASN	GLY	engineered mutation	UNP Q9IK91
B	469	SER	-	expression tag	UNP Q9IK91

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
B	519	ASN	GLY	engineered mutation	UNP Q9IK91
G	469	SER	-	expression tag	UNP Q9IK91
G	519	ASN	GLY	engineered mutation	UNP Q9IK91

- Molecule 2 is water.


Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
2	A	9	Total O 9 9	0	0
2	E	20	Total O 20 20	0	0
2	D	12	Total O 12 12	0	0
2	C	22	Total O 22 22	0	0
2	H	5	Total O 5 5	0	0
2	F	22	Total O 22 22	0	0
2	B	10	Total O 10 10	0	0
2	G	16	Total O 16 16	0	0

3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Phosphoprotein

Chain A: 




- Molecule 1: Phosphoprotein

Chain E: 



- Molecule 1: Phosphoprotein


Chain D: 




- Molecule 1: Phosphoprotein

Chain C: 







● Molecule 1: Phosphoprotein




● Molecule 1: Phosphoprotein



● Molecule 1: Phosphoprotein



● Molecule 1: Phosphoprotein



4 Data and refinement statistics

Property	Value	Source
Space group	P 1	Depositor
Cell constants a, b, c, α , β , γ	48.00Å 76.62Å 80.54Å 100.46° 100.94° 108.05°	Depositor
Resolution (Å)	38.20 – 2.50 38.20 – 2.50	Depositor EDS
% Data completeness (in resolution range)	98.5 (38.20-2.50) 86.8 (38.20-2.50)	Depositor EDS
R_{merge}	0.31	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	0.65 (at 2.51Å)	Xtrriage
Refinement program	PHENIX 1.13_2998	Depositor
R, R_{free}	0.276 , 0.338 0.276 , 0.339	Depositor DCC
R_{free} test set	1771 reflections (2.16%)	wwPDB-VP
Wilson B-factor (Å ²)	26.3	Xtrriage
Anisotropy	0.663	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.37 , 67.5	EDS
L-test for twinning ²	$\langle L \rangle = 0.46$, $\langle L^2 \rangle = 0.29$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
F_o, F_c correlation	0.87	EDS
Total number of atoms	6673	wwPDB-VP
Average B, all atoms (Å ²)	43.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 15.34% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# $ Z > 5$	RMSZ	# $ Z > 5$
1	A	0.42	0/853	0.68	0/1153
1	B	0.42	0/821	0.67	0/1110
1	C	0.45	0/820	0.70	0/1110
1	D	0.51	0/829	0.75	0/1121
1	E	0.54	0/851	0.76	1/1150 (0.1%)
1	F	0.50	0/829	0.75	0/1121
1	G	0.83	4/829 (0.5%)	0.76	0/1121
1	H	0.60	0/829	0.74	1/1121 (0.1%)
All	All	0.55	4/6661 (0.1%)	0.73	2/9007 (0.0%)

All (4) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	G	522	ASN	C-O	-7.70	1.15	1.24
1	G	526	LEU	C-O	-7.67	1.15	1.24
1	G	523	SER	C-O	-6.26	1.16	1.24
1	G	527	ILE	C-O	-5.79	1.17	1.24

All (2) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	H	502	HIS	N-CA-C	5.88	121.50	113.97
1	E	577	MET	N-CA-C	5.18	116.92	111.28

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen

atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	840	0	836	43	0
1	B	808	0	801	29	0
1	C	807	0	799	33	0
1	D	816	0	812	39	1
1	E	838	0	836	57	1
1	F	816	0	812	47	0
1	G	816	0	812	53	0
1	H	816	0	812	57	0
2	A	9	0	0	1	0
2	B	10	0	0	1	0
2	C	22	0	0	1	0
2	D	12	0	0	0	0
2	E	20	0	0	0	0
2	F	22	0	0	1	0
2	G	16	0	0	1	0
2	H	5	0	0	1	0
All	All	6673	0	6520	237	1

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 18.

The worst 5 of 237 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:H:508:LEU:HD22	1:G:508:LEU:CD1	1.56	1.34
1:H:508:LEU:CD2	1:G:508:LEU:HD13	1.71	1.20
1:H:508:LEU:HD22	1:G:508:LEU:HD13	1.14	1.08
1:F:525:LYS:NZ	1:G:523:SER:OG	1.87	1.08
1:H:508:LEU:CD2	1:G:508:LEU:CD1	2.35	1.02

All (1) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:E:577:MET:O	1:D:555:ARG:NH1[1_446]	1.68	0.52

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	102/110 (93%)	101 (99%)	1 (1%)	0	100	100
1	B	98/110 (89%)	98 (100%)	0	0	100	100
1	C	98/110 (89%)	96 (98%)	2 (2%)	0	100	100
1	D	99/110 (90%)	97 (98%)	2 (2%)	0	100	100
1	E	102/110 (93%)	100 (98%)	2 (2%)	0	100	100
1	F	99/110 (90%)	99 (100%)	0	0	100	100
1	G	99/110 (90%)	99 (100%)	0	0	100	100
1	H	99/110 (90%)	96 (97%)	3 (3%)	0	100	100
All	All	796/880 (90%)	786 (99%)	10 (1%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	99/105 (94%)	96 (97%)	3 (3%)	36	64
1	B	95/105 (90%)	94 (99%)	1 (1%)	65	84
1	C	95/105 (90%)	95 (100%)	0	100	100
1	D	96/105 (91%)	94 (98%)	2 (2%)	47	74
1	E	99/105 (94%)	97 (98%)	2 (2%)	48	75
1	F	96/105 (91%)	95 (99%)	1 (1%)	68	86

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	G	96/105 (91%)	95 (99%)	1 (1%)	68	86
1	H	96/105 (91%)	95 (99%)	1 (1%)	68	86
All	All	772/840 (92%)	761 (99%)	11 (1%)	59	81

5 of 11 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	H	493	THR
1	F	561	ASN
1	G	560	THR
1	B	523	SER
1	E	578	ILE

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 9 such sidechains are listed below:

Mol	Chain	Res	Type
1	F	561	ASN
1	G	570	HIS
1	C	570	HIS
1	H	539	GLN
1	H	561	ASN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

There are no ligands in this entry.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled '#RSRZ > 2' contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled 'Q < 0.9' lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	104/110 (94%)	1.30	21 (20%) 3 2	21, 45, 56, 65	0
1	B	100/110 (90%)	1.24	18 (18%) 3 3	26, 42, 56, 68	0
1	C	100/110 (90%)	1.21	16 (16%) 5 4	24, 41, 56, 60	0
1	D	101/110 (91%)	1.35	20 (19%) 3 2	19, 46, 59, 65	0
1	E	103/110 (93%)	1.09	13 (12%) 8 6	20, 40, 54, 63	1 (0%)
1	F	101/110 (91%)	1.07	9 (8%) 15 13	26, 39, 51, 59	0
1	G	101/110 (91%)	1.39	24 (23%) 2 1	23, 47, 58, 73	0
1	H	101/110 (91%)	1.39	16 (15%) 5 4	29, 43, 56, 75	0
All	All	811/880 (92%)	1.25	137 (16%) 4 3	19, 43, 57, 75	1 (0%)

The worst 5 of 137 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	501	ASP	6.0
1	H	496	LEU	4.9
1	D	488	PHE	4.2
1	C	477	TYR	3.9
1	D	576	ILE	3.9

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands

There are no ligands in this entry.

6.5 Other polymers

There are no such residues in this entry.