



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 8, 2026 – 01:16 AM UTC

PDB ID : 8EB3 / pdb_00008eb3
Title : Crystal structure of glutamate racemase from Helicobacter pylori in complex with a fragment
Authors : Cooling, G.T.; Propp, J.; Spies, M.A.
Deposited on : 2022-08-30
Resolution : 2.20 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0
Mogul : 2022.3.0, CSD as543be (2022)
Xtrriage (Phenix) : 2.0
EDS : 3.0
Buster-report : wwPDB partial adaption of 1.1.7 (2018)
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4 : 9.0.010 (Gargrove)
Density-Fitness : 1.0.12
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION



The reported resolution of this entry is 2.20 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	6164 (2.20-2.20)
Clashscore	190562	6851 (2.20-2.20)
Ramachandran outliers	187476	6768 (2.20-2.20)
Sidechain outliers	187428	6769 (2.20-2.20)
RSRZ outliers	180081	6166 (2.20-2.20)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	260	
1	B	260	

2 Entry composition

There are 7 unique types of molecules in this entry. The entry contains 7922 atoms, of which 3846 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Glutamate racemase.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
			Total	C	H	N	O	S			
1	A	255	3772	1253	1858	306	349	6	0	0	0
1	B	255	3870	1276	1923	308	356	7	1	0	0

There are 18 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
A	-4	PRO	-	expression tag	UNP K2K6A3
A	-3	ARG	-	expression tag	UNP K2K6A3
A	-2	GLY	-	expression tag	UNP K2K6A3
A	-1	SER	-	expression tag	UNP K2K6A3
A	0	HIS	-	expression tag	UNP K2K6A3
A	24	ILE	LEU	conflict	UNP K2K6A3
A	27	LYS	GLU	conflict	UNP K2K6A3
A	62	LYS	GLN	conflict	UNP K2K6A3
A	84	HIS	TYR	conflict	UNP K2K6A3
B	-4	PRO	-	expression tag	UNP K2K6A3
B	-3	ARG	-	expression tag	UNP K2K6A3
B	-2	GLY	-	expression tag	UNP K2K6A3
B	-1	SER	-	expression tag	UNP K2K6A3
B	0	HIS	-	expression tag	UNP K2K6A3
B	24	ILE	LEU	conflict	UNP K2K6A3
B	27	LYS	GLU	conflict	UNP K2K6A3
B	62	LYS	GLN	conflict	UNP K2K6A3
B	84	HIS	TYR	conflict	UNP K2K6A3

- Molecule 2 is GLYCEROL (CCD ID: GOL) (formula: C₃H₈O₃).




Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
2	A	1	Total	C	H	O	0	0
			14	3	8	3		
2	B	1	Total	C	H	O	0	0
			13	3	7	3		

- Molecule 3 is DIMETHYL SULFOXIDE (CCD ID: DMS) (formula: C_2H_6OS).




Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
3	A	1	Total	C	H	O	S	0	0
			10	2	6	1	1		
3	A	1	Total	C	H	O	S	0	0
			10	2	6	1	1		

- Molecule 4 is 1-[4-methyl-2-(pyridin-4-yl)-1,3-thiazol-5-yl]methanamine (CCD ID: WFI) (formula: $C_{10}H_{11}N_3S$) (labeled as "Ligand of Interest" by depositor).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
			Total	C	H	N	S		
4	A	1	Total	C	H	N	S	0	0
			26	10	12	3	1		
4	B	1	Total	C	H	N	S	0	0
			26	10	12	3	1		

- Molecule 5 is D-GLUTAMIC ACID (CCD ID: DGL) (formula: $C_5H_9NO_4$).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	
5	A	1	Total	C	H	N	O	0	0
			17	5	7	1	4		
5	B	1	Total	C	H	N	O	0	0
			17	5	7	1	4		

- Molecule 6 is CHLORIDE ION (CCD ID: CL) (formula: Cl).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
6	B	1	Total	Cl	0	0
			1	1		


- Molecule 7 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
7	A	66	Total	O	0	0
			66	66		
7	B	80	Total	O	0	0
			80	80		


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Glutamate racemase



- Molecule 1: Glutamate racemase



4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	53.00Å 95.98Å 57.07Å 90.00° 112.97° 90.00°	Depositor
Resolution (Å)	47.99 – 2.20 47.99 – 2.20	Depositor EDS
% Data completeness (in resolution range)	97.1 (47.99-2.20) 97.1 (47.99-2.20)	Depositor EDS
R_{merge}	0.06	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	3.50 (at 2.20Å)	Xtrriage
Refinement program	PHENIX 1.20.1_4487	Depositor
R, R_{free}	0.214 , 0.256 0.214 , 0.255	Depositor DCC
R_{free} test set	1323 reflections (4.95%)	wwPDB-VP
Wilson B-factor (Å ²)	34.6	Xtrriage
Anisotropy	0.581	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.37 , 31.8	EDS
L-test for twinning ²	$\langle L \rangle = 0.45$, $\langle L^2 \rangle = 0.28$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
F_o, F_c correlation	0.94	EDS
Total number of atoms	7922	wwPDB-VP
Average B, all atoms (Å ²)	47.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 5.34% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: DMS, WFI, DGL, GOL, CL

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.52	0/1957	0.80	1/2665 (0.0%)
1	B	0.39	0/1993	0.67	0/2718
All	All	0.46	0/3950	0.74	1/5383 (0.0%)

There are no bond length outliers.

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	175	GLU	N-CA-C	-6.07	106.21	113.97

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1914	1858	1860	14	0
1	B	1947	1923	1925	16	0
2	A	6	8	8	0	0
2	B	6	7	8	1	0
3	A	8	12	12	0	0
4	A	14	12	0	0	0
4	B	14	12	0	0	0
5	A	10	7	7	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
5	B	10	7	7	0	0
6	B	1	0	0	0	0
7	A	66	0	0	0	0
7	B	80	0	0	2	0
All	All	4076	3846	3827	31	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 4.

All (31) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:113:VAL:HG22	1:B:178:ILE:HD12	1.81	0.61
2:B:301:GOL:O3	2:B:301:GOL:O1	2.17	0.60
1:B:107:LYS:HE3	1:B:132:GLN:O	2.02	0.59
1:B:168:THR:OG1	1:B:169:PRO:HD3	2.04	0.57
1:B:94:GLU:HB2	1:B:95:PRO:HD3	1.89	0.55
1:A:163:MET:HE1	1:A:190:LYS:C	2.33	0.53
1:A:75:ALA:HB2	1:A:93:ILE:HD11	1.93	0.49
1:B:168:THR:N	1:B:169:PRO:CD	2.76	0.49
1:A:67:ILE:HG23	1:A:214:ILE:HD13	1.93	0.49
1:B:172:ILE:O	1:B:174:PRO:HD3	2.12	0.48
1:A:163:MET:HE1	1:A:191:ILE:N	2.29	0.48
1:A:184:PHE:N	1:A:185:PRO:CD	2.77	0.48
1:B:100:ILE:CD1	1:B:178:ILE:HD11	2.43	0.47
1:B:100:ILE:HD11	1:B:178:ILE:HG12	1.98	0.46
1:A:94:GLU:HB2	1:A:95:PRO:HD3	1.98	0.45
1:A:55:LEU:HD13	1:A:85:SER:HB2	1.99	0.44
1:A:120:ILE:HD12	1:A:139:HIS:HB2	2.00	0.44
1:B:184:PHE:N	1:B:185:PRO:CD	2.81	0.43
1:A:3:ILE:HD12	1:A:65:LEU:HB3	2.00	0.42
1:B:177:VAL:HB	1:B:207:LEU:CD2	2.49	0.42
1:A:13:PHE:CZ	1:A:249:ALA:HB2	2.54	0.42
1:A:44:PRO:HB3	1:A:76:LEU:HD12	2.01	0.42
1:B:150:GLU:OE2	7:B:402:HOH:O	2.22	0.42
1:B:107:LYS:CE	1:B:132:GLN:O	2.68	0.42
1:B:219:GLN:HG3	1:B:220:GLN:N	2.35	0.42
1:A:146:VAL:HB	1:A:147:PRO:CD	2.50	0.41
1:B:8:SER:OG	7:B:401:HOH:O	2.21	0.41
1:B:100:ILE:HD11	1:B:178:ILE:CD1	2.50	0.41
1:A:146:VAL:HB	1:A:147:PRO:HD3	2.03	0.41

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:3:ILE:HG13	1:A:25:PHE:CD1	2.56	0.40
1:B:177:VAL:HB	1:B:207:LEU:HD23	2.03	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	249/260 (96%)	242 (97%)	7 (3%)	0	100	100
1	B	253/260 (97%)	244 (96%)	9 (4%)	0	100	100
All	All	502/520 (96%)	486 (97%)	16 (3%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	193/225 (86%)	191 (99%)	2 (1%)	68	81
1	B	203/225 (90%)	202 (100%)	1 (0%)	81	90
All	All	396/450 (88%)	393 (99%)	3 (1%)	73	85

All (3) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	1	MET
1	A	86	LYS
1	B	80	GLU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (3) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	71	ASN
1	A	227	ASN
1	B	132	GLN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 9 ligands modelled in this entry, 1 is monoatomic - leaving 8 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# $ Z > 2$	Counts	RMSZ	# $ Z > 2$
2	GOL	B	301	-	5,5,5	0.81	0	5,5,5	1.06	0
4	WFI	B	302	-	14,15,15	1.42	3 (21%)	15,20,20	6.16	6 (40%)
5	DGL	A	305	-	8,9,9	1.09	1 (12%)	8,11,11	1.17	1 (12%)

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
4	WFI	A	304	-	14,15,15	1.61	5 (35%)	15,20,20	6.49	4 (26%)
3	DMS	A	303	-	3,3,3	0.68	0	3,3,3	0.58	0
5	DGL	B	303	-	8,9,9	1.07	1 (12%)	8,11,11	1.12	1 (12%)
2	GOL	A	301	-	5,5,5	0.81	0	5,5,5	1.02	0
3	DMS	A	302	-	3,3,3	0.68	0	3,3,3	0.56	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	GOL	B	301	-	-	0/4/4/4	-
4	WFI	B	302	-	-	0/4/6/6	0/2/2/2
5	DGL	A	305	-	-	2/9/9/9	-
4	WFI	A	304	-	-	0/4/6/6	0/2/2/2
5	DGL	B	303	-	-	2/9/9/9	-
2	GOL	A	301	-	-	1/4/4/4	-

All (10) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
4	A	304	WFI	C06-C07	-2.92	1.32	1.38
4	B	302	WFI	C06-C07	-2.47	1.33	1.38
4	A	304	WFI	C09-N08	-2.46	1.27	1.33
4	A	304	WFI	C02-N03	-2.26	1.33	1.38
4	B	302	WFI	C09-N08	-2.26	1.27	1.33
4	A	304	WFI	C12-S11	-2.22	1.69	1.74
4	A	304	WFI	C13-C12	2.19	1.58	1.50
5	A	305	DGL	OXT-C	-2.19	1.23	1.30
5	B	303	DGL	OXT-C	-2.11	1.23	1.30
4	B	302	WFI	C13-C12	2.00	1.57	1.50

All (12) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
4	A	304	WFI	C12-S11-C04	21.40	95.74	90.09
4	B	302	WFI	C12-S11-C04	19.20	95.16	90.09
4	B	302	WFI	C04-N03-C02	11.96	116.97	111.20
4	A	304	WFI	C04-N03-C02	10.98	116.49	111.20
4	A	304	WFI	S11-C04-N03	-5.35	106.54	115.14

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
4	B	302	WFI	S11-C04-N03	-5.18	106.80	115.14
4	B	302	WFI	C01-C02-N03	2.91	125.15	119.32
4	A	304	WFI	C05-C04-N03	2.71	127.75	123.29
5	A	305	DGL	OXT-C-O	-2.61	118.15	124.08
5	B	303	DGL	OXT-C-O	-2.47	118.48	124.08
4	B	302	WFI	C05-C04-N03	2.15	126.83	123.29
4	B	302	WFI	C05-C04-S11	2.11	126.36	122.09

There are no chirality outliers.

All (5) torsion outliers are listed below:



Mol	Chain	Res	Type	Atoms
5	A	305	DGL	OE1-CD-CG-CB
5	A	305	DGL	OE2-CD-CG-CB
5	B	303	DGL	OE2-CD-CG-CB
5	B	303	DGL	OE1-CD-CG-CB
2	A	301	GOL	O1-C1-C2-O2

There are no ring outliers.

1 monomer is involved in 1 short contact:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	B	301	GOL	1	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.



5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	255/260 (98%)	0.74	23 (9%) 15 12	29, 47, 72, 92	0
1	B	255/260 (98%)	0.58	12 (4%) 36 33	30, 44, 70, 91	0
All	All	510/520 (98%)	0.66	35 (6%) 23 20	29, 45, 71, 92	0

All (35) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	-4	PRO	4.2
1	A	195	PHE	4.0
1	B	0	HIS	4.0
1	A	109	ALA	3.6
1	B	170	LEU	3.2
1	A	205	PRO	3.2
1	A	204	PRO	3.1
1	B	199	PHE	3.0
1	A	131	GLN	2.9
1	A	108	ASN	2.9
1	B	229	CYS	2.8
1	A	176	VAL	2.6
1	A	199	PHE	2.6
1	A	105	LYS	2.5
1	B	207	LEU	2.5
1	A	230	THR	2.5
1	A	175	GLU	2.4
1	A	173	LEU	2.4
1	B	126	ASP	2.4
1	A	137	VAL	2.4
1	A	111	ILE	2.3
1	B	195	PHE	2.3
1	A	135	LEU	2.3
1	A	172	ILE	2.3

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	B	172	ILE	2.2
1	A	196	MET	2.2
1	A	-2	GLY	2.2
1	A	138	SER	2.1
1	B	200	ALA	2.1
1	B	225	LYS	2.1
1	B	135	LEU	2.1
1	B	201	LEU	2.1
1	A	85	SER	2.0
1	A	225	LYS	2.0
1	A	110	PRO	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.


6.4 Ligands [i](#)


In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
3	DMS	A	302	4/4	0.77	0.17	63,76,92,92	0
4	WFI	A	304	14/14	0.80	0.15	39,49,60,60	0
3	DMS	A	303	4/4	0.84	0.17	47,65,65,67	0
2	GOL	B	301	6/6	0.84	0.18	44,51,55,61	0
4	WFI	B	302	14/14	0.84	0.13	40,51,63,63	0
2	GOL	A	301	6/6	0.87	0.12	43,51,58,63	0
5	DGL	B	303	10/10	0.91	0.08	26,30,37,37	0
6	CL	B	304	1/1	0.91	0.11	53,53,53,53	0
5	DGL	A	305	10/10	0.92	0.09	28,32,38,38	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers

as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.





6.5 Other polymers [i](#)

There are no such residues in this entry.