



# wwPDB X-ray Structure Validation Summary Report ⓘ

Mar 5, 2026 – 04:37 PM UTC

PDB ID : 1ECY / pdb\_00001ecy  
Title : PROTEASE INHIBITOR ECOTIN  
Authors : Shin, D.H.; Suh, S.W.  
Deposited on : 1996-08-06  
Resolution : 2.19 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0  
Mogul : 2022.3.0, CSD as543be (2022)  
Xtriage (Phenix) : 2.0  
EDS : 3.0  
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)  
CCP4 : 9.0.010 (Gargrove)  
Density-Fitness : 1.0.12  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.49


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*








The reported resolution of this entry is 2.19 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.








Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
Clashscore	190562	6851 (2.20-2.20)
Ramachandran outliers	187476	6768 (2.20-2.20)
Sidechain outliers	187428	6769 (2.20-2.20)
RSRZ outliers	180081	6166 (2.20-2.20)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	142	 51% 37% 11% .
2	B	2	 50% 50%
2	C	2	 50% 50%
2	D	2	 50% 50%
2	F	2	 100%
2	G	2	 100%
2	K	2	 100%

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Length	Quality of chain
2	L	2	 50% 50%
2	N	2	 100%
3	O	2	 50% 50%
3	P	2	 100%
3	Q	2	 100%
3	R	2	 50% 50%
3	S	2	 50% 50%

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
2	GLC	D	1	-	-	X	-
3	GLC	O	2	X	-	-	-
3	GLC	P	2	X	-	-	-
3	GLC	Q	2	X	-	-	-
3	GLC	R	1	X	-	-	-
3	GLC	S	2	X	-	X	-
4	GLC	A	329	-	-	X	-

## 2 Entry composition [i](#)


There are 6 unique types of molecules in this entry. The entry contains 1569 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called ECOTIN.


Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	142	1130	721	187	216	6	0	0	0

- Molecule 2 is an oligosaccharide called alpha-D-glucopyranose-(1-1)-alpha-D-glucopyranose.




Mol	Chain	Residues	Atoms			ZeroOcc	AltConf	Trace
2	B	2	Total	C	O	0	0	0
			23	12	11			
2	C	2	Total	C	O	0	0	0
			23	12	11			
2	D	2	Total	C	O	0	0	0
			23	12	11			
2	F	2	Total	C	O	0	0	0
			23	12	11			
2	G	2	Total	C	O	0	0	0
			23	12	11			
2	K	2	Total	C	O	0	0	0
			23	12	11			
2	L	2	Total	C	O	0	0	0
			23	12	11			
2	N	2	Total	C	O	0	0	0
			23	12	11			

- Molecule 3 is an oligosaccharide called alpha-D-glucopyranose-(1-1)-alpha-D-glucopyranose.




Mol	Chain	Residues	Atoms			ZeroOcc	AltConf	Trace
3	O	2	Total	C	O	0	0	0
			23	12	11			
3	P	2	Total	C	O	0	0	0
			23	12	11			
3	Q	2	Total	C	O	0	0	0
			23	12	11			
3	R	2	Total	C	O	0	0	0
			23	12	11			
3	S	2	Total	C	O	0	0	0
			23	12	11			

- Molecule 4 is alpha-D-glucopyranose (CCD ID: GLC) (formula: C<sub>6</sub>H<sub>12</sub>O<sub>6</sub>).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
4	A	1	Total	C	O	0	0
			12	6	6		
4	A	1	Total	C	O	0	0
			12	6	6		
4	A	1	Total	C	O	0	0
			12	6	6		
4	A	1	Total	C	O	0	0
			12	6	6		

- Molecule 5 is beta-D-glucopyranose (CCD ID: BGC) (formula: C<sub>6</sub>H<sub>12</sub>O<sub>6</sub>).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
5	A	1	Total C O 12 6 6	0	0


- Molecule 6 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
6	A	80	Total O 80 80	0	0

### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: ECOTIN



- Molecule 2: alpha-D-glucopyranose-(1-1)-alpha-D-glucopyranose




- Molecule 2: alpha-D-glucopyranose-(1-1)-alpha-D-glucopyranose




- Molecule 2: alpha-D-glucopyranose-(1-1)-alpha-D-glucopyranose



- Molecule 2: alpha-D-glucopyranose-(1-1)-alpha-D-glucopyranose



- Molecule 2: alpha-D-glucopyranose-(1-1)-alpha-D-glucopyranose

Chain G:  100%

GLC1  
GLC2

- Molecule 2: alpha-D-glucopyranose-(1-1)-alpha-D-glucopyranose

Chain K:  100%


GLC1  
GLC2

- Molecule 2: alpha-D-glucopyranose-(1-1)-alpha-D-glucopyranose

Chain L:  50% 50%

GLC1  
GLC2

- Molecule 2: alpha-D-glucopyranose-(1-1)-alpha-D-glucopyranose

Chain N:  100%


GLC1  
GLC2

- Molecule 3: alpha-D-glucopyranose-(1-1)-alpha-D-glucopyranose

Chain O:  50% 50%


GLC1  
GLC2

- Molecule 3: alpha-D-glucopyranose-(1-1)-alpha-D-glucopyranose

Chain P:  100%

GLC1  
GLC2

- Molecule 3: alpha-D-glucopyranose-(1-1)-alpha-D-glucopyranose

Chain Q:  100%

GLC1  
GLC2

- Molecule 3: alpha-D-glucopyranose-(1-1)-alpha-D-glucopyranose

Chain R:  50% 50%



GLC1  
GLC2

- Molecule 3: alpha-D-glucopyranose-(1-1)-alpha-D-glucopyranose

Chain S:  50% 50%



GLC1  
GLC2

## 4 Data and refinement statistics

Property	Value	Source
Space group	P 43 21 2	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	63.32Å 63.32Å 84.13Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	6.00 – 2.19 6.00 – 2.19	Depositor EDS
% Data completeness (in resolution range)	92.3 (6.00-2.19) 86.3 (6.00-2.19)	Depositor EDS
$R_{merge}$	0.07	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	1.34 (at 2.20Å)	Xtrriage
Refinement program	X-PLOR 3.1	Depositor
R, $R_{free}$	0.213 , 0.319 0.253 , (Not available)	Depositor DCC
$R_{free}$ test set	No test flags present.	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	35.6	Xtrriage
Anisotropy	0.489	Xtrriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.89 , 304.9	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.49$ , $\langle L^2 \rangle = 0.33$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
$F_o, F_c$ correlation	0.95	EDS
Total number of atoms	1569	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	41.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 7.07% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: BGC, GLC

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	A	0.57	0/1152	1.17	10/1558 (0.6%)

There are no bond length outliers.

The worst 5 of 10 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	10	ILE	N-CA-C	-8.86	104.10	112.96
1	A	119	THR	CA-C-N	-7.38	113.08	120.31
1	A	119	THR	C-N-CA	-7.38	113.08	120.31
1	A	24	VAL	N-CA-C	6.19	116.78	108.11
1	A	36	LEU	N-CA-C	6.12	118.39	108.41

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1130	0	1144	59	0
2	B	23	0	21	1	0
2	C	23	0	21	0	0
2	D	23	0	21	6	0
2	F	23	0	21	5	0
2	G	23	0	21	0	0

*Continued on next page...*

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
2	K	23	0	21	4	0
2	L	23	0	21	1	0
2	N	23	0	21	0	0
3	O	23	0	21	1	0
3	P	23	0	21	2	0
3	Q	23	0	21	1	0
3	R	23	0	21	4	0
3	S	23	0	21	6	0
4	A	48	0	48	8	0
5	A	12	0	12	1	0
6	A	80	0	0	7	0
All	All	1569	0	1477	79	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 27.

The worst 5 of 79 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
3:R:2:GLC:H61	3:R:2:GLC:H2	1.31	1.10
3:S:1:GLC:H3	3:S:2:GLC:H1	1.36	1.07
5:A:331:BGC:O1	5:A:331:BGC:H6C1	1.70	0.92
1:A:81:VAL:HA	2:D:1:GLC:H2	1.62	0.79
1:A:31:GLU:HG3	4:A:329:GLC:H3	1.69	0.74

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	140/142 (99%)	115 (82%)	21 (15%)	4 (3%)	<b>3</b> <b>2</b>

All (4) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	86	ALA
1	A	17	GLU
1	A	80	PRO
1	A	29	PRO

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	125/125 (100%)	101 (81%)	24 (19%)	<b>1</b> <b>1</b>

5 of 24 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	91	LYS
1	A	108	ARG
1	A	103	ASP
1	A	113	LEU
1	A	41	LEU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (3) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	26	GLN
1	A	44	GLN
1	A	61	ASN

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates [i](#)

26 monosaccharides are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z  > 2$	Counts	RMSZ	$\# Z  > 2$
2	GLC	B	1	2	12,12,12	0.46	0	17,17,17	0.67	0
2	GLC	B	2	2	11,11,12	0.59	0	15,15,17	0.95	1 (6%)
2	GLC	C	1	2	12,12,12	0.35	0	17,17,17	0.44	0
2	GLC	C	2	2	11,11,12	0.54	0	15,15,17	0.71	1 (6%)
2	GLC	D	1	2	12,12,12	0.42	0	17,17,17	0.83	1 (5%)
2	GLC	D	2	2	11,11,12	0.52	0	15,15,17	0.51	0
2	GLC	F	1	2	12,12,12	0.28	0	17,17,17	0.55	0
2	GLC	F	2	2	11,11,12	0.50	0	15,15,17	0.67	0
2	GLC	G	1	2	12,12,12	0.75	1 (8%)	17,17,17	1.12	1 (5%)
2	GLC	G	2	2	11,11,12	0.43	0	15,15,17	0.77	1 (6%)
2	GLC	K	1	2	12,12,12	0.35	0	17,17,17	0.74	1 (5%)
2	GLC	K	2	2	11,11,12	0.84	0	15,15,17	1.81	4 (26%)
2	GLC	L	1	2	12,12,12	0.34	0	17,17,17	0.47	0
2	GLC	L	2	2	11,11,12	0.47	0	15,15,17	0.86	1 (6%)
2	GLC	N	1	2	12,12,12	0.49	0	17,17,17	0.93	2 (11%)
2	GLC	N	2	2	11,11,12	0.53	0	15,15,17	0.94	1 (6%)
3	GLC	O	1	3	11,11,12	0.45	0	15,15,17	0.75	0
3	GLC	O	2	3	12,12,12	0.24	0	17,17,17	0.40	0
3	GLC	P	1	3	11,11,12	0.77	0	15,15,17	1.45	3 (20%)
3	GLC	P	2	3	12,12,12	0.44	0	17,17,17	0.42	0
3	GLC	Q	1	3	11,11,12	0.63	0	15,15,17	1.10	1 (6%)
3	GLC	Q	2	3	12,12,12	0.45	0	17,17,17	0.33	0
3	GLC	R	1	3	11,11,12	0.45	0	15,15,17	0.52	0
3	GLC	R	2	3	12,12,12	0.38	0	17,17,17	0.48	0

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
3	GLC	S	1	3	11,11,12	0.52	0	15,15,17	1.16	2 (13%)
3	GLC	S	2	3	12,12,12	0.43	0	17,17,17	0.50	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	GLC	B	1	2	-	2/2/22/22	0/1/1/1
2	GLC	B	2	2	-	2/2/19/22	0/1/1/1
2	GLC	C	1	2	-	2/2/22/22	0/1/1/1
2	GLC	C	2	2	-	2/2/19/22	0/1/1/1
2	GLC	D	1	2	-	2/2/22/22	0/1/1/1
2	GLC	D	2	2	-	2/2/19/22	0/1/1/1
2	GLC	F	1	2	-	0/2/22/22	0/1/1/1
2	GLC	F	2	2	-	2/2/19/22	0/1/1/1
2	GLC	G	1	2	-	2/2/22/22	0/1/1/1
2	GLC	G	2	2	-	2/2/19/22	0/1/1/1
2	GLC	K	1	2	-	2/2/22/22	0/1/1/1
2	GLC	K	2	2	-	2/2/19/22	0/1/1/1
2	GLC	L	1	2	-	2/2/22/22	0/1/1/1
2	GLC	L	2	2	-	2/2/19/22	0/1/1/1
2	GLC	N	1	2	-	0/2/22/22	0/1/1/1
2	GLC	N	2	2	-	2/2/19/22	0/1/1/1
3	GLC	O	1	3	-	2/2/19/22	0/1/1/1
3	GLC	O	2	3	1/1/5/5	0/2/22/22	0/1/1/1
3	GLC	P	1	3	-	0/2/19/22	0/1/1/1
3	GLC	P	2	3	1/1/5/5	2/2/22/22	0/1/1/1
3	GLC	Q	1	3	-	2/2/19/22	0/1/1/1
3	GLC	Q	2	3	1/1/5/5	2/2/22/22	0/1/1/1
3	GLC	R	1	3	1/1/4/5	2/2/19/22	0/1/1/1
3	GLC	R	2	3	-	0/2/22/22	0/1/1/1
3	GLC	S	1	3	-	2/2/19/22	0/1/1/1
3	GLC	S	2	3	1/1/5/5	2/2/22/22	0/1/1/1

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	G	1	GLC	C1-C2	2.09	1.57	1.52

The worst 5 of 20 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	P	1	GLC	C1-C2-C3	3.93	115.36	109.64
2	K	2	GLC	C1-C2-C3	-3.46	104.61	109.64
3	S	1	GLC	C2-C3-C4	-3.28	105.09	110.86
2	K	2	GLC	C1-O5-C5	3.24	116.53	112.19
2	G	1	GLC	C1-C2-C3	2.95	116.38	110.36

All (5) chirality outliers are listed below:

Mol	Chain	Res	Type	Atom
3	O	2	GLC	C1
3	P	2	GLC	C1
3	Q	2	GLC	C1
3	R	1	GLC	C1
3	S	2	GLC	C1

5 of 42 torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
3	S	1	GLC	O5-C5-C6-O6
2	L	2	GLC	O5-C5-C6-O6
3	O	1	GLC	O5-C5-C6-O6
3	R	1	GLC	O5-C5-C6-O6
2	D	1	GLC	O5-C5-C6-O6

There are no ring outliers.

14 monomers are involved in 30 short contacts:


Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	Q	2	GLC	1	0
2	D	1	GLC	6	0
3	R	2	GLC	4	0
2	F	1	GLC	5	0
2	F	2	GLC	1	0
2	K	2	GLC	1	0
2	D	2	GLC	2	0
3	P	2	GLC	2	0
2	K	1	GLC	3	0



*Continued on next page...*


Continued from previous page...


Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	B	2	GLC	1	0
3	S	1	GLC	5	0
3	S	2	GLC	6	0
2	L	2	GLC	1	0
3	O	2	GLC	1	0


The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for oligosaccharide.

















## 5.6 Ligand geometry

5 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
5	BGC	A	331	-	12,12,12	0.33	0	17,17,17	0.36	0
4	GLC	A	328	-	12,12,12	0.41	0	17,17,17	0.49	0
4	GLC	A	327	-	12,12,12	0.43	0	17,17,17	0.85	0
4	GLC	A	329	-	12,12,12	0.21	0	17,17,17	0.46	0
4	GLC	A	330	-	12,12,12	0.35	0	17,17,17	0.65	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '2' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
5	BGC	A	331	-	-	2/2/22/22	0/1/1/1
4	GLC	A	328	-	-	2/2/22/22	0/1/1/1
4	GLC	A	327	-	-	2/2/22/22	0/1/1/1
4	GLC	A	329	-	-	2/2/22/22	0/1/1/1
4	GLC	A	330	-	-	2/2/22/22	0/1/1/1

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

5 of 10 torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
4	A	328	GLC	O5-C5-C6-O6
4	A	327	GLC	C4-C5-C6-O6
4	A	327	GLC	O5-C5-C6-O6
4	A	328	GLC	C4-C5-C6-O6
4	A	330	GLC	C4-C5-C6-O6

There are no ring outliers.

2 monomers are involved in 9 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
5	A	331	BGC	1	0
4	A	329	GLC	8	0

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled '#RSRZ > 2' contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled 'Q < 0.9' lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	142/142 (100%)	-0.76	0 <b>100</b> <b>100</b>	17, 32, 55, 75	0

There are no RSRZ outliers to report.

### 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q < 0.9' lists the number of atoms with occupancy less than 0.9.


Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å <sup>2</sup> )	Q<0.9
2	GLC	K	1	12/12	0.70	0.10	30,63,70,75	0
2	GLC	K	2	11/12	0.78	0.08	50,61,65,81	0
2	GLC	B	2	11/12	0.79	0.07	50,63,70,74	0
2	GLC	L	2	11/12	0.79	0.09	30,38,48,80	0
2	GLC	F	1	12/12	0.80	0.08	46,65,73,91	0
2	GLC	C	2	11/12	0.82	0.07	57,66,77,77	0
2	GLC	G	2	11/12	0.84	0.07	35,46,57,69	0
2	GLC	G	1	12/12	0.86	0.08	45,57,68,72	0
2	GLC	N	2	11/12	0.86	0.07	40,51,68,74	0
2	GLC	D	1	12/12	0.88	0.08	33,55,72,72	0
2	GLC	N	1	12/12	0.88	0.07	35,39,51,63	0
2	GLC	C	1	12/12	0.88	0.06	31,64,73,74	0
2	GLC	B	1	12/12	0.89	0.08	32,62,72,75	0
2	GLC	F	2	11/12	0.89	0.07	42,50,65,75	0

*Continued on next page...*

Continued from previous page...




Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors( $\text{\AA}^2$ )	Q<0.9
2	GLC	L	1	12/12	0.89	0.07	40,53,60,60	0
2	GLC	D	2	11/12	0.92	0.06	37,50,65,69	0
3	GLC	O	1	11/12	-	-	35,40,61,62	0
3	GLC	O	2	12/12	-	-	32,51,58,70	0
3	GLC	P	1	11/12	-	-	52,63,72,78	0
3	GLC	P	2	12/12	-	-	51,71,80,90	0
3	GLC	Q	1	11/12	-	-	33,49,57,65	0
3	GLC	Q	2	12/12	-	-	27,64,73,74	0
3	GLC	R	1	11/12	-	-	54,63,79,90	0
3	GLC	R	2	12/12	-	-	39,48,66,71	0
3	GLC	S	1	11/12	-	-	26,43,48,49	0
3	GLC	S	2	12/12	-	-	43,62,69,87	0

The following is a graphical depiction of the model fit to experimental electron density for oligosaccharide. Each fit is shown from different orientation to approximate a three-dimensional view.







**Electron density around Chain C:**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)

**Electron density around Chain D:**




$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)








**Electron density around Chain G:**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)




**Electron density around Chain K:**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)






**Electron density around Chain L:**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)




**Electron density around Chain N:**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)






**Electron density around Chain O:**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)




**Electron density around Chain P:**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)






**Electron density around Chain Q:**


$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)



**Electron density around Chain R:**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)





## 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q<0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å <sup>2</sup> )	Q<0.9
4	GLC	A	330	12/12	0.58	0.11	80,94,99,99	0
4	GLC	A	328	12/12	0.73	0.11	45,69,79,96	0
5	BGC	A	331	12/12	0.75	0.08	43,61,74,92	0
4	GLC	A	329	12/12	0.84	0.10	49,63,81,88	0
4	GLC	A	327	12/12	0.86	0.07	42,52,64,65	0

## 6.5 Other polymers [i](#)

There are no such residues in this entry.