



# wwPDB X-ray Structure Validation Summary Report ⓘ

Apr 27, 2026 – 05:14 PM EDT

PDB ID : 6EDB / pdb\_00006edb  
Title : Crystal structure of SRY.hcGAS-21bp dsDNA complex  
Authors : Xie, W.; Lama, L.; Adura, C.; Glickman, J.F.; Tuschl, T.; Patel, D.J.  
Deposited on : 2018-08-09  
Resolution : 3.21 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0  
Xtriage (Phenix) : 2.0  
EDS : 3.0  
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)  
CCP4 : 9.0.010 (Gargrove)  
Density-Fitness : 1.0.12  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.49


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 3.21 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	180053	1466 (3.20-3.20)
Clashscore	190562	1573 (3.20-3.20)
Ramachandran outliers	187476	1548 (3.20-3.20)
Sidechain outliers	187428	1547 (3.20-3.20)
RSRZ outliers	180081	1466 (3.20-3.20)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	455	
1	B	455	
2	C	22	
3	D	22	

## 2 Entry composition [i](#)

There are 4 unique types of molecules in this entry. The entry contains 4379 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Sex-determining region Y protein, Cyclic GMP-AMP synthase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	357	2944	1882	510	537	15	0	0	0
1	B	67	579	367	110	97	5	0	0	0

There are 28 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	68	SER	-	expression tag	UNP Q05066
A	144	LYS	-	linker	UNP Q05066
A	145	GLY	-	linker	UNP Q05066
A	146	GLU	-	linker	UNP Q05066
A	147	THR	-	linker	UNP Q05066
A	148	LYS	-	linker	UNP Q05066
A	149	LYS	-	linker	UNP Q05066
A	150	LYS	-	linker	UNP Q05066
A	151	PHE	-	linker	UNP Q05066
A	152	LYS	-	linker	UNP Q05066
A	153	ASP	-	linker	UNP Q05066
A	154	PRO	-	linker	UNP Q05066
A	155	ASN	-	linker	UNP Q05066
A	156	ALA	-	linker	UNP Q05066
B	0	SER	-	expression tag	UNP Q05066
B	76	LYS	-	linker	UNP Q05066
B	77	GLY	-	linker	UNP Q05066
B	78	GLU	-	linker	UNP Q05066
B	79	THR	-	linker	UNP Q05066
B	80	LYS	-	linker	UNP Q05066
B	81	LYS	-	linker	UNP Q05066
B	82	LYS	-	linker	UNP Q05066
B	83	PHE	-	linker	UNP Q05066
B	84	LYS	-	linker	UNP Q05066
B	85	ASP	-	linker	UNP Q05066

*Continued on next page...*

*Continued from previous page...*

Chain	Residue	Modelled	Actual	Comment	Reference
B	86	PRO	-	linker	UNP Q05066
B	87	ASN	-	linker	UNP Q05066
B	88	ALA	-	linker	UNP Q05066

- Molecule 2 is a DNA chain called DNA (5'-D(\*CP\*GP\*AP\*TP\*CP\*CP\*GP\*GP\*GP\*AP\*TP\*CP\*TP\*AP\*AP\*AP\*CP\*AP\*AP\*TP\*G)-3').

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	P			
2	C	21	429	205	83	121	20	0	0	0

- Molecule 3 is a DNA chain called DNA (5'-D(\*GP\*CP\*AP\*TP\*TP\*GP\*TP\*TP\*TP\*AP\*GP\*AP\*TP\*CP\*CP\*CP\*GP\*GP\*AP\*TP\*C)-3').

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	P			
3	D	21	426	205	74	127	20	0	0	0


- Molecule 4 is ZINC ION (CCD ID: ZN) (formula: Zn).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	1	Total	Zn	0	0
			1	1		



VAL THR TYR PHE LEU GLN CYS LEU ARG THR GLU LYS LEU GLU ASN TYR PHE ILE PRO GLU PHE ASN LEU PHE SER SER ASN ILE ASP LYS ARG SER LYS LEU PHE LEU THR LYS GLN ILE GLU TYR GLU ARG ASN ASN GLU PHE PRO VAL PHE ASP GLU PHE

- Molecule 2: DNA (5'-D(\*CP\*GP\*AP\*TP\*CP\*CP\*GP\*GP\*GP\*AP\*TP\*CP\*TP\*AP\*AP\*AP\*CP\*AP\*AP\*TP\*G)-3')

Chain C:  18% 77% 5%

C1 G2 A3 T4 C5 C6 C7 C8 C9 A10 T11 C12 T13 A14 A15 A16 C17 A18 A19 T20 G21 DC

- Molecule 3: DNA (5'-D(\*GP\*CP\*AP\*TP\*TP\*GP\*TP\*TP\*TP\*AP\*GP\*AP\*TP\*CP\*CP\*CP\*GP\*GP\*AP\*TP\*C)-3')

Chain D:  55% 41% 5%

D6 G2 C3 T6 T10 A11 G12 C17 G18 G19 C22

## 4 Data and refinement statistics

Property	Value	Source
Space group	I 2 2 2	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	90.31Å 115.83Å 159.08Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	30.01 – 3.21 30.01 – 3.21	Depositor EDS
% Data completeness (in resolution range)	90.3 (30.01-3.21) 82.3 (30.01-3.21)	Depositor EDS
$R_{merge}$	(Not available)	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	1.83 (at 3.24Å)	Xtrriage
Refinement program	PHENIX (1.10.1_2155: ???)	Depositor
R, $R_{free}$	0.255 , 0.302 0.260 , 0.300	Depositor DCC
$R_{free}$ test set	1264 reflections (10.00%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	120.7	Xtrriage
Anisotropy	0.498	Xtrriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.27 , 35.1	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.48$ , $\langle L^2 \rangle = 0.31$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
$F_o, F_c$ correlation	0.94	EDS
Total number of atoms	4379	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	105.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 5.12% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section:  
ZN

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	A	0.33	0/3003	0.57	7/4027 (0.2%)
1	B	0.12	0/594	0.27	0/794
2	C	0.14	0/482	0.31	0/742
3	D	0.19	0/476	0.40	0/733
All	All	0.28	0/4555	0.50	7/6296 (0.1%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	1

There are no bond length outliers.

The worst 5 of 7 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	257	PRO	N-CA-C	8.30	125.07	113.53
1	A	335	GLN	N-CA-C	6.66	118.54	111.28
1	A	331	PRO	CA-C-O	6.48	130.71	122.15
1	A	331	PRO	N-CA-C	5.53	119.36	111.13
1	A	256	ASN	N-CA-C	5.51	121.98	109.81

There are no chirality outliers.

All (1) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	330	TRP	Mainchain

## 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2944	0	2990	91	0
1	B	579	0	575	12	0
2	C	429	0	237	14	0
3	D	426	0	240	7	0
4	A	1	0	0	0	0
All	All	4379	0	4042	118	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 14.

The worst 5 of 118 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:444:HIS:O	1:A:447:THR:HG22	1.40	1.18
1:A:430:LEU:HD21	1:A:504:LEU:HD13	1.40	1.03
1:A:241:GLU:OE1	1:A:246:ARG:HD2	1.68	0.92
1:A:416:LEU:O	1:A:420:LEU:HD13	1.72	0.88
1:A:220:ILE:HD11	1:A:418:GLU:OE2	1.79	0.82

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	355/455 (78%)	324 (91%)	30 (8%)	1 (0%)	36 68

*Continued on next page...*

Continued from previous page...

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	B	65/455 (14%)	64 (98%)	1 (2%)	0	100	100
All	All	420/910 (46%)	388 (92%)	31 (7%)	1 (0%)	43	73

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	303	GLY

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	331/417 (79%)	325 (98%)	6 (2%)	51	74
1	B	60/417 (14%)	58 (97%)	2 (3%)	33	65
All	All	391/834 (47%)	383 (98%)	8 (2%)	48	72

5 of 8 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	B	57	GLU
1	B	44	LYS
1	A	476	ARG
1	A	410	LEU
1	A	483	TYR

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (2) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	514	ASN
1	B	27	ASN

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

### 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

### 5.6 Ligand geometry [i](#)

Of 1 ligands modelled in this entry, 1 is monoatomic - leaving 0 for Mogul analysis.

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

### 5.7 Other polymers [i](#)

There are no such residues in this entry.

### 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	357/455 (78%)	-0.41	3 (0%) 82 67	63, 100, 137, 148	0
1	B	67/455 (14%)	-0.48	0 100 100	90, 116, 135, 157	0
2	C	21/22 (95%)	-0.76	0 100 100	87, 101, 123, 127	0
3	D	21/22 (95%)	-0.67	0 100 100	87, 102, 121, 141	0
All	All	466/954 (48%)	-0.44	3 (0%) 85 73	63, 103, 137, 157	0

All (3) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	276	MET	3.5
1	A	201	SER	2.1
1	A	262	LEU	2.0

### 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

### 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors( $\text{\AA}^2$ )	Q<0.9
4	ZN	A	601	1/1	0.97	0.09	30,30,30,30	0

## 6.5 Other polymers [i](#)

There are no such residues in this entry.