



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 9, 2026 – 01:38 PM UTC

PDB ID : 6EDI / pdb_00006edi
Title : Crystal structure of Leishmania braziliensis glucokinase
Authors : Buechner, G.S.; Millington, M.E.; Perry, K.; D'Antonio, E.L.
Deposited on : 2018-08-09
Resolution : 1.85 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.85 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	3428 (1.86-1.86)
Clashscore	190562	3579 (1.86-1.86)
Ramachandran outliers	187476	3553 (1.86-1.86)
Sidechain outliers	187428	3553 (1.86-1.86)
RSRZ outliers	180081	3429 (1.86-1.86)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	412	4% 77% 16% • •
1	B	412	3% 78% 16% • •

2 Entry composition

There are 2 unique types of molecules in this entry. The entry contains 6126 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Glucokinase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	394	Total	C	N	O	S	0	0	0
			2975	1890	512	559	14			
1	B	394	Total	C	N	O	S	0	0	0
			2975	1890	512	559	14			

There are 28 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	-13	MET	-	expression tag	UNP A4HPA0
A	-12	GLY	-	expression tag	UNP A4HPA0
A	-11	ARG	-	expression tag	UNP A4HPA0
A	-10	GLY	-	expression tag	UNP A4HPA0
A	-9	SER	-	expression tag	UNP A4HPA0
A	-8	HIS	-	expression tag	UNP A4HPA0
A	-7	HIS	-	expression tag	UNP A4HPA0
A	-6	HIS	-	expression tag	UNP A4HPA0
A	-5	HIS	-	expression tag	UNP A4HPA0
A	-4	HIS	-	expression tag	UNP A4HPA0
A	-3	HIS	-	expression tag	UNP A4HPA0
A	-2	GLY	-	expression tag	UNP A4HPA0
A	-1	MET	-	expression tag	UNP A4HPA0
A	0	ALA	-	expression tag	UNP A4HPA0
B	-13	MET	-	expression tag	UNP A4HPA0
B	-12	GLY	-	expression tag	UNP A4HPA0
B	-11	ARG	-	expression tag	UNP A4HPA0
B	-10	GLY	-	expression tag	UNP A4HPA0
B	-9	SER	-	expression tag	UNP A4HPA0
B	-8	HIS	-	expression tag	UNP A4HPA0
B	-7	HIS	-	expression tag	UNP A4HPA0
B	-6	HIS	-	expression tag	UNP A4HPA0
B	-5	HIS	-	expression tag	UNP A4HPA0
B	-4	HIS	-	expression tag	UNP A4HPA0
B	-3	HIS	-	expression tag	UNP A4HPA0

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
B	-2	GLY	-	expression tag	UNP A4HPA0
B	-1	MET	-	expression tag	UNP A4HPA0
B	0	ALA	-	expression tag	UNP A4HPA0


- Molecule 2 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	A	91	Total	O	0	0
			91	91		
2	B	85	Total	O	0	0
			85	85		


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: Glucokinase



• Molecule 1: Glucokinase



4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	68.48Å 63.26Å 85.57Å 90.00° 107.81° 90.00°	Depositor
Resolution (Å)	49.97 – 1.85 49.97 – 1.85	Depositor EDS
% Data completeness (in resolution range)	97.6 (49.97-1.85) 97.6 (49.97-1.85)	Depositor EDS
R_{merge}	0.10	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.22 (at 1.84Å)	Xtriage
Refinement program	PHENIX 1.8.2_1309	Depositor
R, R_{free}	0.212 , 0.262 0.215 , 0.265	Depositor DCC
R_{free} test set	2968 reflections (4.93%)	wwPDB-VP
Wilson B-factor (Å ²)	31.4	Xtriage
Anisotropy	0.300	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.33 , 25.3	EDS
L-test for twinning ²	$\langle L \rangle = 0.48$, $\langle L^2 \rangle = 0.31$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.95	EDS
Total number of atoms	6126	wwPDB-VP
Average B, all atoms (Å ²)	35.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 5.53% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.47	0/3033	0.82	0/4106
1	B	0.47	0/3033	0.86	4/4106 (0.1%)
All	All	0.47	0/6066	0.84	4/8212 (0.0%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	B	0	1

There are no bond length outliers.

All (4) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed($^{\circ}$)	Ideal($^{\circ}$)
1	B	285	ALA	CA-C-N	7.68	128.08	119.32
1	B	285	ALA	C-N-CA	7.68	128.08	119.32
1	B	85	LEU	CA-C-N	6.70	128.22	119.84
1	B	85	LEU	C-N-CA	6.70	128.22	119.84

There are no chirality outliers.

All (1) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	B	113	ALA	Peptide

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen

atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2975	0	2986	45	0
1	B	2975	0	2986	41	0
2	A	91	0	0	0	0
2	B	85	0	0	3	0
All	All	6126	0	5972	83	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 7.

All (83) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:238:GLN:NE2	1:B:255:SER:O	2.05	0.89
1:B:87:ASP:OD1	1:B:87:ASP:N	2.25	0.70
1:B:117:ASN:ND2	1:B:118:ASN:OD1	2.23	0.70
1:A:276:TYR:HA	1:A:280:GLY:HA3	1.74	0.69
1:B:310:ASP:OD2	1:B:354:ARG:NH1	2.26	0.68
1:B:137:ARG:HD3	1:B:397:LEU:HD13	1.75	0.67
1:A:280:GLY:O	1:A:282:LYS:N	2.28	0.66
1:A:282:LYS:HE3	1:A:283:ALA:H	1.62	0.64
1:B:78:LEU:O	1:B:82:LYS:HG2	1.99	0.62
1:B:188:ARG:NH1	1:B:212:ASP:OD1	2.32	0.62
1:A:113:ALA:HB3	1:A:123:ALA:HB3	1.82	0.61
1:A:251:LYS:O	1:A:253:LEU:N	2.30	0.58
1:A:167:MET:SD	1:A:378:ARG:HG3	2.44	0.57
1:B:113:ALA:HB3	1:B:123:ALA:HB3	1.86	0.57
1:A:78:LEU:HD21	1:A:131:GLU:HB2	1.86	0.57
1:A:74:PHE:HB3	1:A:129:PRO:HG2	1.87	0.57
1:A:137:ARG:HH11	1:A:137:ARG:HG2	1.69	0.57
1:A:254:ASP:HB3	1:A:256:THR:H	1.70	0.57
1:B:22:TRP:CZ2	1:B:56:ILE:HG12	2.39	0.57
1:B:32:ASP:OD2	1:B:39:ARG:NH2	2.37	0.57
1:A:78:LEU:O	1:A:82:LYS:HG2	2.05	0.57
1:B:243:TYR:HD2	1:B:244:MET:HE2	1.71	0.56
1:A:254:ASP:HB2	1:A:257:VAL:HG23	1.86	0.56
1:A:90:ALA:O	1:A:94:ARG:HG2	2.06	0.55
1:A:346:PHE:CG	1:A:378:ARG:HD2	2.42	0.54
1:B:280:GLY:O	1:B:282:LYS:N	2.38	0.54
1:B:303:THR:HA	1:B:306:ILE:HD12	1.89	0.53

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:22:TRP:O	1:B:95:ARG:NH2	2.40	0.53
1:B:97:ALA:O	1:B:137:ARG:HD2	2.10	0.50
1:B:30:VAL:HG12	1:B:390:CYS:HB2	1.93	0.50
1:A:137:ARG:HG2	1:A:137:ARG:NH1	2.25	0.50
1:A:251:LYS:C	1:A:253:LEU:H	2.19	0.50
1:B:295:LYS:NZ	2:B:401:HOH:O	2.41	0.50
1:A:146:ALA:O	1:A:389:GLY:HA3	2.12	0.49
1:A:34:GLY:O	1:A:37:SER:OG	2.31	0.49
1:A:48:HIS:HB2	1:A:50:ARG:H	1.78	0.48
1:B:12:LEU:HB3	1:B:80:HIS:CD2	2.48	0.48
1:B:137:ARG:CD	1:B:397:LEU:HD13	2.41	0.48
1:A:301:SER:OG	1:A:302:ASP:N	2.44	0.47
1:B:196:PRO:HG3	1:B:316:LEU:HD22	1.96	0.47
1:A:51:SER:HB3	1:A:157:ALA:HB1	1.95	0.47
1:B:135:LYS:HB3	1:B:135:LYS:HE2	1.60	0.47
1:A:280:GLY:C	1:A:282:LYS:N	2.73	0.47
1:B:118:ASN:OD1	1:B:118:ASN:N	2.47	0.47
1:A:252:GLY:O	1:B:366:ARG:NH1	2.46	0.47
1:B:240:LEU:HD11	1:B:244:MET:HE3	1.96	0.46
1:A:197:GLY:HA2	1:A:338:ASN:OD1	2.14	0.46
1:A:78:LEU:HD22	1:A:132:LEU:HG	1.97	0.46
1:B:31:CYS:HA	1:B:39:ARG:O	2.17	0.45
1:B:360:HIS:HA	1:B:365:GLU:HG2	1.99	0.45
1:A:3:GLN:HB2	1:A:54:HIS:HD2	1.80	0.45
1:B:231:ASN:O	1:B:235:LYS:HG2	2.17	0.45
1:A:31:CYS:HA	1:A:39:ARG:O	2.17	0.44
1:A:159:ILE:HD12	1:A:159:ILE:HA	1.77	0.44
1:A:22:TRP:CZ2	1:A:56:ILE:HG12	2.53	0.44
1:A:137:ARG:HH12	1:A:397:LEU:HB3	1.83	0.44
1:B:4:THR:HG21	1:B:384:ASN:ND2	2.33	0.44
1:B:64:LYS:NZ	2:B:403:HOH:O	2.51	0.44
1:A:137:ARG:HH11	1:A:137:ARG:CG	2.32	0.43
1:B:28:TYR:HB2	1:B:43:SER:OG	2.18	0.43
1:B:278:LYS:HE3	2:B:444:HOH:O	2.18	0.43
1:A:302:ASP:O	1:A:306:ILE:HG13	2.18	0.43
1:B:285:ALA:HA	1:B:286:PRO:HD3	1.83	0.43
1:A:137:ARG:NH1	1:A:397:LEU:HB3	2.34	0.43
1:A:10:GLU:H	1:A:10:GLU:CD	2.27	0.42
1:B:241:ALA:HB2	1:B:259:PRO:HD3	2.00	0.42
1:A:366:ARG:HD3	1:B:252:GLY:O	2.19	0.42
1:B:101:VAL:HB	1:B:140:LEU:HD23	2.02	0.42

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:75:ASP:O	1:A:79:GLN:HG2	2.20	0.42
1:A:22:TRP:HZ2	1:A:56:ILE:HG12	1.84	0.42
1:B:146:ALA:O	1:B:389:GLY:HA3	2.20	0.42
1:B:78:LEU:HD12	1:B:78:LEU:HA	1.87	0.42
1:A:12:LEU:HB3	1:A:80:HIS:CD2	2.54	0.41
1:B:172:GLN:HG2	1:B:176:LEU:HD22	2.01	0.41
1:A:171:THR:OG1	1:A:373:ARG:HA	2.20	0.41
1:A:251:LYS:C	1:A:253:LEU:N	2.78	0.41
1:B:173:TRP:O	1:B:177:SER:HB3	2.21	0.41
1:A:348:LYS:HD2	1:A:351:ASN:ND2	2.36	0.41
1:A:7:ILE:CD1	1:A:12:LEU:HA	2.50	0.41
1:A:367:GLN:HG2	1:B:251:LYS:HB2	2.02	0.41
1:B:276:TYR:CE1	1:B:281:GLN:HA	2.55	0.41
1:A:53:LEU:HD13	1:A:388:LEU:HD22	2.03	0.40
1:A:253:LEU:HD22	1:A:253:LEU:HA	1.94	0.40

There are no symmetry-related clashes.

5.3 Torsion angles

5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	390/412 (95%)	371 (95%)	12 (3%)	7 (2%)	6	1
1	B	390/412 (95%)	374 (96%)	15 (4%)	1 (0%)	36	25
All	All	780/824 (95%)	745 (96%)	27 (4%)	8 (1%)	12	4

All (8) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	282	LYS
1	A	52	GLY
1	A	281	GLN

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	A	287	LEU
1	A	301	SER
1	B	51	SER
1	A	252	GLY
1	A	302	ASP

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	315/327 (96%)	295 (94%)	20 (6%)	16	4
1	B	315/327 (96%)	302 (96%)	13 (4%)	27	12
All	All	630/654 (96%)	597 (95%)	33 (5%)	21	7

All (33) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	14	LEU
1	A	36	THR
1	A	49	ASP
1	A	50	ARG
1	A	78	LEU
1	A	120	LYS
1	A	130	VAL
1	A	137	ARG
1	A	143	ASP
1	A	144	LEU
1	A	159	ILE
1	A	198	THR
1	A	251	LYS
1	A	253	LEU
1	A	284	SER
1	A	288	LYS
1	A	299	SER
1	A	303	THR

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	A	350	GLU
1	A	369	LYS
1	B	14	LEU
1	B	21	SER
1	B	37	SER
1	B	56	ILE
1	B	78	LEU
1	B	117	ASN
1	B	135	LYS
1	B	144	LEU
1	B	159	ILE
1	B	176	LEU
1	B	296	LEU
1	B	334	LEU
1	B	343	SER

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (5) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	326	GLN
1	B	117	ASN
1	B	172	GLN
1	B	238	GLN
1	B	326	GLN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no oligosaccharides in this entry.

5.6 Ligand geometry

There are no ligands in this entry.

5.7 Other polymers

There are no such residues in this entry.

5.8 Polymer linkage issues

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	394/412 (95%)	0.32	16 (4%) 41 45	21, 34, 52, 75	0
1	B	394/412 (95%)	0.24	14 (3%) 46 50	22, 33, 50, 83	0
All	All	788/824 (95%)	0.28	30 (3%) 44 48	21, 33, 52, 83	0

All (30) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	114	GLY	5.3
1	A	49	ASP	3.9
1	A	280	GLY	3.7
1	A	287	LEU	3.7
1	A	283	ALA	3.3
1	A	48	HIS	3.2
1	A	47	GLN	3.1
1	A	285	ALA	3.1
1	B	48	HIS	3.0
1	A	50	ARG	2.9
1	A	286	PRO	2.9
1	B	49	ASP	2.7
1	B	51	SER	2.7
1	B	252	GLY	2.6
1	B	50	ARG	2.6
1	B	286	PRO	2.6
1	B	34	GLY	2.6
1	B	254	ASP	2.6
1	A	284	SER	2.6
1	B	46	SER	2.6
1	A	51	SER	2.5
1	A	253	LEU	2.3
1	A	246	SER	2.2
1	B	47	GLN	2.2

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	B	20	ALA	2.2
1	A	254	ASP	2.1
1	B	45	ALA	2.1
1	B	159	ILE	2.0
1	A	281	GLN	2.0
1	A	251	LYS	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

There are no ligands in this entry.

6.5 Other polymers [i](#)

There are no such residues in this entry.